

**PCT**WORLD INTELLECTUAL  
PROPERTY ORGANIZATION  
International BureauExp Mail EV335610938US  
USAN 09/895,814

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/67, 9/64, 15/63, C07K 16/40,</b> <b>C12Q 1/68, G01N 33/53, A61K 39/395,</b> <b>48/00, 38/48, A01K 67/027</b>		<b>A2</b>	<b>(11) International Publication Number: WO 00/53776</b> <b>(43) International Publication Date: 14 September 2000 (14.09.00)</b>
<b>(21) International Application Number: PCT/CA00/00258</b> <b>(22) International Filing Date: 9 March 2000 (09.03.00)</b>  <b>(30) Priority Data:</b> 60/124,260 11 March 1999 (11.03.99) US 60/127,386 1 April 1999 (01.04.99) US 60/144,919 21 July 1999 (21.07.99) US  <b>(71) Applicant (for all designated States except US):</b> MOUNT SINAI HOSPITAL [CA/CA]; Samuel Lunenfeld Research Institute, Office Of Technology Transfer & Industrial Liaison, 600 University Avenue, Toronto, Ontario M5G 1X5 (CA).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> YOUSEF, George, M. [EG/CA]; Suite 1701, 50 Stephanie St., Toronto, Ontario M5T 1B3 (CA). DIAMANDIS, Eleftherios, P. [CA/CA]; Suite 44, 1504 Gerrard St. W., Toronto, Ontario M5G 2X2 (CA).  <b>(74) Agents:</b> VAN ZANT, Joan M., et al.; Swabey Ogilvy Renault, 77 Bloor Street West, Suite 1407, Toronto, Ontario M5S 1M2 (CA).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title: NOVEL HUMAN KALLIKREIN-LIKE GENES</b>			
<b>(57) Abstract</b>  The invention relates to nucleic acid molecules, kallikrein-like proteins encoded by such nucleic acid molecules; and use of the proteins and nucleic acid molecules			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**TITLE: Novel Human Kallikrein-Like Genes****FIELD OF THE INVENTION**

5 The invention relates to nucleic acid molecules, proteins encoded by such nucleic acid molecules; and use of the proteins and nucleic acid molecules

**BACKGROUND OF THE INVENTION**

10 Kallikreins and kallikrein-like proteins are a subgroup of the serine protease enzyme family and exhibit a high degree of substrate specificity (1). The biological role of these kallikreins is the selective cleavage of specific polypeptide precursors (substrates) to release peptides with potent biological activity (2). In mouse and rat, kallikreins are encoded by large multigene families. In the mouse genome, at least 24 genes have been identified (3). Expression of 11 of these genes has been confirmed; the rest are presumed to be pseudogenes (4). A similar family of 15-20 kallikreins has been found in the rat genome (5) where at least 4 of these are known to be expressed (6).

15 Three human kallikrein genes have been described, i.e. prostatic specific antigen (PSA or KLK3) (7), human glandular kallikrein (KLK2) (8) and tissue (pancreatic-renal) kallikrein (KLK1) (9). The PSA gene spans 5.8 Kb of sequence which has been published (7); the KLK2 gene has a size of 5.2 Kb and its complete structure has also been elucidated (8). The KLK1 gene is approximately 4.5 Kb long and the exon sequences and the exon/intron junctions of this gene have been determined (9).

20 The mouse kallikrein genes are clustered in groups of up to 11 genes on chromosome 7 and the distance between the genes in the various clusters can be as small as 3-7 Kb (3). All three human kallikrein genes have been assigned to chromosome 19q13.2 – 19q13.4 and the distance between PSA and KLK2 has been estimated to be 12 Kb (9).

25 A major difference between mouse and human kallikreins is that two of the human kallikreins (KLK2 and KLK3) are expressed almost exclusively in the prostate while in animals none of the kallikreins is localized in this organ. Other candidate new members of the human kallikrein gene family include protease M (10) (also named Zyme (11) or neurosin (12) and the normal epithelial cell-specific gene-1 (NES1) (13). Both genes have been assigned to chromosome 19q13.3 (10,14) and show structural homology with other serine proteases and the kallikrein gene family (10-14).

**SUMMARY OF THE INVENTION**

30 In efforts to precisely define the relative genomic location of PSA, KLK2, Zyme and NES1 genes, an area spanning approximately 300 Kb of contiguous sequence on human chromosome 19 (19q13.3 –q13.4) was examined. The present inventors were able to identify the relative location of the known kallikrein genes and, in addition, they identified other kallikrein-like genes which exhibit both location proximity and structural similarity with the known members of the human kallikrein family. The novel  
35 genes exhibit homology with the currently known members of the kallikrein family and they are co-localized in the same genomic region. These new genes, like the already known kallikreins have utility in various cancers including those of the breast, testicular, and prostate.

The kallikrein-like proteins described herein are individually referred to as "KLK-L1, KLK-L2,

KLK-L3, KLK-L4, KLK-L5, or KLK-L6", and collectively as "kallikrein-like proteins" or "KLK-L Proteins". The genes encoding the proteins are referred to as "*klk-11*, *klk-12*, *klk-13*, *klk-14*, *klk-15*, or *klk-16*", and collectively as "kallikrein-like genes" or "*klk-l* genes".

Broadly stated the present invention relates to an isolated nucleic acid molecule which comprises:

- 5 (i) a nucleic acid sequence encoding a protein having substantial sequence identity with an amino acid sequence of KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively;
- (ii) a nucleic acid sequence encoding a protein comprising an amino acid sequence of KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 as shown in SEQ.ID.NO. 2, 3, 10 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively;
- (iii) nucleic acid sequences complementary to (i);
- (iv) a degenerate form of a nucleic acid sequence of (i);
- (v) a nucleic acid sequence capable of hybridizing under stringent conditions to a nucleic acid sequence in (i), (ii) or (iii);
- 15 (vi) a nucleic acid sequence encoding a truncation, an analog, an allelic or species variation of a protein comprising an amino acid sequence of KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively; or
- (vii) a fragment, or allelic or species variation of (i), (ii) or (iii).

20 Preferably, a purified and isolated nucleic acid molecule of the invention comprises:

- (i) a nucleic acid sequence comprising the sequence of SEQ.ID.NO. 1, 13, 21, 43, 56, or 65 wherein T can also be U;
- (ii) nucleic acid sequences complementary to (i), preferably complementary to the full nucleic acid sequence of SEQ.ID.NO. 1, 13, 21, 43, 56, or 65;
- 25 (iii) a nucleic acid capable of hybridizing under stringent conditions to a nucleic acid of (i) or (ii) and preferably having at least 18 nucleotides; or
- (iv) a nucleic acid molecule differing from any of the nucleic acids of (i) to (iii) in codon sequences due to the degeneracy of the genetic code.

The invention also contemplates a nucleic acid molecule comprising a sequence encoding a truncation of a KLK-L protein, an analog, or a homolog of a KLK-L Protein or a truncation thereof. (KLK-L Proteins and truncations, analogs and homologs of KLK-L Proteins are also collectively referred to herein as "KLK-L Related Proteins").

The nucleic acid molecules of the invention may be inserted into an appropriate expression vector, i.e. a vector that contains the necessary elements for the transcription and translation of the inserted coding sequence. Accordingly, recombinant expression vectors adapted for transformation of a host cell may be constructed which comprise a nucleic acid molecule of the invention and one or more transcription and translation elements linked to the nucleic acid molecule.

The recombinant expression vector can be used to prepare transformed host cells expressing KLK-

L Related Proteins. Therefore, the invention further provides host cells containing a recombinant molecule of the invention. The invention also contemplates transgenic non-human mammals whose germ cells and somatic cells contain a recombinant molecule comprising a nucleic acid molecule of the invention, in particular one which encodes an analog of a KLK-L Protein, or a truncation of a KLK-L Protein.

5           The invention further provides a method for preparing KLK-L Related Proteins utilizing the purified and isolated nucleic acid molecules of the invention. In an embodiment a method for preparing a KLK-L Related Protein is provided comprising (a) transferring a recombinant expression vector of the invention into a host cell; (b) selecting transformed host cells from untransformed host cells; (c) culturing a selected transformed host cell under conditions which allow expression of the KLK-L Related Protein;  
10       and (d) isolating the KLK-L Related Protein.

          The invention further broadly contemplates an isolated KLK-L Protein comprising an amino acid sequence as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67.

          The KLK-L Related Proteins of the invention may be conjugated with other molecules, such as proteins, to prepare fusion proteins. This may be accomplished, for example, by the synthesis of N-terminal  
15       or C-terminal fusion proteins.

          The invention further contemplates antibodies having specificity against an epitope of a KLK-L Related Protein of the invention. Antibodies may be labeled with a detectable substance and used to detect proteins of the invention in tissues and cells.

          The invention also permits the construction of nucleotide probes which are unique to the nucleic acid molecules of the invention and/or to proteins of the invention. Therefore, the invention also relates to  
20       a probe comprising a nucleic acid sequence of the invention, or a nucleic acid sequence encoding a protein of the invention, or a part thereof. The probe may be labeled, for example, with a detectable substance and it may be used to select from a mixture of nucleotide sequences a nucleic acid molecule of the invention including nucleic acid molecules coding for a protein which displays one or more of the properties of a  
25       protein of the invention.

          The invention still further provides a method for identifying a substance which binds to a protein of the invention comprising reacting the protein with at least one substance which potentially can bind with the protein, under conditions which permit the formation of complexes between the substance and protein and detecting binding. Binding may be detected by assaying for complexes, for free substance, or for non-  
30       complexed protein. The invention also contemplates methods for identifying substances that bind to other intracellular proteins that interact with a KLK-L Related Protein. Methods can also be utilized which identify compounds which bind to KLK-L gene regulatory sequences (e.g. promoter sequences).

          Still further the invention provides a method for evaluating a compound for its ability to modulate the biological activity of a KLK-L Related Protein of the invention. For example a substance which inhibits  
35       or enhances the interaction of the protein and a substance which binds to the protein may be evaluated. In an embodiment, the method comprises providing a known concentration of a KLK-L Related Protein, with a substance which binds to the protein and a test compound under conditions which permit the formation of complexes between the substance and protein, and removing and/or detecting complexes.

Compounds which modulate the biological activity of a protein of the invention may also be identified using the methods of the invention by comparing the pattern and level of expression of the protein of the invention in tissues and cells, in the presence, and in the absence of the compounds.

5 The proteins of the invention and substances and compounds identified using the methods of the invention, and peptides of the invention may be used to modulate the biological activity of a KLK-L Related Protein of the invention, and they may be used in the treatment of conditions such as cancer (e.g. breast, testicular, and prostate cancer). Accordingly, the substances and compounds may be formulated into compositions for administration to individuals suffering from cancer.

10 Therefore, the present invention also relates to a composition comprising one or more of a protein of the invention, a peptide of the invention, or a substance or compound identified using the methods of the invention, and a pharmaceutically acceptable carrier, excipient or diluent. A method for treating or preventing cancer is also provided comprising administering to a patient in need thereof, a KLK-L Related Protein of the invention, or a composition of the invention.

15 Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

20 The invention will now be described in relation to the drawings in which:

Figure 1 shows an approximate 300 Kb of contiguous genomic sequence around chromosome 19q13.3 - q13.4 represented by 8 contigs, each one shown with its length in Kb. The contig numbers refer to those reported in the Lawrence Livermore National Laboratory website. Note the localization of the seven known genes (PSA, KLK2, Zyme, NES1, HSCCE, neuropsin and TLSP) (see abbreviations for full names of these genes). All genes are represented with arrows denoting the direction of transcription. The gene with no homology to human kallikreins is termed UG (unknown gene). The five new kallikrein-like genes (KLK-L1 to KLK-L5) were numbered from the most centromeric to the most telomeric. Numbers just below or just above the arrows indicate appropriate Kb lengths in each contig. Gene lengths and distances between genes are rounded to the nearest 6.5 kb. The site of the gap is marked with an asterisk.

25 30 Figure 2 shows a contiguous genomic sequence around chromosome 19q13.3- q13.4. Genes are represented by horizontal arrows denoting the direction of the coding sequence. Distances between genes are in base pairs.

Figure 3 shows tissue expression of the prostate/KLK-L1 gene as determined by RT-PCR. Actin and PSA are control genes. Interpretations are presented in Table 9.

35 Figure 4 shows the sequence of PCR product obtained with cDNA from female breast tissue using prostate/KLK-L1 primers. Primer sequences are underlined. The sequence is identical to the sequence obtained from prostatic tissue.

Figure 5 is a blot showing the results of experiments for hormonal regulation of the prostate/KLK-

L1 gene in the BT-474 breast carcinoma cell lines. DHT = dihydrotestosterone. Steroids were added at  $10^{-8}$  M final concentrations. Actin (not regulated by steroid hormones), pS2 (up-regulated by estrogens) and PSA (up-regulated by androgens and progestins), are control genes. Prostase/KLK-L1 is up-regulated by androgens and progestins.

5        Figure 6 is a schematic diagram showing comparison of the genomic structure of PSA, KLK1, KLK2, zyme, neuropsin and prostase/KLK-L1 genes. Exons are shown by open boxes and introns by the connecting lines. Arrow head shows the start codons and the vertical arrow represents stop codons. Letters above boxes indicate relative positions of the catalytic triad; H denotes histidine, D aspartic acid and S serine. Roman numbers indicate intron phases. The intron phase refers to the location of the intron within  
10 the codon; I denotes that the intron occurs after the first nucleotide of the codon, II the intron occurs after the second nucleotide, 0 the intron occurs between codons. Numbers inside boxes indicate exon lengths in base pairs.

Figure 7 shows the genomic organization and partial genomic sequence of the KLK-L2 gene. Intronic sequences are not shown except for the splice junctions. Introns are shown with lower case letters  
15 and exons with capital letters. The start and stop codons are encircled and the exon-intron junctions are boxed. The translated amino acids of the coding region are shown underneath by a single letter abbreviation. The catalytic residues are inside triangles. Putative polyadenylation signal is underlined.

Figure 8 shows an approximate 300 Kb region of almost contiguous genomic sequence around chromosome 19q13.3- q13.4. Genes are represented by horizontal arrows denoting the direction of the  
20 coding sequence. Distances between genes are mentioned in base pairs .

Figure 9 shows the alignment of the deduced amino acid sequence of KLK-L2 with members of the kallikrein multi-gene family. Genes are (from top to bottom) : Prostase/KLK-L1, enamel matrix serine proteinase 1 (EMSP1) (GenBank accession # NP\_004908), KLK-L2, zyme (GenBank accession # Q92876), neuropsin (GenBank accession # BAA28673), trypsin-like serine protease (TLSP) (GenBank  
25 accession # BAA33404), PSA (GenBank accession # P07288), KLK2 (GenBank accession # P20151), KLK1 (GenBank accession # NP\_002248), and trypsinogen ( GenBank accession # P07477). (See SEQ.ID. NOs. 68-77) Dashes represent gaps to bring the sequences to better alignment. The residues of the catalytic triad are represented by (✚) and the 29 invariant serine protease residues by (I or ✚). Conserved areas around the catalytic triad are boxed. The predicted cleavage sites are indicated by (✚). The dotted area  
30 represents the kallikrein loop sequence. The trypsin like cleavage pattern is indicated by (⊙).

Figure 10(A) shows a dendrogram of the predicted phylogenetic tree for some kallikrein genes. Neighbor-joining/UPGMA method was used to align KLK-L2 with other members of the kallikrein gene family. Gene names and accession numbers are listed in Figure 9. The tree grouped the classical kallikreins (KLK1, KLK2, and PSA) together and aligned the KLK-L2 gene in one group with EMSP, prostase, and  
35 TLSP. (B) Plot of hydrophobicity and hydrophilicity of KLK-L2.

Figure 11 is a blot showing tissue expression of KLK-L2 gene as determined by RT-PCR. Actin and PSA are control genes. Interpretations are presented in Table 12.

Figure 12 is a blot showing hormonal regulation of the KLK-L2 gene in BT-474 breast carcinoma

cell lines. DHT = dihydrotestosterone. Steroids were at  $10^{-8}$  M final concentrations. Actin (not regulated by steroid hormones), pS2 (up-regulated by estrogens) and PSA (upregulated by androgens and progestins), are control genes. KLK-L2 is upregulated by estrogens and progestins.

Figure 13 are blots of EtBr-stained agarose gels. Total RNA was extracted from normal, benign, and cancer tissues and used to generate cDNA. PCR was performed on cDNA

Figure 14 shows an approximate 300 Kb region of almost contiguous genomic sequence around chromosome 19q13.3- q13.4. Genes are represented by horizontal arrows denoting the direction of the coding sequence. Gene lengths and distances between genes are rounded to the nearest 0.5 kb. The site of the gap is marked with an asterisk. Telomeric to TLSP there are likely another three kallikrein-like genes.

Figure 15 shows the genomic organization and partial genomic sequence of the KLK-L3 gene. Intronic sequences are not shown except for the splice junctions. Introns are shown with lower case letters and exons with capital letters. For the full sequence, see SEQ.ID. NO. 21. The start and stop codons are encircled and the exon-intron junctions are boxed. The translated amino acids of the coding region are shown underneath by a single letter abbreviation. The catalytic residues are inside triangles. Putative polyadenylation signal is underlined.

Figure 16 is a plot of hydrophobicity and hydrophilicity, comparing the pattern of the KLK-L3 with that of the zyme gene. Note the hydrophobic region around the first twenty amino acids, likely representing the signal peptide.

Figure 17 is an alignment of the deduced amino acid sequence of KLK-L3 with members of the kallikrein multi-gene family. Genes are (from top to bottom and in brackets is the GenBank accession #): PSA (P07288), KLK2 (P20151), KLK1 (NP002248), trypsinogen (P07477), KLK-L3 (AF135026), trypsin-like serine protease (TLSP) (BAA33404), neuropsin (BAA28673), zyme (Q92876), human stratum corneum chymotryptic enzyme (HSCCE) (AAD49718), and/prostase/KLK-L1 (AAD21581). (See SEQ.ID. NOs. 78 to 84). Dashes represent gaps to bring the sequences to better alignment. The residues of the catalytic triad are bold and in italics, and the 29 invariant serine protease residues are denoted by ( $\diamond$ ). Cysteine residues are marked by ( $\bullet$ ). Conserved areas around the catalytic triad are highlighted in black. The arrow heads ( $\wedge$ ) represent the potential cleavage sites. The dotted area represents the kallikrein loop sequence.

Figure 18 is a dendrogram of the predicted phylogenetic tree for some serine proteases and kallikrein genes. Neighbor-joining/UPGMA method was used to align KLK-L3 with other members of the kallikrein gene family. Gene names and accession numbers are listed in Figure 17. The tree grouped the classical kallikreins (KLK1, KLK2, and PSA) together and aligned the KLK-L3 gene in one group with TLSP, neuropsin, and NES 1 genes. KLK-L4 (SEQ.ID.NO. 43) lies further telomeric to TLSP (21).

Figure 19 is a blot showing tissue expression of the KLK-L3 gene as determined by RT-PCR. Actin and PSA are control genes.

Figure 20 shows hormonal regulation of the KLK-L3 gene in the BT-474 breast carcinoma cell line. DHT = dihydrotestosterone. Steroids were at  $10^{-8}$  M final concentrations. Actin (not regulated by steroid hormones), pS2 (up-regulated by estrogens) and PSA (upregulated by androgens and progestins),

are control genes. KLK-L3 is upregulated by progestins, estrogens and androgens, in that order.

Figure 21 is a schematic diagram showing the comparison of the genomic structure of PSA, KLK2, neuropsin, NES1, and KLK-L3 genes. Exons are shown by black boxes and introns by the connecting lines. Arrowheads show the start codon, and arrows show the stop codon. Letters above boxes indicate relative positions of the catalytic triad; H denotes histidine, D aspartic acid and S serine. Roman numbers indicate intron phases. The intron phase refers to the location of the intron within the codon; I denotes that the intron occurs after the first nucleotide of the codon, II the intron occurs after the second nucleotide, 0 the intron occurs between codons. Numbers inside boxes indicate exon lengths in base pairs.

Figure 22 shows a comparative genomic structure of the ESTs (Table 16), the clone from The German Genome Project, and the long form of KLK-L4. Exons are represented by solid bars and introns by the connecting lines. Exon numbers on top of solid bars refer to GenBank submission #AF135024. The EST IDs represent GenBank accession numbers. Asterisks represent the positions of stop codons. Horizontal arrows indicate the direction of the PCR primers (described in Table 15) and arrowheads their position along the exons. Vertical dotted lines show alignment of identical fragments.

Figure 23 shows tissue expression of the KLK-L4 gene as determined by RT-PCR. Actin and PSA are control genes. KLK-L4 is highly expressed in breast, prostate, salivary gland and testis.

Figure 24 in the Upper Panel is a Diagram showing the comparative genomic structure of the long KLK-L4 form and the short KLK-L4 variant. Exons are represented by boxes and introns by the connecting lines. Exon numbers refer to SEQ. ID. NO. 43 and GenBank Accession No. AF135024. The black region indicates the extra fragment (214 bp) that is found in the long, but not in the short form of the gene. The positions of the stop codons of the two forms are marked with asterisks. Frame shifting occurs as a result of utilization of an alternative splice site, and a stop codon is generated at the beginning of exon 4 in the short form. The Lower Panel shows PCR products of the amplification of the KLK-L4 gene using L4-R1 and L4-X1 primers (Figure 22 and Table 15). Note the predominant long form and a minor band representing the short form of KLK-L4 mRNA. (M); Markers with sizes in bp shown on the left. Tissues used: (1), salivary gland; (2), mammary gland; (3), prostate; (4), testis; (5), uterus; (6), breast cancer tissue; (7), negative control.

Figure 25 shows the genomic organization and partial genomic sequence of the KLK-L4 gene. Intronic sequences are not shown except for the splice junction areas. Introns are shown with lower case letters and exons with capital letters. For full sequence, see SEQ. ID. NO.43 or GenBank Accession #AF135024. The start and stop codons are encircled and the exon-intron junctions are underlined. The translated amino acids of the coding region are shown underneath by a single letter abbreviation. The catalytic residues are boxed. The putative polyadenylation signal is underlined.

Figure 26 is a plot of hydrophobicity and hydrophilicity of the KLK-L4 protein, as compared with the glandular kallikrein gene 2 (KLK2). Note the hydrophobic region at the amino terminus, suggesting presence of a signal peptide.

Figure 27 shows an alignment of the deduced amino acid sequence of KLK-L4 with members of the kallikrein multi-gene family. Genes are (from top to bottom, and in brackets are the GenBank accession

#): KLK-L1/prostase (AAD21581), enamel matrix serine proteinase I (EMSP) (NP\_004908), KLK-L2 (AF135028), PSA (P07288), KLK2 (P20151), KLK1 (NP\_002248), trypsinogen (P07477), zyme (Q92876), KLK-L4 (AF135024), trypsin-like serine protease (TLSP) (BAA33404), KLK-L3 (AF135026), neuropsin (BAA28673), and the normal epithelial cell-specific 1 gene (NES1) (O43240). (See SEQ.ID. NOs. 78-88). Dashes represent gaps to bring the sequences to better alignment. The residues of the catalytic triad are typed in bold, and conserved motifs around them are highlighted in grey. The 29 invariant serine protease residues are denoted by (•), and the cysteine residues by (♦). The predicted cleavage sites are indicated by (▲). The dotted area represents the kallikrein loop sequence. The trypsin-like cleavage pattern of KLK-L4 with the D residue, is indicated by (⊙).

Figure 28 shows an approximate 300 Kb region of almost contiguous genomic sequence around chromosome 19q13.3- q13.4. Genes are represented by horizontal arrows denoting the direction of the coding sequence. Their lengths are shown on top of each arrow. Distances between genes are mentioned in base pairs below the arrows. The distance between KLK1 and PSA is not accurately known. For gene names, see under abbreviations.

Figure 29 shows is a dendrogram of the predicted phylogenetic tree for some kallikrein and serine protease genes. The neighbor-joining/UPGMA method was used to align KLK-L4 with other serine proteases and members of the kallikrein gene family. The tree grouped the classical kallikreins (KLK1, KLK2, and PSA) together and aligned the KLK-L4 gene in one group with zyme, NES1, neuropsin, KLK-L3, and TLSP. Other serine proteases were aligned in different groups, as shown.

Figure 30 is a blot showing the hormonal regulation of the KLK-L4 gene in the BT-474 breast carcinoma cell line. DHT = dihydrotestosterone. Steroids were added at  $10^{-8}$  M final concentrations. Actin (not regulated by steroid hormones), pS2 (up-regulated by estrogens) and PSA (upregulated by androgens and progestins) are control genes. KLK-L4 is up-regulated by androgens and progestins and to a lesser extent by estrogens. H<sub>2</sub>O was used to check for PCR specificity in all PCR reactions. For more details, see text.

Figure 31 is a schematic diagram showing the comparison of the genomic structure of PSA, KLK2, neuropsin, NES1, and KLK-L4 genes. Exons are shown by black boxes and introns by the connecting lines. The arrowhead shows the start codons and the arrow the stop codons. Letters above boxes indicate the relative positions of the amino acids of the catalytic triad; H denotes histidine, D aspartic acid and S serine. Roman numbers indicate intron phases. The intron phase refers to the location of the intron within the codon; I, the intron occurs after the first nucleotide of the codon, II the intron occurs after the second nucleotide, 0 the intron occurs between codons. Numbers inside boxes indicate exon lengths in base pairs. The question mark indicates the possibility of more untranslated bases.

Figure 32 is a diagram showing the comparative genomic structure of the three splice forms of KLK-L5; the classical kallikrein form, related protein-1, and related protein-2. Exons are represented by solid bars and introns by the connecting lines. Exon numbers refer to SEQ.ID. NO.56 and GenBank Accession #AF135025. Start codons are represented by the inverted arrowhead (▼) and stop codons are represented by asterisks (\*). Primer locations are represented by vertical arrowheads (▲) and their

directions by horizontal arrows. For primer sequences and codes see Table 17 and SEQ.ID. NOs. 61-64, and 9-12.

Figure 33 shows the genomic organization and partial genomic sequence of the KLK-L5 gene. Intronic sequences are not shown except for short sequences around the splice junctions. Introns are shown with lower case letters and exons with capital letters. For full sequence, see SEQ.ID.NO. 56. The start and stop codons are encircled and the exon-intron junctions are underlined. The translated amino acids of the coding region are shown underneath by a single letter abbreviation. The catalytic residues are boxed. Putative polyadenylation signal is underlined. The extra intron of the related protein-1 form is represented by non-bold capital letters between brackets. When this intron is spliced, the frame continues with codon AAC (asparagine, N, instead of lysine, K) until it encounters the stop codon TAA (encircled).

Figure 34 is a schematic diagram showing the comparison of the genomic structure of PSA, KLK2, neuropsin, NES1, KLK-L4 and KLK-L5 genes. Exons are shown by solid bars and introns by the connecting lines. Arrowhead marks the site of the start codon, and the arrow represents the stop codon. Letters above boxes indicate relative positions of the catalytic triad; H denotes histidine, D aspartic acid and S serine. Roman numbers indicate intron phases. The intron phase refers to the location of the intron within the codon; I denotes that the intron occurs after the first nucleotide of the codon, II the intron occurs after the second nucleotide, 0 the intron occurs between codons. Numbers inside boxes indicate exon lengths in base pairs. Question marks indicate that exact length is not accurately known.

Figure 35 is a plot of hydrophobicity and hydrophilicity of KLK-L5 protein compared to prostate specific antigen (PSA). The hydrophobic N-terminus may harbor a signal and activation peptide.

Figure 36 shows an alignment of the deduced amino acid sequence of KLK-L5 with members of the kallikrein multigene family. (See SEQ.ID. NOs. 78-81, 83, 84). Dashes represent gaps to bring the sequences to better alignment. The residues of the catalytic triad are represented by bold letters, and the 29 invariant serine protease residues are marked with (•). The cysteine residues are marked by (♦). Conserved areas are highlighted in grey. The predicted cleavage sites in signal peptide are indicated by (♣). The dotted area represents the kallikrein loop sequence. A vertical arrow marks the trypsin like cleavage site.

Figure 37 is a dendrogram of the predicted phylogenetic tree for some serine proteases and other kallikrein proteins. Neighbor-joining/UPGMA method was used to align KLK-L5 with other serine proteases and members of the kallikrein gene family. The tree grouped the classical kallikreins (hK1, hK2, and PSA) together and aligned the KLK-L5 protein in one group with NES1 and neuropsin. Other serine proteases were aligned in different groups.

Figure 38 shows tissue expression of the KLK-L5 gene as determined by RT-PCR. The upper band (905 base pairs, bp) is the classical form (see Figure 32, the middle (776 bp) the related protein-1, and the lower band (644 bp) the related protein-2. For splice variant discussion see text. The primers used were L5-F2 and L5-R2, as shown in Table 17.

Figure 39 shows hormonal regulation of the KLK-L5 gene in the LnCaP prostatic carcinoma cell line, BT-474 and T-47D breast carcinoma cell lines. Steroids were at  $10^{-8}$  M final concentration. Actin (not

regulated by steroid hormones) was used as a control gene. Note detection of three isoforms only in LNCaP.

Figure 40 shows the expression of the KLK-L5 gene in breast cancer (1-17) and normal (18) tissues. Note complete absence of expression in 12 cancer tissues. For isoforms see also Figure 38.

5 Figure 41 shows the full structure of a KLK-L6 nucleic acid molecule;

Figure 42 is a plot of hydrophobicity and hydrophilicity of KLK-L6 protein compared to prostate specific antigen (PSA).

10 Figure 43 shows an alignment of the deduced amino acid sequence of KLK-L6 with members of the kallikrein multigene family. (See SEQ.ID. NOs. 78-81, 83, 84). Dashes represent gaps to bring the sequences to better alignment.

Figure 44 is a dendrogram of the predicted phylogenetic tree for some serine proteases and other kallikrein proteins. Neighbor-joining/UPGMA method was used to align KLK-L6 with other serine proteases and members of the kallikrein gene family.

#### DETAILED DESCRIPTION OF THE INVENTION

15 In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See for example, Sambrook, Fritsch, & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *DNA Cloning: A Practical Approach*, Volumes I and II (D.N. Glover ed. 1985); *Oligonucleotide Synthesis* (M..J. Gait ed. 1984); *Nucleic Acid Hybridization* B.D. Hames & S.J. Higgins eds. (1985); *Transcription and Translation* B.D. Hames & S.J. Higgins eds (1984); *Animal Cell Culture* R.I. Freshney, ed. (1986); *Immobilized Cells and enzymes* IRL Press, (1986); and B. Perbal, *A Practical Guide to Molecular Cloning* (1984).

##### 1. Nucleic Acid Molecules of the Invention

25 As hereinbefore mentioned, the invention provides an isolated nucleic acid molecule having a sequence encoding a KLK-L Protein. The term "isolated" refers to a nucleic acid substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical reactants, or other chemicals when chemically synthesized. An "isolated" nucleic acid may also be free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid molecule) from which the nucleic acid is derived. The term "nucleic acid" is intended to include DNA and RNA and can be either double stranded or single stranded. In an embodiment, a nucleic acid molecule encodes a KLK-L Protein comprising an amino acid sequence as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, preferably a nucleic acid molecule comprising a nucleic acid sequence as shown in SEQ.ID.NO. 1, 13, 21, 43, 56, or 65.

35 The invention includes nucleic acid sequences complementary to a nucleic acid encoding a KLK-L Protein comprising an amino acid sequence as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, preferably the nucleic acid sequences complementary to a full nucleic acid sequence shown in SEQ.ID.NO. 1, 13, 21, 43, 56, or 65.

The invention includes nucleic acid molecules having substantial sequence identity or homology to nucleic acid sequences of the invention or encoding proteins having substantial identity or similarity to the amino acid sequence shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67. Preferably, the nucleic acids have substantial sequence identity for example at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% nucleic acid identity; more preferably 90% nucleic acid identity; and most preferably at least 95%, 96%, 97%, 98%, or 99% sequence identity. "Identity" as known in the art and used herein, is a relationship between two or more amino acid sequences or two or more nucleic acid sequences, as determined by comparing the sequences. It also refers to the degree of sequence relatedness between amino acid or nucleic acid sequences, as the case may be, as determined by the match between strings of such sequences. Identity and similarity are well known terms to skilled artisans and they can be calculated by conventional methods (for example see Computational Molecular Biology, Lesk, A.M. ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D.W. ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A.M. and Griffin, H.G. eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G. Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J. eds. M. Stockton Press, New York, 1991, Carillo, H. and Lipman, D., SIAM J. Applied Math. 48:1073, 1988). Methods which are designed to give the largest match between the sequences are generally preferred. Methods to determine identity and similarity are codified in publicly available computer programs including the GCG program package (Devereux J. et al., Nucleic Acids Research 12(1): 387, 1984); BLASTP, BLASTN, and FASTA (Atschul, S.F. et al. J. Molec. Biol. 215: 403-410, 1990). The BLAST X program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S. et al. NCBI NLM NIH Bethesda, Md. 20894; Altschul, S. et al. J. Mol. Biol. 215: 403-410, 1990).

Isolated nucleic acid molecules encoding a KLK-L Protein, and having a sequence which differs from a nucleic acid sequence of the invention due to degeneracy in the genetic code are also within the scope of the invention. Such nucleic acids encode functionally equivalent proteins (e.g., a KLK-L Protein) but differ in sequence from the sequence of a KLK-L Protein due to degeneracy in the genetic code. As one example, DNA sequence polymorphisms within the nucleotide sequence of a KLK-L Protein may result in silent mutations which do not affect the amino acid sequence. Variations in one or more nucleotides may exist among individuals within a population due to natural allelic variation. Any and all such nucleic acid variations are within the scope of the invention. DNA sequence polymorphisms may also occur which lead to changes in the amino acid sequence of a KLK-L Protein. These amino acid polymorphisms are also within the scope of the present invention.

Another aspect of the invention provides a nucleic acid molecule which hybridizes under stringent conditions, preferably high stringency conditions to a nucleic acid molecule which comprises a sequence which encodes a KLK-L Protein having an amino acid sequence shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67. Appropriate stringency conditions which promote DNA hybridization are known to those skilled in the art, or can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C,

followed by a wash of 2.0 x SSC at 50°C may be employed. The stringency may be selected based on the conditions used in the wash step. By way of example, the salt concentration in the wash step can be selected from a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be at high stringency conditions, at about 65°C.

5 It will be appreciated that the invention includes nucleic acid molecules encoding a KLK-L Related Protein including truncations of a KLK-L Protein, and analogs of a KLK-L Protein as described herein. It will further be appreciated that variant forms of the nucleic acid molecules of the invention which arise by alternative splicing of an mRNA corresponding to a cDNA of the invention are encompassed by the invention. (See for example, splice variants of KLK-L5, SEQ.ID.NO. 58, 59, and 60.)

10 An isolated nucleic acid molecule of the invention which comprises DNA can be isolated by preparing a labelled nucleic acid probe based on all or part of a nucleic acid sequence of the invention. The labeled nucleic acid probe is used to screen an appropriate DNA library (e.g. a cDNA or genomic DNA library). For example, a cDNA library can be used to isolate a cDNA encoding a KLK-L Related Protein by screening the library with the labeled probe using standard techniques. Alternatively, a genomic DNA  
15 library can be similarly screened to isolate a genomic clone encompassing a gene encoding a KLK-L Related Protein. Nucleic acids isolated by screening of a cDNA or genomic DNA library can be sequenced by standard techniques.

An isolated nucleic acid molecule of the invention which is DNA can also be isolated by selectively amplifying a nucleic acid encoding a KLK-L Related Protein using the polymerase chain  
20 reaction (PCR) methods and cDNA or genomic DNA. It is possible to design synthetic oligonucleotide primers from the nucleotide sequence of the invention for use in PCR. A nucleic acid can be amplified from cDNA or genomic DNA using these oligonucleotide primers and standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. cDNA may be prepared from mRNA, by isolating total cellular mRNA by a variety of techniques,  
25 for example, by using the guanidinium-thiocyanate extraction procedure of Chirgwin et al., Biochemistry, 18, 5294-5299 (1979). cDNA is then synthesized from the mRNA using reverse transcriptase (for example, Moloney MLV reverse transcriptase available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase available from Seikagaku America, Inc., St. Petersburg, FL).

An isolated nucleic acid molecule of the invention which is RNA can be isolated by cloning a  
30 cDNA encoding a KLK-L Related Protein into an appropriate vector which allows for transcription of the cDNA to produce an RNA molecule which encodes a KLK-L Related Protein. For example, a cDNA can be cloned downstream of a bacteriophage promoter, (e.g. a T7 promoter) in a vector, cDNA can be transcribed *in vitro* with T7 polymerase, and the resultant RNA can be isolated by conventional techniques.

Nucleic acid molecules of the invention may be chemically synthesized using standard techniques.  
35 Methods of chemically synthesizing polydeoxynucleotides are known, including but not limited to solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071).

Determination of whether a particular nucleic acid molecule encodes a KLK-L Related Protein can be accomplished by expressing the cDNA in an appropriate host cell by standard techniques, and testing the expressed protein in the methods described herein. A cDNA encoding a KLK-L Related Protein can be sequenced by standard techniques, such as dideoxynucleotide chain termination or Maxam-Gilbert  
5 chemical sequencing, to determine the nucleic acid sequence and the predicted amino acid sequence of the encoded protein.

The initiation codon and untranslated sequences of a KLK-L Related Protein may be determined using computer software designed for the purpose, such as PC/Gene (IntelliGenetics Inc., Calif.). The intron-exon structure and the transcription regulatory sequences of a gene encoding a KLK-L Related  
10 Protein may be confirmed by using a nucleic acid molecule of the invention encoding a KLK-L Related Protein to probe a genomic DNA clone library. Regulatory elements can be identified using standard techniques. The function of the elements can be confirmed by using these elements to express a reporter gene such as the lacZ gene which is operatively linked to the elements. These constructs may be introduced into cultured cells using conventional procedures or into non-human transgenic animal models. In addition  
15 to identifying regulatory elements in DNA, such constructs may also be used to identify nuclear proteins interacting with the elements, using techniques known in the art.

In a particular embodiment of the invention, the nucleic acid molecules isolated using the methods described herein are mutant *klk-l* gene alleles. The mutant alleles may be isolated from individuals either known or proposed to have a genotype which contributes to the symptoms of for example, cancer (e.g.,  
20 breast, testicular, brain, colon, and prostate cancer). Mutant alleles and mutant allele products may be used in therapeutic and diagnostic methods described herein. For example, a cDNA of a mutant *klk-l* gene may be isolated using PCR as described herein, and the DNA sequence of the mutant allele may be compared to the normal allele to ascertain the mutation(s) responsible for the loss or alteration of function of the mutant gene product. A genomic library can also be constructed using DNA from an individual suspected  
25 of or known to carry a mutant allele, or a cDNA library can be constructed using RNA from tissue known, or suspected to express the mutant allele. A nucleic acid encoding a normal *klk-l* gene or any suitable fragment thereof, may then be labeled and used as a probe to identify the corresponding mutant allele in such libraries. Clones containing mutant sequences can be purified and subjected to sequence analysis. In addition, an expression library can be constructed using cDNA from RNA isolated from a tissue of an  
30 individual known or suspected to express a mutant *klk-l* allele. Gene products made by the putatively mutant tissue may be expressed and screened, for example using antibodies specific for a KLK-L Related Protein as described herein. Library clones identified using the antibodies can be purified and subjected to sequence analysis.

The sequence of a nucleic acid molecule of the invention, or a fragment of the molecule, may be  
35 inverted relative to its normal presentation for transcription to produce an antisense nucleic acid molecule. An antisense nucleic acid molecule may be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art.

## 2. Proteins of the Invention

An amino acid sequence of a KLK-L Protein comprises a sequence as shown in Tables 1 to 5 or SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67.

In addition to proteins comprising an amino acid sequence as shown in Tables 1 to 5 or SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, the proteins of the present invention include  
5 truncations of a KLK-L Protein, analogs of a KLK-L Protein, and proteins having sequence identity or similarity to a KLK-L Protein, and truncations thereof as described herein (i.e. included in KLK-L Related Proteins). Truncated proteins may comprise peptides of between 3 and 70 amino acid residues, ranging in size from a tripeptide to a 70 mer polypeptide.

The truncated proteins may have an amino group (-NH<sub>2</sub>), a hydrophobic group (for example, carbobenzoxyl, dansyl, or T-butyloxycarbonyl), an acetyl group, a 9-fluorenylmethoxy-carbonyl (PMOC) group, or a macromolecule including but not limited to lipid-fatty acid conjugates, polyethylene glycol, or carbohydrates at the amino terminal end. The truncated proteins may have a carboxyl group, an amido group, a T-butyloxycarbonyl group, or a macromolecule including but not limited to lipid-fatty acid conjugates, polyethylene glycol, or carbohydrates at the carboxy terminal end.  
10

The proteins of the invention may also include analogs of a KLK-L Protein, and/or truncations thereof as described herein, which may include, but are not limited to a KLK-L Protein, containing one or more amino acid substitutions, insertions, and/or deletions. Amino acid substitutions may be of a conserved or non-conserved nature. Conserved amino acid substitutions involve replacing one or more amino acids of a KLK-L Protein amino acid sequence with amino acids of similar charge, size, and/or hydrophobicity characteristics. When only conserved substitutions are made the resulting analog is preferably functionally equivalent to a KLK-L Protein. Non-conserved substitutions involve replacing one or more amino acids of the KLK-L Protein amino acid sequence with one or more amino acids which possess dissimilar charge, size, and/or hydrophobicity characteristics.  
15  
20

One or more amino acid insertions may be introduced into a KLK-L Protein. Amino acid  
25 insertions may consist of single amino acid residues or sequential amino acids ranging from 2 to 15 amino acids in length.

Deletions may consist of the removal of one or more amino acids, or discrete portions from a KLK-L Protein sequence. The deleted amino acids may or may not be contiguous. The lower limit length of the resulting analog with a deletion mutation is about 10 amino acids, preferably 20 to 40 amino acids.  
30

The proteins of the invention include proteins with sequence identity or similarity to a KLK-L Protein and/or truncations thereof as described herein. Such KLK-L Proteins include proteins whose amino acid sequences are comprised of the amino acid sequences of KLK-L Protein regions from other species that hybridize under selected hybridization conditions (see discussion of stringent hybridization conditions herein) with a probe used to obtain a KLK-L Protein. These proteins will generally have the same regions  
35 which are characteristic of a KLK-L Protein. Preferably a protein will have substantial sequence identity for example, about 30%, 35%, 40%, 45%, 50%, 60%, 65%, 70%, 75%, 80%, or 85% identity, preferably 90% identity, more preferably at least 95%, 96%, 97%, 98%, or 99% identity, and most preferably 98% identity with an amino acid sequence shown in Tables 1 to 5 or SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57,

58, 59, 60, 66, or 67.

A percent amino acid sequence homology, similarity or identity is calculated as the percentage of aligned amino acids that match the reference sequence using known methods as described herein.

5 The invention also contemplates isoforms of the proteins of the invention. An isoform contains the same number and kinds of amino acids as a protein of the invention, but the isoform has a different molecular structure. Isoforms contemplated by the present invention preferably have the same properties as a protein of the invention as described herein.

10 The present invention also includes KLK-L Related Proteins conjugated with a selected protein, or a marker protein (see below) to produce fusion proteins. Additionally, immunogenic portions of a KLK-L Protein and a KLK-L Protein Related Protein are within the scope of the invention.

15 A KLK-L Related Protein of the invention may be prepared using recombinant DNA methods. Accordingly, the nucleic acid molecules of the present invention having a sequence which encodes a KLK-L Related Protein of the invention may be incorporated in a known manner into an appropriate expression vector which ensures good expression of the protein. Possible expression vectors include but are not limited to cosmids, plasmids, or modified viruses (e.g. replication defective retroviruses, adenoviruses and adeno-associated viruses), so long as the vector is compatible with the host cell used.

20 The invention therefore contemplates a recombinant expression vector of the invention containing a nucleic acid molecule of the invention, and the necessary regulatory sequences for the transcription and translation of the inserted protein-sequence. Suitable regulatory sequences may be derived from a variety of sources, including bacterial, fungal, viral, mammalian, or insect genes (For example, see the regulatory sequences described in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). Selection of appropriate regulatory sequences is dependent on the host cell chosen as discussed below, and may be readily accomplished by one of ordinary skill in the art. The necessary regulatory sequences may be supplied by the native KLK-L Protein and/or its flanking regions.

25 The invention further provides a recombinant expression vector comprising a DNA nucleic acid molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is linked to a regulatory sequence in a manner which allows for expression, by transcription of the DNA molecule, of an RNA molecule which is antisense to the nucleic acid sequence of a protein of the invention or a fragment thereof. Regulatory sequences linked to the antisense nucleic acid can be chosen 30 which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance a viral promoter and/or enhancer, or regulatory sequences can be chosen which direct tissue or cell type specific expression of antisense RNA.

35 The recombinant expression vectors of the invention may also contain a marker gene which facilitates the selection of host cells transformed or transfected with a recombinant molecule of the invention. Examples of marker genes are genes encoding a protein such as G418 and hygromycin which confer resistance to certain drugs,  $\beta$ -galactosidase, chloramphenicol acetyltransferase, firefly luciferase, or an immunoglobulin or portion thereof such as the Fc portion of an immunoglobulin preferably IgG. The markers can be introduced on a separate vector from the nucleic acid of interest.

The recombinant expression vectors may also contain genes which encode a fusion moiety which provides increased expression of the recombinant protein; increased solubility of the recombinant protein; and aid in the purification of the target recombinant protein by acting as a ligand in affinity purification. For example, a proteolytic cleavage site may be added to the target recombinant protein to allow separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the recombinant protein.

The recombinant expression vectors may be introduced into host cells to produce a transformant host cell. "Transformant host cells" include host cells which have been transformed or transfected with a recombinant expression vector of the invention. The terms "transformed with", "transfected with", "transformation" and "transfection" encompass the introduction of a nucleic acid (e.g. a vector) into a cell by one of many standard techniques. Prokaryotic cells can be transformed with a nucleic acid by, for example, electroporation or calcium-chloride mediated transformation. A nucleic acid can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride coprecipitation, DEAE-dextran-mediated transfection, lipofectin, electroporation or microinjection. Suitable methods for transforming and transfecting host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

Suitable host cells include a wide variety of prokaryotic and eukaryotic host cells. For example, the proteins of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus), yeast cells, or mammalian cells. Other suitable host cells can be found in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1991).

A host cell may also be chosen which modulates the expression of an inserted nucleic acid sequence, or modifies (e.g. glycosylation or phosphorylation) and processes (e.g. cleaves) the protein in a desired fashion. Host systems or cell lines may be selected which have specific and characteristic mechanisms for post-translational processing and modification of proteins. For example, eukaryotic host cells including CHO, VERO, BHK, HeLA, COS, MDCK, 293, 3T3, and WI38 may be used. For long-term high-yield stable expression of the protein, cell lines and host systems which stably express the gene product may be engineered.

Host cells and in particular cell lines produced using the methods described herein may be particularly useful in screening and evaluating compounds that modulate the activity of a KLK-L Related Protein.

The proteins of the invention may also be expressed in non-human transgenic animals including but not limited to mice, rats, rabbits, guinea pigs, micro-pigs, goats, sheep, pigs, non-human primates (e.g. baboons, monkeys, and chimpanzees) [see Hammer et al. (Nature 315:680-683, 1985), Palmiter et al. (Science 222:809-814, 1983), Brinster et al. (Proc Natl. Acad. Sci USA 82:44384442, 1985), Palmiter and Brinster (Cell. 41:343-345, 1985) and U.S. Patent No. 4,736,866)]. Procedures known in the art may be

used to introduce a nucleic acid molecule of the invention encoding a KLK-L Related Protein into animals to produce the founder lines of transgenic animals. Such procedures include pronuclear microinjection, retrovirus mediated gene transfer into germ lines, gene targeting in embryonic stem cells, electroporation of embryos, and sperm-mediated gene transfer.

5           The present invention contemplates a transgenic animal that carries the *KLK-L* gene in all their cells, and animals which carry the transgene in some but not all their cells. The transgene may be integrated as a single transgene or in concatamers. The transgene may be selectively introduced into and activated in specific cell types (See for example, Lasko et al, 1992 Proc. Natl. Acad. Sci. USA 89: 6236). The transgene may be integrated into the chromosomal site of the endogenous gene by gene targeting. The transgene may be selectively introduced into a particular cell type inactivating the endogenous gene in that cell type (See  
10           Gu et al Science 265: 103-106).

          The expression of a recombinant KLK-L Related Protein in a transgenic animal may be assayed using standard techniques. Initial screening may be conducted by Southern Blot analysis, or PCR methods to analyze whether the transgene has been integrated. The level of mRNA expression in the tissues of  
15           transgenic animals may also be assessed using techniques including Northern blot analysis of tissue samples, *in situ* hybridization, and RT-PCR. Tissue may also be evaluated immunocytochemically using antibodies against KLK-L Protein.

          Proteins of the invention may also be prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, 1964, J. Am. Chem. Assoc. 85:2149-  
20           2154) or synthesis in homogenous solution (Houbenweyl, 1987, Methods of Organic Chemistry, ed. E. Wansch, Vol. 15 I and II, Thieme, Stuttgart).

          N-terminal or C-terminal fusion proteins comprising a KLK-L Related Protein of the invention conjugated with other molecules, such as proteins, may be prepared by fusing, through recombinant techniques, the N-terminal or C-terminal of a KLK-L Related Protein, and the sequence of a selected  
25           protein or marker protein with a desired biological function. The resultant fusion proteins contain KLK-L Protein fused to the selected protein or marker protein as described herein. Examples of proteins which may be used to prepare fusion proteins include immunoglobulins, glutathione-S-transferase (GST), hemagglutinin (HA), and truncated myc.

### 3.       Antibodies

30           KLK-L Related Proteins of the invention can be used to prepare antibodies specific for the proteins. Antibodies can be prepared which bind a distinct epitope in an unconserved region of the protein. An unconserved region of the protein is one that does not have substantial sequence homology to other proteins. A region from a conserved region such as a well-characterized domain can also be used to prepare an antibody to a conserved region of a KLK-L Related Protein. Antibodies having specificity for a KLK-L  
35           Related Protein may also be raised from fusion proteins created by expressing fusion proteins in bacteria as described herein.

          The invention can employ intact monoclonal or polyclonal antibodies, and immunologically active fragments (e.g. a Fab, (Fab)<sub>2</sub> fragment, or Fab expression library fragments and epitope-binding fragments

thereof), an antibody heavy chain, and antibody light chain, a genetically engineered single chain Fv molecule (Ladner et al, U.S. Pat. No. 4,946,778), or a chimeric antibody, for example, an antibody which contains the binding specificity of a murine antibody, but in which the remaining portions are of human origin. Antibodies including monoclonal and polyclonal antibodies, fragments and chimeras, may be prepared using methods known to those skilled in the art.

#### 4. Applications of the Nucleic Acid Molecules, KLK-L Related Proteins, and Antibodies of the Invention

The nucleic acid molecules, KLK-L Related Proteins, and antibodies of the invention may be used in the prognostic and diagnostic evaluation of cancer (e.g. breast, testicular, and prostate cancer) or other conditions, and the identification of subjects with a predisposition to cancer (Section 4.1.1 and 4.1.2). Methods for detecting nucleic acid molecules and KLK-L Related Proteins of the invention, can be used to monitor conditions including cancer, by detecting KLK-L Related Proteins and nucleic acid molecules encoding KLK-L Related Proteins. It would also be apparent to one skilled in the art that the methods described herein may be used to study the developmental expression of KLK-L Related Proteins and, accordingly, will provide further insight into the role of KLK-L Related Proteins. The applications of the present invention also include methods for the identification of compounds that modulate the biological activity of *KLK-L* or KLK-L Related Proteins (Section 4.2). The compounds, antibodies etc. may be used for the treatment of cancer (Section 4.3).

##### 4.1 Diagnostic Methods

A variety of methods can be employed for the diagnostic and prognostic evaluation of conditions including cancer (e.g. breast, testicular, and prostate cancer), and the identification of subjects with a predisposition to such conditions. Such methods may, for example, utilize nucleic acid molecules of the invention, and fragments thereof, and antibodies directed against KLK-L Related Proteins, including peptide fragments. In particular, the nucleic acids and antibodies may be used, for example, for: (1) the detection of the presence of *KLK-L* mutations, or the detection of either over- or under-expression of *KLK-L* mRNA relative to a non-disorder state or the qualitative or quantitative detection of alternatively spliced forms of *KLK-L* transcripts which may correlate with certain conditions or susceptibility toward such conditions; and (2) the detection of either an over- or an under-abundance of KLK-L Related Proteins relative to a non- disorder state or the presence of a modified (e.g., less than full length) KLK-L Protein which correlates with a disorder state, or a progression toward a disorder state.

The methods described herein may be performed by utilizing pre-packaged diagnostic kits comprising at least one specific *KLK-L* nucleic acid or antibody described herein, which may be conveniently used, e.g., in clinical settings, to screen and diagnose patients and to screen and identify those individuals exhibiting a predisposition to developing a disorder.

Nucleic acid-based detection techniques are described, below, in Section 4.1.1. Peptide detection techniques are described, below, in Section 4.1.2. The samples that may be analyzed using the methods of the invention include those which are known or suspected to express *KLK-L* or contain KLK-L Related Proteins. The samples may be derived from a patient or a cell culture, and include but are not limited to

biological fluids, tissue extracts, freshly harvested cells, and lysates of cells which have been incubated in cell cultures.

Oligonucleotides or longer fragments derived from any of the nucleic acid molecules of the invention may be used as targets in a microarray. The microarray can be used to simultaneously monitor the expression levels of large numbers of genes and to identify genetic variants, mutations, and polymorphisms. The information from the microarray may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

The preparation, use, and analysis of microarrays are well known to a person skilled in the art. (See, for example, Brennan, T. M. et al. (1995) U.S. Pat. No. 5,474,796; Schena, et al. (1996) Proc. Natl. Acad. Sci. 93:10614-10619; Baldeschweiler et al. (1995), PCT Application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R. A. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-2155; and Heller, M. J. et al. (1997) U.S. Pat. No. 5,605,662.)

#### **4.1.1 Methods for Detecting Nucleic Acid Molecules of the Invention**

The nucleic acid molecules of the invention allow those skilled in the art to construct nucleotide probes for use in the detection of nucleic acid sequences of the invention in samples. Suitable probes include nucleic acid molecules based on nucleic acid sequences encoding at least 5 sequential amino acids from regions of a KLK-L Protein, preferably they comprise 15 to 30 nucleotides. A nucleotide probe may be labeled with a detectable substance such as a radioactive label which provides for an adequate signal and has sufficient half-life such as  $^{32}\text{P}$ ,  $^3\text{H}$ ,  $^{14}\text{C}$  or the like. Other detectable substances which may be used include antigens that are recognized by a specific labeled antibody, fluorescent compounds, enzymes, antibodies specific for a labeled antigen, and luminescent compounds. An appropriate label may be selected having regard to the rate of hybridization and binding of the probe to the nucleotide to be detected and the amount of nucleotide available for hybridization. Labeled probes may be hybridized to nucleic acids on solid supports such as nitrocellulose filters or nylon membranes as generally described in Sambrook et al, 1989, Molecular Cloning, A Laboratory Manual (2nd ed.). The nucleic acid probes may be used to detect genes, preferably in human cells, that encode KLK-L Related Proteins. The nucleotide probes may also be useful in the diagnosis of cancer; in monitoring the progression of cancer; or monitoring a therapeutic treatment.

The probe may be used in hybridization techniques to detect genes that encode KLK-L Related Proteins. The technique generally involves contacting and incubating nucleic acids (e.g. recombinant DNA molecules, cloned genes) obtained from a sample from a patient or other cellular source with a probe of the present invention under conditions favorable for the specific annealing of the probes to complementary sequences in the nucleic acids. After incubation, the non-annealed nucleic acids are removed, and the presence of nucleic acids that have hybridized to the probe if any are detected.

The detection of nucleic acid molecules of the invention may involve the amplification of specific gene sequences using an amplification method such as PCR, followed by the analysis of the amplified molecules using techniques known to those skilled in the art. Suitable primers can be routinely designed

by one of skill in the art.

Genomic DNA may be used in hybridization or amplification assays of biological samples to detect abnormalities involving *klk-l* structure, including point mutations, insertions, deletions, and chromosomal rearrangements. For example, direct sequencing, single stranded conformational polymorphism analyses, heteroduplex analysis, denaturing gradient gel electrophoresis, chemical mismatch cleavage, and oligonucleotide hybridization may be utilized.

Genotyping techniques known to one skilled in the art can be used to type polymorphisms that are in close proximity to the mutations in a *klk-l* gene. The polymorphisms may be used to identify individuals in families that are likely to carry mutations. If a polymorphism exhibits linkage disequilibrium with mutations in a *klk-l* gene, it can also be used to screen for individuals in the general population likely to carry mutations. Polymorphisms which may be used include restriction fragment length polymorphisms (RFLPs), single-base polymorphisms, and simple sequence repeat polymorphisms (SSLPs).

A probe of the invention may be used to directly identify RFLPs. A probe or primer of the invention can additionally be used to isolate genomic clones such as YACs, BACs, PACs, cosmids, phage or plasmids. The DNA in the clones can be screened for SSLPs using hybridization or sequencing procedures.

Hybridization and amplification techniques described herein may be used to assay qualitative and quantitative aspects of *klk-l* expression. For example, RNA may be isolated from a cell type or tissue known to express *klk-l* and tested utilizing the hybridization (e.g. standard Northern analyses) or PCR techniques referred to herein. The techniques may be used to detect differences in transcript size which may be due to normal or abnormal alternative splicing. The techniques may be used to detect quantitative differences between levels of full length and/or alternatively splice transcripts detected in normal individuals relative to those individuals exhibiting cancer symptoms or other disease conditions.

The primers and probes may be used in the above described methods *in situ* i.e directly on tissue sections (fixed and/or frozen) of patient tissue obtained from biopsies or resections.

#### 4.1.2 Methods for Detecting KLK-L Related Proteins

Antibodies specifically reactive with a KLK-L Related Protein, or derivatives, such as enzyme conjugates or labeled derivatives, may be used to detect KLK-L Related Proteins in various samples (e.g. biological materials). They may be used as diagnostic or prognostic reagents and they may be used to detect abnormalities in the level of KLK-L Related Proteins expression, or abnormalities in the structure, and/or temporal, tissue, cellular, or subcellular location of a KLK-L Related Protein. Antibodies may also be used to screen potentially therapeutic compounds *in vitro* to determine their effects on cancer, and other conditions. *In vitro* immunoassays may also be used to assess or monitor the efficacy of particular therapies. The antibodies of the invention may also be used *in vitro* to determine the level of *KLK-L* expression in cells genetically engineered to produce a KLK-L Related Protein.

The antibodies may be used in any known immunoassays which rely on the binding interaction between an antigenic determinant of a KLK-L Related Protein and the antibodies. Examples of such assays are radioimmunoassays, enzyme immunoassays (e.g. ELISA), immunofluorescence, immunoprecipitation,

latex agglutination, hemagglutination, and histochemical tests. The antibodies may be used to detect and quantify KLK-L Related Proteins in a sample in order to determine its role in particular cellular events or pathological states, and to diagnose and treat such pathological states.

5 In particular, the antibodies of the invention may be used in immuno-histochemical analyses, for example, at the cellular and sub-subcellular level, to detect a KLK-L Related Protein, to localize it to particular cells and tissues, and to specific subcellular locations, and to quantitate the level of expression.

Cytochemical techniques known in the art for localizing antigens using light and electron microscopy may be used to detect a KLK-L Related Protein. Generally, an antibody of the invention may be labeled with a detectable substance and a KLK-L Related Protein may be localised in tissues and cells  
10 based upon the presence of the detectable substance. Examples of detectable substances include, but are not limited to, the following: radioisotopes (e.g.,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{35}\text{S}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), luminescent labels such as luminol; enzymatic labels (e.g., horseradish peroxidase, beta-galactosidase, luciferase, alkaline phosphatase, acetylcholinesterase), biotinyl groups (which can be detected by marked avidin e.g., streptavidin containing a fluorescent marker or enzymatic  
15 activity that can be detected by optical or calorimetric methods), predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached via spacer arms of various lengths to reduce potential steric hindrance. Antibodies may also be coupled to electron dense substances, such as ferritin or colloidal gold, which are readily visualised by electron microscopy.

20 The antibody or sample may be immobilized on a carrier or solid support which is capable of immobilizing cells, antibodies etc. For example, the carrier or support may be nitrocellulose, or glass, polyacrylamides, gabbros, and magnetite. The support material may have any possible configuration including spherical (e.g. bead), cylindrical (e.g. inside surface of a test tube or well, or the external surface of a rod), or flat (e.g. sheet, test strip). Indirect methods may also be employed in which the primary  
25 antigen-antibody reaction is amplified by the introduction of a second antibody, having specificity for the antibody reactive against KLK-L Related Protein. By way of example, if the antibody having specificity against a KLK-L Related Protein is a rabbit IgG antibody, the second antibody may be goat anti-rabbit gamma-globulin labeled with a detectable substance as described herein.

Where a radioactive label is used as a detectable substance, a KLK-L Related Protein may be  
30 localized by radioautography. The results of radioautography may be quantitated by determining the density of particles in the radioautographs by various optical methods, or by counting the grains.

#### 4.2 Methods for Identifying or Evaluating Substances/Compounds

The methods described herein are designed to identify substances that modulate the biological activity of a KLK-L Related Protein including substances that bind to KLK-L Related Proteins, or bind to  
35 other proteins that interact with a KLK-L Related Protein, to compounds that interfere with, or enhance the interaction of a KLK-L Related Protein and substances that bind to the KLK-L Related Protein or other proteins that interact with a KLK-L Related Protein. Methods are also utilized that identify compounds that bind to *KLK-L* regulatory sequences.

The substances and compounds identified using the methods of the invention include but are not limited to peptides such as soluble peptides including Ig-tailed fusion peptides, members of random peptide libraries and combinatorial chemistry-derived molecular libraries made of D- and/or L-configuration amino acids, phosphopeptides (including members of random or partially degenerate, directed phosphopeptide libraries), antibodies [e.g. polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, single chain antibodies, fragments, (e.g. Fab, F(ab)<sub>2</sub>, and Fab expression library fragments, and epitope-binding fragments thereof)], and small organic or inorganic molecules. The substance or compound may be an endogenous physiological compound or it may be a natural or synthetic compound.

Substances which modulate a KLK-L Related Protein can be identified based on their ability to bind to a KLK-L Related Protein. Therefore, the invention also provides methods for identifying substances which bind to a KLK-L Related Protein. Substances identified using the methods of the invention may be isolated, cloned and sequenced using conventional techniques. A substance that associates with a polypeptide of the invention may be an agonist or antagonist of the biological or immunological activity of a polypeptide of the invention.

The term "agonist", refers to a molecule that increases the amount of, or prolongs the duration of, the activity of the polypeptide. The term "antagonist" refers to a molecule which decreases the biological or immunological activity of the polypeptide. Agonists and antagonists may include proteins, nucleic acids, carbohydrates, or any other molecules that associate with a polypeptide of the invention.

Substances which can bind with a KLK-L Related Protein may be identified by reacting a KLK-L Related Protein with a test substance which potentially binds to a KLK-L Related Protein, under conditions which permit the formation of substance-KLK-L Related Protein complexes and removing and/or detecting the complexes. The complexes can be detected by assaying for substance-KLK-L Related Protein complexes, for free substance, or for non-complexed KLK-L Related Protein. Conditions which permit the formation of substance-KLK-L Related Protein complexes may be selected having regard to factors such as the nature and amounts of the substance and the protein.

The substance-protein complex, free substance or non-complexed proteins may be isolated by conventional isolation techniques, for example, salting out, chromatography, electrophoresis, gel filtration, fractionation, absorption, polyacrylamide gel electrophoresis, agglutination, or combinations thereof. To facilitate the assay of the components, antibody against KLK-L Related Protein or the substance, or labeled KLK-L Related Protein, or a labeled substance may be utilized. The antibodies, proteins, or substances may be labeled with a detectable substance as described above.

A KLK-L Related Protein, or the substance used in the method of the invention may be insolubilized. For example, a KLK-L Related Protein, or substance may be bound to a suitable carrier such as agarose, cellulose, dextran, Sephadex, Sepharose, carboxymethyl cellulose polystyrene, filter paper, ion-exchange resin, plastic film, plastic tube, glass beads, polyamine-methyl vinyl-ether-maleic acid copolymer, amino acid copolymer, ethylene-maleic acid copolymer, nylon, silk, etc. The carrier may be in the shape of, for example, a tube, test plate, beads, disc, sphere etc. The insolubilized protein or substance may be prepared by reacting the material with a suitable insoluble carrier using known chemical or physical

methods, for example, cyanogen bromide coupling.

The invention also contemplates a method for evaluating a compound for its ability to modulate the biological activity of a KLK-L Related Protein of the invention, by assaying for an agonist or antagonist (i.e. enhancer or inhibitor) of the binding of a KLK-L Related Protein with a substance which binds with a KLK-L Related Protein. The basic method for evaluating if a compound is an agonist or antagonist of the binding of a KLK-L Related Protein and a substance that binds to the protein, is to prepare a reaction mixture containing the KLK-L Related Protein and the substance under conditions which permit the formation of substance-KLK-L Related Protein complexes, in the presence of a test compound. The test compound may be initially added to the mixture, or may be added subsequent to the addition of the KLK-L Related Protein and substance. Control reaction mixtures without the test compound or with a placebo are also prepared. The formation of complexes is detected and the formation of complexes in the control reaction but not in the reaction mixture indicates that the test compound interferes with the interaction of the KLK-L Related Protein and substance. The reactions may be carried out in the liquid phase or the KLK-L Related Protein, substance, or test compound may be immobilized as described herein. The ability of a compound to modulate the biological activity of a KLK-L Related Protein of the invention may be tested by determining the biological effects on cells.

It will be understood that the agonists and antagonists i.e. inhibitors and enhancers that can be assayed using the methods of the invention may act on one or more of the binding sites on the protein or substance including agonist binding sites, competitive antagonist binding sites, non-competitive antagonist binding sites or allosteric sites.

The invention also makes it possible to screen for antagonists that inhibit the effects of an agonist of the interaction of KLK-L Related Protein with a substance which is capable of binding to the KLK-L Related Protein. Thus, the invention may be used to assay for a compound that competes for the same binding site of a KLK-L Related Protein.

The invention also contemplates methods for identifying compounds that bind to proteins that interact with a KLK-L Related Protein. Protein-protein interactions may be identified using conventional methods such as co-immunoprecipitation, crosslinking and co-purification through gradients or chromatographic columns. Methods may also be employed that result in the simultaneous identification of genes which encode proteins interacting with a KLK-L Related Protein. These methods include probing expression libraries with labeled KLK-L Related Protein.

Two-hybrid systems may also be used to detect protein interactions *in vivo*. Generally, plasmids are constructed that encode two hybrid proteins. A first hybrid protein consists of the DNA-binding domain of a transcription activator protein fused to a KLK-L Related Protein, and the second hybrid protein consists of the transcription activator protein's activator domain fused to an unknown protein encoded by a cDNA which has been recombined into the plasmid as part of a cDNA library. The plasmids are transformed into a strain of yeast (e.g. *S. cerevisiae*) that contains a reporter gene (e.g. lacZ, luciferase, alkaline phosphatase, horseradish peroxidase) whose regulatory region contains the transcription activator's binding site. The hybrid proteins alone cannot activate the transcription of the reporter gene. However,

interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product.

It will be appreciated that fusion proteins may be used in the above-described methods. In particular, KLK-L Related Proteins fused to a glutathione-S-transferase may be used in the methods.

5 The reagents suitable for applying the methods of the invention to evaluate compounds that modulate a KLK-L Related Protein may be packaged into convenient kits providing the necessary materials packaged into suitable containers. The kits may also include suitable supports useful in performing the methods of the invention.

#### 4.3 Compositions and Treatments

10 The proteins of the invention, substances or compounds identified by the methods described herein, antibodies, and antisense nucleic acid molecules of the invention may be used for modulating the biological activity of a KLK-L Related Protein, and they may be used in the treatment of conditions such as cancer (e.g. prostate, testicular, brain, uterine, thymus, ovarian, colon, ovarian, or breast cancer). Accordingly, the substances, antibodies, peptides, and compounds may be formulated into pharmaceutical  
15 compositions for administration to subjects in a biologically compatible form suitable for administration *in vivo*. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the active substance to be administered in which any toxic effects are outweighed by the therapeutic effects. The active substances may be administered to living organisms including humans, and animals. Administration of a therapeutically active amount of a pharmaceutical composition of the present invention is defined as  
20 an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a substance may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of antibody to elicit a desired response in the individual. Dosage regima may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies  
25 of the therapeutic situation.

The active substance may be administered in a convenient manner such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the active substance may be coated in a material to protect the substance from the action of enzymes, acids and other natural conditions that may inactivate  
30 the substance.

The compositions described herein can be prepared by per se known methods for the preparation of pharmaceutically acceptable compositions which can be administered to subjects, such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's  
35 Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985). On this basis, the compositions include, albeit not exclusively, solutions of the active substances in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffered solutions with a suitable pH and iso-osmotic with the physiological fluids.

Based upon their homology to genes encoding kallikrein, nucleic acid molecules of the invention may be also useful in the treatment of conditions such as hypertension, cardiac hypertrophy, arthritis, inflammatory disorders, neurological disorders, and blood clotting disorders.

5 Vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids, may be used to deliver nucleic acid molecules to a targeted organ, tissue, or cell population. Methods well known to those skilled in the art may be used to construct recombinant vectors which will express antisense nucleic acid molecules of the invention. (See, for example, the techniques described in Sambrook et al (supra) and Ausubel et al (supra)).

10 The nucleic acid molecules comprising full length cDNA sequences and/or their regulatory elements enable a skilled artisan to use sequences encoding a protein of the invention as an investigative tool in sense (Yousoufian H and H F Lodish 1993 Mol Cell Biol 13:98-104) or antisense (Eguchi et al (1991) Annu Rev Biochem 60:631-652) regulation of gene function. Such technology is well known in the art, and sense or antisense oligomers, or larger fragments, can be designed from various locations along the coding or control regions.

15 Genes encoding a protein of the invention can be turned off by transfecting a cell or tissue with vectors which express high levels of a desired KLK-L-encoding fragment. Such constructs can inundate cells with untranslatable sense or antisense sequences. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until all copies are disabled by endogenous nucleases.

20 Modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA or PNA, to the regulatory regions of a gene encoding a protein of the invention, ie, the promoters, enhancers, and introns. Preferably, oligonucleotides are derived from the transcription initiation site, eg, between -10 and +10 regions of the leader sequence. The antisense molecules may also be designed so that they block translation of mRNA by preventing the transcript from binding to ribosomes. Inhibition may also  
25 be achieved using "triple helix" base-pairing methodology. Triple helix pairing compromises the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Therapeutic advances using triplex DNA were reviewed by Gee J E et al (In: Huber B E and B I Carr (1994) Molecular and Immunologic Approaches, Futura Publishing Co, Mt Kisco N.Y.).

30 Ribozymes are enzymatic RNA molecules that catalyze the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. The invention therefore contemplates engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding a protein of the invention.

35 Specific ribozyme cleavage sites within any potential RNA target may initially be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC. Once the sites are identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets

may also be determined by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

5 Methods for introducing vectors into cells or tissues include those methods discussed herein and which are suitable for *in vivo*, *in vitro* and *ex vivo* therapy. For *ex vivo* therapy, vectors may be introduced into stem cells obtained from a patient and clonally propagated for autologous transplant into the same patient (See U.S. Pat. Nos. 5,399,493 and 5,437,994). Delivery by transfection and by liposome are well known in the art.

10 The nucleic acid molecules disclosed herein may also be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including but not limited to such properties as the triplet genetic code and specific base pair interactions.

15 The invention also provides methods for studying the function of a polypeptide of the invention. Cells, tissues, and non-human animals lacking in expression or partially lacking in expression of a nucleic acid molecule or gene of the invention may be developed using recombinant expression vectors of the invention having specific deletion or insertion mutations in the gene. A recombinant expression vector may be used to inactivate or alter the endogenous gene by homologous recombination, and thereby create a deficient cell, tissue, or animal.

20 Null alleles may be generated in cells, such as embryonic stem cells by deletion mutation. A recombinant gene may also be engineered to contain an insertion mutation that inactivates the gene. Such a construct may then be introduced into a cell, such as an embryonic stem cell, by a technique such as transfection, electroporation, injection etc. Cells lacking an intact gene may then be identified, for example by Southern blotting, Northern Blotting, or by assaying for expression of the encoded polypeptide using the methods described herein. Such cells may then be fused to embryonic stem cells to generate transgenic non-human animals deficient in a polypeptide of the invention. Germline transmission of the mutation may be achieved, for example, by aggregating the embryonic stem cells with early stage embryos, such as 8 cell embryos, *in vitro*; transferring the resulting blastocysts into recipient females and; generating germline transmission of the resulting aggregation chimeras. Such a mutant animal may be used to define specific cell populations, developmental patterns and *in vivo* processes, normally dependent on gene expression.

25 The invention thus provides a transgenic non-human mammal all of whose germ cells and somatic cells contain a recombinant expression vector that inactivates or alters a gene encoding a KLK-L Related Protein. In an embodiment the invention provides a transgenic non-human mammal all of whose germ cells and somatic cells contain a recombinant expression vector that inactivates or alters a gene encoding a KLK-L Related Protein resulting in a KLK-L Related Protein associated pathology. Further the invention provides a transgenic non-human mammal which does not express a KLK-L Related Protein of the invention. 30 In an embodiment, the invention provides a transgenic non-human mammal which does not express a KLK-L Related Protein of the invention resulting in a KLK-L Related Protein associated pathology. A KLK-L Related Protein pathology refers to a phenotype observed for a KLK-L Related Protein homozygous mutant. 35

A transgenic non-human animal includes but is not limited to mouse, rat, rabbit, sheep, hamster, dog, guinea pig, micro-pig, pig, cat, goat, and non-human primates, preferably mouse.

The invention also provides a transgenic non-human animal assay system which provides a model system for testing for an agent that reduces or inhibits a pathology associated with an KLK-L Related Protein, preferably a KLK-L Related Protein associated pathology, comprising:

- (a) administering the agent to a transgenic non-human animal of the invention; and
- (b) determining whether said agent reduces or inhibits the pathology (e.g. KLK-L Related Protein associated pathology) in the transgenic non-human animal relative to a transgenic non-human animal of step (a) which has not been administered the agent.

The agent may be useful in the treatment and prophylaxis of conditions such as cancer as discussed herein. The agents may also be incorporated in a pharmaceutical composition as described herein.

The activity of the proteins, substances, compounds, antibodies, nucleic acid molecules, agents, and compositions of the invention may be confirmed in animal experimental model systems. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The therapeutic index is the dose ratio of therapeutic to toxic effects and it can be expressed as the  $ED_{50}/LD_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred.

The following non-limiting examples are illustrative of the present invention:

## Examples

### Example 1

#### MATERIALS AND METHODS

##### Identification of positive PAC and BAC genomic clones from a human genomic DNA library

The sequence of PSA, KLK1, KLK2, NES1 and Zyme genes is already known. Polymerase chain reaction (PCR)-based amplification protocols have been developed which allowed generation of PCR products specific for each one of these genes. Using these PCR products as probes, labeled with  $^{32}P$ , a human genomic DNA PAC library and a human genomic DNA BAC library was screened for the purpose of identifying positive clones of approximately 100-150 Kb long. The general strategies for these experiments have been published elsewhere (14). The genomic libraries were spotted in duplicate on nylon membranes and positive clones were further confirmed by Southern blot analysis as described (14).

##### DNA sequences on chromosome 19

The Lawrence Livermore National Laboratory participates in the sequencing of the human genome project and focuses on sequencing chromosome 19. Large sequencing information on this chromosome is available at the website of the Lawrence Livermore National Laboratory (<http://www-bio.llnl.gov/genome/gemnome.html>).

Approximately 300 Kb of genomic sequences were obtained from that website, encompassing a region on chromosome 19q13.3 - 13.4, where the known kallikrein genes are localized. This 300 Kb of sequence is represented by 8 contigs of variable lengths. By using a number of different computer

programs, an almost contiguous sequence of the region was established as shown diagrammatically in Figure 1 and Figure 28. Some of the contigs were reversed as shown in Figure 1 in order to reconstruct the area on both strands of DNA.

5 By using the published sequences of PSA, KLK2, NES1 and Zyme and the computer software BLAST 2, using alignment strategies, the relative positions of these genes on the contiguous map were identified (Figure 28). These known genes served as hallmarks for further studies. An EcoR1 restriction map of the area is also available at the website of the Lawrence Livermore National Laboratory. Using this restriction map and the computer program WebCutter (<http://www.firstmarket.com/cutter/cut2.html>), a restriction study analysis of the available sequence was performed to further confirm the assignment and  
10 relative positions of these contigs along chromosome 19. The obtained configuration and the relative location of the known genes are presented in Figure 1.

#### Gene prediction analysis

For exon prediction analysis of the whole genomic area, a number of different computer programs were used. All the programs were initially tested using known genomic sequences of the PSA, Zyme, and  
15 NES1 genes. The more reliable computer programs, GeneBuilder (gene prediction), GeneBuilder (exon prediction), Grail 2 and GENEID-3 were selected for further use.

#### Protein homology searching

Putative exons of the new genes were first translated to the corresponding amino acid sequences. BLAST homology searching for the proteins encoded by the exons of the putative new genes were  
20 performed using the BLASTP program and the Genbank databases.

### RESULTS

#### Relative position of PSA, KLK2, Zyme and NES1 on Chromosome 19

Screening of the human BAC library identified two clones which were positive for the Zyme gene (clones BAC 288H1 and BAC 76F7). These BACs were further analyzed by PCR and primers specific for  
25 PSA, NES1, KLK1 and KLK2. These analyses indicated that both BACs were positive for Zyme, PSA and KLK2 and negative for KLK1 and NES1 genes.

Screening of the human PAC genomic library identified a PAC clone which was positive for NES1 (clone PAC 34B1). Further PCR analysis indicated that this PAC clone was positive for NES1 and KLK1 genes and negative for PSA, KLK2 and Zyme. Combination of this information with the EcoR1  
30 restriction map of the region allowed establishment of the relative positions of these four genes. PSA is the most centromeric, followed by KLK2, Zyme and NES1. Further alignment of the known sequences of these genes with the 300 Kb contig enabled precise localization of all four genes and determination of the direction of transcription, as shown by the arrows in Figure 1. The KLK1 gene sequence was not identified on any of these contigs and appears to be further telomeric to NES1 (since it is co-localized on the same  
35 PAC as NES1).

#### Identification of new genes

A set of rules was used to consider the presence of a new gene in the genomic area of interest as follows:

1. Clusters of at least 3 exons should be found.
2. Only exons with high prediction score ("good" or "excellent" quality, as indicated by the searching programs) were considered for the construction of the putative new genes.
3. Exons predicted were reliable only if they were identified by at least two different exon prediction programs.

By using this strategy, eleven putative new genes were identified of which three were found on subsequent homology analysis to be known genes not previously mapped i.e. the human stratum corneum chymotrypsin enzyme (HSCCE), human neuropsin, and trypsin-like serine protease (TLSP). Their relative location is shown in Figure 1. The five genes all have variable homologies with known human or animal kallikrein proteins and/or other known serine proteases (depicted as KLK-L1, KLK-L2, KLK-L3, KLK-L4 and KLK-L5 in Figure 1 and KLK-L1 to KLK-L6 in Figure 28).

In Tables 1 to 5, the preliminary exon structure and partial protein sequence for each one of the newly identified genes is shown. In Table 6, some proteins are presented which appear, on preliminary analysis, to be homologous to the proteins encoded by the putative new genes. SEQ. ID. NOs. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, and 67 show amino acid sequences of KLK-L1-KLK-L6, and SEQ. ID. NOs. 1, 13, 21, 43, 56, and 65 show nucleic acid sequences of the genes encoding KLK-L1- KLK-L6.

## DISCUSSION

Prediction of protein-coding genes in newly sequenced DNA becomes very important after the establishment of large genome sequencing projects. This problem is complicated due to the exon-intron structure of the eukaryotic genes which interrupts the coding sequence in many unequal parts. In order to predict the protein-coding exons and overall gene structure, a number of computer programs were developed. All these programs are based on the combination of potential functional signals with the global statistical properties of known protein-coding regions (15). However, the most powerful approach for gene structure prediction is to combine information about potential functional signals (splice sites, translation start or stop signal etc.) together with the statistical properties of coding sequences (coding potential) along with information about homologies between the predicted protein and known protein families (16).

In mouse and rat, kallikreins are encoded by large multigene families and these genes tend to cluster in groups with a distance as small as 3.3 – 7.0 Kb (3). A strong conservation of gene order between human chromosome 19q13.1 – q13.4 and 17 loci in a 20-cM proximal part of mouse chromosome 7, including the kallikrein locus, has been documented (17).

In humans, only a few kallikrein genes were identified. In fact, only KLK1, KLK2 and KLK3 (PSA) are considered to represent the human kallikrein gene family (9). The work described herein provides strong evidence that a large number of kallikrein-like genes are clustered within a 300Kb region around chromosome 19q13.2 – q13.4. The three established human kallikreins (KLK1, KLK2, KLK3), Zyme and NES1, as well as the stratum corneum chymotryptic enzyme, neuropsin, and TLSP (trypsin-like serine protease) and another five new genes, KLK-L1 to KLK-L5, may constitute a large gene family. This will bring the total number of kallikrein or kallikrein-like genes in this region of chromosome 19 to thirteen.

The human stratum corneum chymotryptic enzyme (19), neuropsin (20) and trypsin-like serine

protease (TLSP) (21) are three previously characterized genes which have many structural similarities with the kallikreins and other members of the serine protease family. However, they have not been mapped in the past. Their precise mapping in the region of the kallikrein gene family indicates that these three genes, along with the ones that were newly identified, or are already known, constitute a family that likely originated by duplication of an ancestral gene. The relative localization of all these genes is depicted in Figure 1.

Kallikrein genes are a subfamily of serine proteases, traditionally characterized by their ability to liberate lysyl-bradykinin (kallidin) from kininogen (18). More recently, however, a new, structural concept has emerged to describe kallikreins. From accumulated sequence data, it is now clear that the mouse has many genes with high homology to kallikrein coding sequences (19-20). Richard and co-workers have contributed to the concept of a "kallikrein multigene family" to refer to these genes (21-22). This definition is not based much on specific enzymatic function of the gene product, but more on its sequence homology and their close linkage on mouse chromosome 7. In humans, only KLK1 meets the functional definition of a kallikrein. KLK2 has trypsin-like enzymatic activity and KLK3 (PSA) has very weak chymotrypsin-like enzymatic activity. These activities of KLK2 and KLK3 are not known to liberate biologically active peptides from precursors. Based on the newer definition, members of the kallikrein family include, not only the gene for the kallikrein enzyme, but also genes encoding other homologous proteases, including the enzyme that processes the precursors of the nerve growth factor and epidermal growth factor (8). Therefore, it is important to note the clear distinction between the enzyme kallikrein and a kallikrein or a kallikrein-like gene.

In carrying out the study only exons were considered which were predicted with "good" or "excellent" quality and only exons were considered which were predicted by at least two different programs. Moreover, the presence of a putative gene was only considered when at least three exons clustered coordinately in that region. Additional evidence that these new genes are indeed homologous to the known kallikreins and other serine proteases comes from comparison of the intron phases. As published previously (14), trypsinogen, PSA and NES1 have 5 coding exons of which the first has intron phase I (the intron occurs after the first nucleotide of the codon), the second has intron phase II (the intron occurs after the second nucleotide and the codon), the third has intron phase I and the fourth has intron phase 0 (the intron occurs between codons). The fifth exon contains the stop codon. The intron phases of the predicted new kallikrein-like genes follow these rules and are shown in the respective tables. Further support comes from the identification in the new genes, of the conserved amino acids of the catalytic domain of the serine proteases, as presented in Tables 1 - 5.

In order to test the accuracy of the computer programs, known genomic areas containing the PSA, Zyme and KLK2 genes were tested. Two of these programs (Grail 2 and GeneBuilder) were able to detect about 95% of the tested known genes. Matches with expressed sequence tag sequences (EST) can also be employed for gene structure prediction in the GeneBuilder program and this can significantly improve the power of the program especially at high stringency (e.g. >95% homology).

In mouse, ten of the kallikrein genes appear to be pseudogenes (9).

## Example 2

### PROSTASE/KLK-L1 in prostate and breast tissues

The fine mapping of the prostase/KLK-L1 gene and its chromosomal localization in relation to a number of other homologous genes also mapping to the same region are described. In addition, extensive tissue expression studies were carried out that demonstrate that, in addition to prostate (which shows the highest expression), that prostase/KLK-L1 is also expressed in female breasts, testis, adrenals, uterus, colon, thyroid, brain, spinal cord and salivary glands. Furthermore, the gene is up-regulated by androgens and progestins in the breast carcinoma cell line BT-474.

### Materials and Methods

#### 10 DNA sequences on chromosome 19

Large DNA sequencing data for chromosome 19 is available at the web site of the Lawrence Livermore National Laboratory (LLNL) (<http://www-bio.llnl.gov/genome/genome.html>). Approximately 300 Kb of genomic sequence was obtained from that web site, encompassing a region on chromosome 19q13.3 - 13.4, where the known kallikrein genes are localized. This sequence is represented by 9 contigs of variable lengths. By using the sequences of PSA, KLK2, NES1 and protease M and the alignment program BLAST 2 (37), the relative positions of these genes on the contiguous map were located.

### Gene prediction analysis

For exon prediction analysis of the whole genomic area, a number of different computer programs were used. Originally all these programs were tested using the known genomic sequences of the PSA, protease M and NES1 genes. The most reliable computer programs GeneBuilder (gene prediction) [<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html> ] GeneBuilder (exon prediction) [<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html> ], Grail 2 [<http://compbio.ornl.gov> ], and GENEID-3 [<http://apollo.imim.es/geneid.html>] were selected for further use.

### Protein homology searching

Putative exons of the newly identified gene were first translated to the corresponding amino acid sequences. BLAST homology searching for the proteins encoded by the exons were performed using the BLASTP program and the GenBank databases (37).

### Searching expressed sequence tags (ESTs)

Sequence homology searching was performed using the BLASTN algorithm (37) on the National Center for Biotechnology Information web server (<http://www.ncbi.nlm.nih.gov/BLAST/>) against the human EST database (dbEST). Clones with > 95% homology were obtained from the I.M.A.G.E. (38) consortium through Research Genetics Inc, Huntsville, AL and from The Institute for Genomic Research (TIGR) (<http://WWW.TIGR.ORG/tdb/tdb.html>) (Table 7). Clones were propagated, purified and then sequenced from both directions with an automated sequencer, using insert-flanking vector primers.

#### 35 Breast cancer cell line and stimulation experiments

The breast cancer cell line BT-474 was purchased from the American Type Culture Collection (ATCC), Rockville, MD. BT-474 cells were cultured in RPMI media (Gibco BRL, Gaithersburg, MD) supplemented with glutamine (200 mmol/L), bovine insulin (10 mg/L), fetal bovine serum (10%),

antibiotics and antimycotics, in plastic flasks, to near confluency. The cells were then aliquoted into 24-well tissue culture plates and cultured to 50% confluency. 24 hours before the experiments, the culture media were changed into phenol red-free media containing 10% charcoal-stripped fetal bovine serum. For stimulation experiments, various steroid hormones dissolved in 100% ethanol were added into the culture media, at a final concentration of  $10^{-8}$  M. Cells stimulated with 100% ethanol were included as controls. The cells were cultured for 24 hours, then harvested for mRNA extraction.

#### **Reverse transcriptase polymerase chain reaction**

Total RNA was extracted from the breast cancer cells using Trizol reagent (Gibco BRL) following the manufacturer's instructions. RNA concentration was determined spectrophotometrically. 2 µg of total RNA was reverse transcribed into first strand cDNA using the Superscript<sup>TM</sup> preamplification system (Gibco BRL). The final volume was 20 µl. Based on the combined information obtained from the predicted genomic structure of the new gene and the EST sequences, two gene-specific primers were designed (Table 8), PCR was carried out in a reaction mixture containing 1 µl of cDNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200 µM dNTPs (deoxynucleoside triphosphates), 150 ng of primers and 2.5 units of AmpliTaq Gold DNA polymerase (Roche Molecular Systems, Branchburg, NJ, USA) on a Perkin-Elmer 9600 thermal cycler. The cycling conditions were 94°C for 9 minutes to activate the Taq Gold DNA polymerase, followed by 43 cycles of 94°C for 30 s, 63°C for 1 minute and a final extension at 63°C for 10 min. Equal amounts of PCR products were electrophoresed on 2% agarose gels and visualized by ethidium bromide staining. All primers for RT-PCR spanned at least 2 exons to avoid contamination by genomic DNA.

#### **Tissue expression of KLK-L1**

Total RNA isolated from 26 different human tissues was purchased from Clontech, Palo Alto, CA. cDNA was prepared as described above for the tissue culture experiments and used for PCR reactions with the primers described in Table 8 (SEQ. ID. Nos 5-12). Tissue cDNAs were amplified at various dilutions.

#### **Cloning and sequencing of the PCR products**

To verify the identity of the PCR products, they were cloned into the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The inserts were sequenced from both directions using vector-specific primers, by an automated DNA sequencer.

### **Results**

#### **Identification of the prostase/KLK-L1 gene**

The exon prediction strategy of the 300Kb DNA sequences around chromosome 19q13.3 - q13.4 identified a novel gene with a structure reminiscent of a serine protease. The major features of this gene were its homology, at the amino acid and DNA level, with other human kallikrein genes; the conservation of the catalytic triad (histidine, aspartic acid, and serine), the number of exons and the complete conservation of the intron phases.

#### **EST sequence homology search**

EST sequence homology search of the putative exons obtained from the gene prediction programs (as described above) against the human EST database (dbEST) revealed five expressed sequence tags

(ESTs) with >95 % identity to the putative exons of the gene (Table 7). Positive clones were obtained and the inserts were sequenced from both directions. Alignment was used to compare between the EST sequences and the exons predicted by the computer programs, and final selection of the exon-intron splice sites was made according to the EST sequences. Furthermore, many of the ESTs were overlapping, further ensuring the accuracy of the data.

The coding sequence of the klk-L2 gene is shown in SEQ. ID. NO. 1 and GenBank Accession # AF135023. The exons of the gene are as follows: exon 1 (939-999); exon 2 (2263-2425); exon 3 (2847-3097); exon 4 (3181-3317); and exon 5 (4588-4740). The amino acid sequence of KLK-L2 proteins are shown in SEQ. ID. Nos. 2 and 3.

#### 10 Mapping and chromosomal localization of prostate/KLK-L1 gene

Alignment of the prostate/KLK-L1 sequence and the sequences of other known kallikrein genes within the 300 Kb area of the contigs constructed at the Lawrence Livermore National Laboratory enabled precise localization of all genes and to determine the direction of transcription, as shown in Figure 2. The distance between PSA and KLK2 genes was calculated to be 12,508 bp. The prostate/KLK-L1 gene is 15 26,229 bp more telomeric to KLK2 and transcribes in the opposite direction. The zyme gene is about 51 Kb more telomeric to the prostate gene and transcribes in the same direction. The human stratum corneum chymotryptic enzyme gene, the neuropsin gene and the NES 1 gene are all further telomeric to zyme and all transcribe in the same direction as zyme.

#### Tissue expression of the prostate/KLK-L1 gene

20 The tissues that express the prostate/KLK-L1 gene were assessed by RT-PCR. The experiments were performed at various dilutions of the cDNAs to obtain some information about the relative levels of expression. RT-PCR for actin was used as a positive control and RT-PCR for the PSA cDNA was used as another positive control with tissue restricted specificity. Positive ESTs for prostate/KLK-L1 were used as controls for the PCR procedure. The PSA gene was found to be highly expressed in the prostate, as 25 expected, and to a lower extent in mammary and salivary glands as also expected from recent literature reports (24, 25). Very low expression of PSA in the thyroid gland, trachea and testis was also found, a finding that accords with recent RT-PCR data by others (26).

The tissue expression of prostate/KLK-L1 is summarized in Table 9 and Figure 3. This protease is primarily expressed in the prostate, testis, adrenals, uterus, thyroid, colon, central nervous system and 30 mammary tissues, and, at much lower levels in other tissues. The specificity of the RT-PCR procedure was verified for prostate/KLK-L1 by cloning the PCR products from mammary, testicular and prostate tissues and sequencing them. One example with mammary tissue is shown in Figure 4. All cloned PCR products were identical in sequence to the cDNA sequence reported for the prostate/KLK-L1.

#### Hormonal regulation of the prostate/KLK-L1 gene

35 The steroid hormone receptor-positive breast carcinoma cell line BT-474 was used as a model system to evaluate whether prostate/KLK-L1 expression is under steroid hormone regulation. As shown in Figure 5, the controls worked as expected i. e., actin positivity without hormonal regulation in all cDNAs, only estrogen up-regulation of the pS2 gene and up-regulation of the PSA gene by androgens and

progestins. Prostase/KLK-L1 is up-regulated primarily by androgens and progestins, similarly to PSA. This up-regulation was dose-dependent and it was evident at steroid hormone levels  $\geq 10^{-10}$  M.

## DISCUSSION

5 The KLK3 gene encodes for PSA, a protein that currently represents the best tumor marker available (24). Since in rodents there are so many kallikrein genes, the restriction of this family to only 3 genes in humans was somewhat surprising. More recently, new candidate kallikrein genes in humans have been discovered, including NES1 (13) and zyme/protease M/neurosin (10-12). The known kallikreins and the newly discovered kallikrein-like genes share the following similarities: (a) they encode serine proteases (b) they have five coding exons (c) they share significant DNA and protein homologies with each other (d) 10 they map in the same locus on chromosome 19q13.3-q13.4, a region that is structurally similar to an area on mouse chromosome 7, where all the mouse kallikrein genes are localized (e) they appear to be regulated by steroid hormones. Prostase/KLK-L1 is a member of the same family since these common characteristics are also shared by the newly discovered gene.

15 The exact localization of the KLK-L1 gene and its position in relation to other genes in the area (Figure 2) was determined. Prostase/KLK-L1 lies between KLK2 and zyme.

Irwin et al. (27) have proposed that the serine protease genes can be classified into five different groups according to intron position. The established kallikreins ( KLK1, KLK2, and PSA), trypsinogen and chymotrypsinogen belong to a group that has: (1) an intron just downstream from the codon for the active site histidine residue, (2) a second intron downstream from the exon containing the codon for the active site aspartic acid residue, and (3) a third intron just upstream from the exon containing the codon for the active site serine residue. As seen in Figure 6, the genomic organization of prostase/KLK-L1 gene is very similar to this group of genes. The lengths of the coding parts of exons 1-5 are 61,163, 263, 137 and 153 bp, respectively, which are close or identical to the lengths of the exons of the kallikrein genes and also, similar or identical to those of other newly discovered genes in the same chromosomal region like the 20 NES1(14), zyme/protease M/neurosin (10-12) and neuropsin (28) genes.

25 The sensitive RT-PCR protocol reveals that the KLK-L1 enzyme is expressed in prostatic tissue and it is also expressed in significant amounts in other tissues, including testis, female mammary gland, adrenals, uterus, thyroid, colon, brain, lung and salivary glands (Figure 3 and Table 9). The specificity of the RT-PCR primers was verified by sequencing the obtained PCR products, with one example shown in 30 Figure 4 (SEQ.ID.NO. 4). Tissue culture studies with the breast carcinoma cell line BT-474 further confirm not only the ability of these cells to produce prostase/KLK-L1 but also its hormonal regulation (Figure 5).

35 An interesting theme is now developing involving the group of homologous genes on chromosome 19q13.3(PSA, KLK2, prostase, zyme, and NES1). The combined data suggest that all of them are expressed in prostate and breast tissues, and all of them are hormonally regulated. All these genes may be part of a cascade pathway that plays a role in cell proliferation, differentiation or apoptosis by regulating (positively or negatively) growth factors or their receptors or cytokines, through proteolysis (30). Also interesting is the linkage of locus 19q13 to solid tumors and gliomas (31) which raises the possibility that some of the genes in the region may be disrupted by rearrangements.

The KLK-1L gene encodes for a serine protease that shows homology with other members of the kallikrein gene family and maps to the same chromosomal location. Many structural features of the kallikreins are conserved in prostase/KLK-L1. The precise mapping of this gene between the two known genes KLK2 and zyme is presented. It is further demonstrated that prostase/KLK-L1 is expressed in many tissues, in addition to the prostate, including the female breast. This gene is also herein referred to as 'prostase'. It has been further demonstrated, using breast carcinoma cell lines, that prostase/KLK-L1 can be produced by these cells and that its expression is significantly up-regulated by androgens and progestins. Based on information for other homologous genes in the area (PSA, zyme, and NES1), prostase/KLK-L1 may be involved in the pathogenesis and/or progression of prostate, breast and possibly other cancers.

### Example 3

#### IDENTIFICATION OF THE KLK-L2 GENE

##### Materials and Methods

##### DNA sequence on chromosome 19

Sequencing data of approximately 300Kb of nucleotides on chromosome 19q13.3-q13.4 was obtained from the web site of the Lawrence Livermore National Laboratory (LLNL) (<http://www-bio.llnl.gov/genome/genome.html>). This sequence was in the form of 9 contigs of different lengths. A restriction analysis study of the available sequences was performed using the "WebCutter" computer program (<http://www.firstmarket.com/cutter/cut2.html>) and with the aid of the EcoRI restriction map of this area (also available from the LLNL web site) an almost contiguous stretch of genomic sequences was constructed. The relative positions of the known kallikrein genes: PSA (GenBank accession # X14810), KLK2 (GenBank accession # M18157), and zyme (GenBank accession # U60801) was determined using the alignment program BLAST 2 (37).

##### New Gene Identification

A number of computer programs were used to predict the presence of putative new genes in the genomic area of interest. These programs were initially tested using the known genomic sequences of the PSA, protease M and NES1 genes. The most reliable computer programs GeneBuilder (gene prediction) (<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html>) GeneBuilder (exon prediction) (<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html>), Grail 2 (<http://compbio.ornl.gov>) and GENEID-3 (<http://apolo.imim.es/geneid.html>) were selected for further use.

##### Expressed sequence tag (EST) searching

The predicted exons of the putative new gene were subjected to homology search using the BLASTN algorithm (37) on the National Center for Biotechnology Information web server (<http://www.ncbi.nlm.nih.gov/BLAST/>) against the human EST database (dbEST). Clones with > 95% homology were obtained from the I.M.A.G.E. consortium (38) through Research Genetics Inc, Huntsville, AL (Table 10). The clones were propagated, purified and sequenced from both directions with an automated sequencer, using insert-flanking vector primers.

##### Rapid amplification of cDNA ends (5' RACE)

According to the EST sequence data and the predicted structure of the gene, two gene-specific primers were designed (R1 & R2) (Table 11). Two rounds of RACE reactions (nested PCR) were performed with 5µl Marathon Ready™ cDNA of human testis (Clontech, Palo Alto, CA, USA) as a template. The reaction mix and PCR conditions were conducted according to the manufacturer's recommendations. In brief, denaturation was done for 5 min at 94°C followed by 94° C for 5 sec followed by 72°C for 2 min for 5 cycles, then 94°C for 5 sec followed by 70° C for 2 min for 5 cycles then 94°C for 5 sec followed by 65°C for 2 min for 30 cycles for the first reaction and 25 cycles for the nested PCR reaction.

#### Tissue expression

Total RNA isolated from 26 different human tissues was purchased from Clontech, Palo Alto, CA. cDNA was prepared as described below for the tissue culture experiments and used for PCR reactions with the primers described in Table 11 (SEQ. ID. Nos 9-12, 15-20). Tissue cDNAs were amplified at various dilutions.

#### Breast cancer cell line and hormonal stimulation experiments

The breast cancer cell line BT-474 was purchased from the American Type Culture Collection (ATCC), Rockville, MD. Cells were cultured in RPMI media (Gibco BRL, Gaithersburg, MD) supplemented with glutamine (200 mmol/L), bovine insulin (10 mg/L), fetal bovine serum (10%), antibiotics and antimycotics, in plastic flasks, to near confluency. The cells were then aliquoted into 24-well tissue culture plates and cultured to 50% confluency. 24 hours before the experiments, the culture media were changed into phenol red-free media containing 10% charcoal-stripped fetal bovine serum. For stimulation experiments, various steroid hormones dissolved in 100% ethanol were added into the culture media, at a final concentration of  $10^{-8}$  M. Cells stimulated with 100% ethanol were included as controls. The cells were cultured for 24 hours, then harvested for mRNA extraction.

#### Reverse transcriptase polymerase chain reaction

Total RNA was extracted from the breast cancer cells using Trizol reagent (Gibco BRL) following the manufacturer's instructions. RNA concentration was determined spectrophotometrically. 2 µg of total RNA was reverse-transcribed into first strand cDNA using the Superscript™ preamplification system (Gibco BRL). The final volume was 20 µl. Based on the combined information obtained from the predicted genomic structure of the new gene and the EST sequences, two gene-specific primers were designed (Table 11) and PCR was carried out in a reaction mixture containing 1 µl of cDNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200 µM dNTPs (deoxynucleoside triphosphates), 150 ng of primers and 2.5 units of AmpliTaq Gold DNA polymerase (Roche Molecular Systems, Branchburg, NJ, USA) on a Perkin-Elmer 9600 thermal cycler. The cycling conditions were 94°C for 9 minutes to activate the Taq Gold DNA polymerase, followed by 43 cycles of 94°C for 30 s, 63°C for 1 minute and a final extension at 63°C for 10 min. Equal amounts of PCR products were electrophoresed on 2% agarose gels and visualized by ethidium bromide staining. All primers for RT-PCR spanned at least 2 exons to avoid contamination by genomic DNA.

To verify the identity of the PCR products, they were cloned into the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The inserts were sequenced from both directions using vector-specific primers, with an automated DNA sequencer.

#### Structure analysis

5 Multiple alignment was performed using the Clustal X software package available at: <ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/> (clustalx1.64b.msw.exe) and the multiple alignment program available from the Baylor College of Medicine (BCM), Houston, TX, USA ([kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html](http://kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html)). Phylogenetic studies were performed using the Phylip software package available at: <http://evolution.genetics.washington.edu/phylip/getme.html>. Distance matrix analysis  
10 was performed using the "Neighbor-Joining/UPGMA" program and parsimony analysis was done using the "Protpars" program. Hydrophobicity study was performed using the BCM search launcher programs (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>). Signal peptide was predicted using the "SignalP" server (<http://www.cbs.dtu.dk/services/signal>). Protein structure analysis was performed by "SAPS" (structural analysis of protein sequence) program (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>).  
15

#### RESULTS

Computer analysis of the genomic sequence predicted a putative new gene consisting of four exons. This gene was detected by all programs used and all exons had high prediction scores. EST sequence homology search of the putative exons against the human EST database (dbEST) revealed nine  
20 expressed sequence tag (EST) clones from different tissues with >95 % identity to the putative exons of the gene (Table 10). Positive clones were obtained and the inserts were sequenced from both directions. The "Blast 2 sequences" program was used to compare the EST sequences with the predicted exons, and final selection of the exon-intron splice sites was done according to the EST sequences. The presence of many areas of overlap between the various EST sequences allowed further verification of the structure of  
25 the new gene. The coding sequence of the gene is shown in SEQ. ID. NO. 13 and GenBank Accession #AF135028. The 3' end of the gene was verified by the presence of poly A stretches that are not present in the genomic sequence at the end of two of the sequenced ESTs. One of the sequenced ESTs revealed the presence of an additional exon at the 5' end. The nucleotide sequence of this exon matches exactly with the genomic sequence. To further identify the 5' end of the gene, 5' RACE was performed but no additional  
30 sequence could be obtained. However, as is the case with other kallikreins, the presence of further upstream untranslated exon(s) could not be excluded. The amino acid sequence of KLK-L2 is shown in SEQ. ID. No. 14.

#### Mapping and chromosomal localization of the KLK-L2 gene

Alignment of KLK-L2 gene and the sequences of other known kallikrein genes within the 300 Kb  
35 area of interest enabled precise localization of all genes and determination of the direction of transcription, as shown by the arrows in Figure 8. The PSA gene was found to be the most centromeric, separated by 12,508 base pairs (bp) from KLK2, and both genes are transcribed in the same direction (centromere to telomere). The prostase/KLK-L1 gene is 26,229 bp more telomeric and transcribes in the opposite

direction, followed by KLK-L2. The distance between KLK-L1 and KLK-L2 is about 35 Kilobases (Kb). The zyme gene is 5,981 bp more telomeric and the latter 3 genes are all transcribed in the same direction (Figure 8).

#### Structural characterization of the KLK-L2 gene and its protein product

5       The KLK-L2 gene, as presented in Figure 7, is formed of 5 coding exons and 4 intervening introns, spanning an area of 9,349 bp of genomic sequence on chromosome 19q13.3-q13.4. The lengths of the exons are 73, 262, 257, 134, and 156 bp, respectively. The intron/exon splice sites (mGT....AGm) and their flanking sequences are closely related to the consensus splicing sites (-mGTAAGT ...CAGm-) (32). The presumptive protein coding region of the KLK-L2 gene is formed of 879 bp nucleotide sequence encoding  
10       a deduced 293-amino acid polypeptide with a predicted molecular weight of 32 KDa. There are two potential translation initiation codons (ATG) at positions 1 and 25 of the predicted first exon (numbers refer to SEQ. ID. NO. 13 and GenBank Accession #AF135028). It is assumed that the first ATG will be the initiation codon, since: (1) the flanking sequence of that codon (GCGGCCATGG) matches closely with the Kozak consensus sequence for initiation of translation (GCC A/G CCATGG) (33) and is exactly the  
15       same as that of the homologous zyme gene. At this initiation codon, the putative signal sequence at the N-terminus is similar to other trypsin-like serine proteases (prostase and EMSP) (Figure 9). The cDNA ends with a 328 bp of 3' untranslated region containing a conserved poly adenylation signal (AATAAA) located 11 bp up-stream of the poly A tail ( at a position exactly the same as that of the zyme poly A tail)(11).

20       A hydrophobicity study of the KLK-L2 gene shows a hydrophobic region in the N-terminal region of the protein (Figure 10), suggesting that a presumed signal peptide is present. By computer analysis, a 29-amino acid signal peptide is predicted with a cleavage site at the carboxyl end of Ala<sup>29</sup>. For better characterization of the predicted structural motif of the KLK-L2 protein, it was aligned with other members of the kallikrein multi-gene family, (Figure 9), and the predicted signal peptide cleavage site was found to match with the predicted signal cleavage sites of zyme (11), KLK1(1), KLK2 (8), and KLK-L1. Also,  
25       sequence alignment supports, by analogy, the presence of a cleavage site at the carboxyl end of Ser<sup>66</sup>, which is the exact site predicted for cleavage of the activation peptide of all the other kallikreins aligned in Figure 9. Interestingly, the starting amino acid sequence of the mature protein (I I N G (S) D C ) is conserved in the prostase and enamel matrix serine proteinase 1 (EMSP) genes. Thus, like other kallikreins, KLK-L2 is likely also synthesized as a preproenzyme that contains an N-terminal signal peptide (prezymogen)  
30       followed by an activation peptide and the enzymatic domain.

35       The presence of aspartate (D) in position 239 suggests that KLK-L2 will possess a trypsin-like cleavage pattern like most of the other kallikreins (e.g., KLK1, KLK2, TLSP, neuropsin, zyme, prostase, and EMSP) but different from PSA which has a serine (S) residue in the corresponding position, and is known to have a chymotrypsin like activity (Figure 9). The dotted region in Figure 9 indicates an 11-amino acid loop characteristic of the classical kallikreins (PSA, KLK1, and KLK2) but not found in KLK-L2 or other members of the kallikrein-like gene family (11).

#### Homology with the kallikrein multi-gene family

The mature 227-amino acid sequence of the predicted protein was aligned against the GenBank

database and the known kallikreins using the "BLASTP" and "BLAST 2 sequence" programs. KLK-L2 is found to have 54% amino acid sequence identity and 68% similarity with the enamel matrix serine proteinase 1 (EMSP1) gene, 50% identity with both trypsin like serine protease (TLSP) and neuropsin genes and 47%, 46%, and 42% identity with trypsinogen, zyme, and PSA genes, respectively. The multiple alignment study shows that the typical catalytic triad of serine proteases is conserved in the KLK-L2 gene (H<sup>108</sup>, D<sup>153</sup>, and S<sup>245</sup>) and, as the case with all other kallikreins, a well conserved peptide motif is found around the amino acid residues of the catalytic triad [i.e., histidine (WLLTAAHC), serine(GDSGGP), and aspartate(DLMLI)] (10, 11).

Twelve cysteine residues are present in the putative mature KLK-L2 protein, ten of them are conserved in all the serine proteases that are aligned in Figure 9, and would be expected to form disulphide bridges. The other two cysteines (C<sup>178</sup> and C<sup>279</sup>) are not found in PSA, KLK1, KLK2 or trypsinogen, however, they are found in similar positions in prostase, EMSP1, zyme, neuropsin, and TLSP genes and are expected to form an additional disulphide bond. Twenty nine "invariant" amino acids surrounding the active site of serine proteases have been described (39). Of these, twenty-six are conserved in KLK-L2. One of the non-conserved amino acids (Ser<sup>210</sup> instead of Pro) is also found in prostase and EMSP1 genes, the second (Leu<sup>103</sup> instead of Val) is also found in TLSP gene, and the third (Val<sup>174</sup> instead of Leu) is also not conserved in prostase or EMSP1 genes. According to protein evolution studies, each of these amino acid changes represents a conserved evolutionary substitution to a protein of the same group (39).

#### Evolution of the KLK-L2 gene

To predict the phylogenetic relatedness of the KLK-L2 gene with other serine proteases, the amino acid sequences of the kallikrein genes were aligned together using the "Clustal X" multiple alignment program and a distance matrix tree was predicted using the Neighbor-joining/UPGMA method (Figure 10). Phylogenetic analysis separated the classical kallikreins (KLK1, KLK2, and PSA) and grouped the KLK-L2 with KLK-L1, EMSP1, and TLSP (40, 41).

#### Tissue expression of the KLK-L2 gene

As shown in Table 12 and Figure 11, the KLK-L2 gene is primarily expressed in the brain, mammary gland, and testis but lower levels of expression are found in many other tissues. In order to verify the RT-PCR specificity, the PCR products were cloned and sequenced.

#### Hormonal regulation of the KLK-L2 gene

A steroid hormone receptor positive breast cancer cell line (BT-474) was used as a model to verify whether the KLK-L2 gene is under steroid hormone regulation. PSA was used as a control known to be upregulated by androgens and progestins and pS2 as an estrogen upregulated control. The results indicate that KLK-L2 is up-regulated by estrogens and progestins (Figure 12).

#### Expression of KLK-L2 in Ovarian Tissues

KLK-L2 is up-regulated (overexpressed) in ovarian tumors (Figure 13).

#### Discussion

With the aid of computer programs for gene prediction and the available EST database, a new gene, named KLK-L2 (for kallikrein like gene 2) was identified. The 3' end of the gene was verified by the

presence of "poly A" stretches in the sequenced ESTs which were not found in the genomic sequence, and the start of translation was identified by the presence of a start codon in a well conserved consensus Kozak sequence.

As is the case with other kallikreins, the KLK-L2 gene is composed of 5 coding exons and 4  
5 intervening introns and, except for the second coding exon, the exon lengths are comparable to those of other members of the kallikrein gene family (Figure 6). The exon-intron splice junctions were identified by comparing the genomic sequence with the EST sequence and were further confirmed by the conservation of the consensus splice sequence (-mGT.....AGm-) (32), and the fully conserved intron phases, as shown in Figure 6. Furthermore, the position of the catalytic triad residues in relation to the different exons is also  
10 conserved (Figure 6). As is the case with most other kallikreins, except PSA and HSCCE, KLK-L2 is more functionally related to trypsin than to chymotrypsin (34). The wide range of tissue expression of KLK-L2 should not be surprising since, by using the more sensitive RT-PCR technique instead of Northern blot analysis, many kallikrein genes were found to be expressed in a wide variety of tissues including salivary gland, kidney, pancreas, brain, and tissues of the reproductive system (uterus, mammary gland, ovary, and  
15 testis) (34). KLK-L2 is highly expressed in the brain. Another kallikrein, neuropsin, was also found to be highly expressed in the brain and has been shown to have important roles in neural plasticity in mice (35). Also, the zyme gene is highly expressed in the brain and appears to have amyloidogenic potential (11). Taken together, these data point to a possible role of KLK-L2 in the central nervous system.

It was initially thought that each kallikrein enzyme has one specific physiological substrate.  
20 However, the increasing number of substrates, which purified proteins can cleave *in vitro*, has led to the suggestion that they may perform a variety of functions in different tissues or physiological circumstances. Serine proteases encode protein cleaving enzymes that are involved in digestion, tissue remodeling, blood clotting etc., and many of the kallikrein genes are synthesized as precursor proteins that must be activated by cleavage of the propeptide. The predicted trypsin-like cleavage specificity of KLK-L2 makes it a  
25 candidate activator of other kallikreins or it may be involved in a "cascade" of enzymatic reactions similar to those found in fibrinolysis and blood clotting (36).

In conclusion, a new member of the human kallikrein gene family, KLK-L2 was characterized. This gene is hormonally regulated and it is mostly expressed in the brain, mammary gland and testis. KLK-L2 may be useful as a tumor marker.

30

#### Example 4

##### Materials and methods

##### Strategy for new gene discovery

Sequencing data of approximately 300 kb, around chromosome 19q13.3-q13.4, was obtained from the web site of the Lawrence Livermore National Laboratory (LLNL) (<http://www-bio.llnl.gov/genome/genome.html>). Different computer programs were used for putative new gene prediction, as previously  
35 described.

##### RT-PCR for KLK-L3 cDNA

Total RNA isolated from 26 different human tissues was purchased from Clontech, Palo Alto, CA.

- 41 -

cDNA was prepared as described below and used for PCR amplification. A primer set (L3-F1 and L3-R1) was used to identify the presence of the gene in tissues, and the reverse primer (L3-R1) was used with another primer (L3-F2) to amplify and clone the full cDNA of the gene. These primer sequences are shown in Table 13 (SEQ. ID. Nos. 9-12, 24-26). Tissue cDNAs were amplified at various dilutions.

#### 5 Reverse transcriptase polymerase chain reaction.

2 µg of total RNA was reverse-transcribed into first strand cDNA using the Superscript™ preamplification system (Gibco BRL, Gaithersburg, MD). The final volume was 20 µl. Based on the combined information obtained from the predicted genomic structure of the new gene and the EST sequence, two gene-specific primers (L3-F1 and L3-R1) were designed (Table 13, SEQ. ID. Nos. 9-12, 24-26) and PCR was carried out in a reaction mixture containing 1 µl of cDNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200 µM dNTPs (deoxynucleoside triphosphates), 150 ng of primers and 2.5 units of AmpliTaq Gold DNA polymerase (Roche Molecular Systems, Branchburg, NJ, USA) on a Perkin-Elmer 9600 thermal cycler. The cycling conditions were 94°C for 9 minutes, followed by 43 cycles of 94°C for 30 s, 63°C for 1 minute, and a final extension at 63°C for 10 minutes. Equal amounts of PCR products were electrophoresed on 2% agarose gels and visualized by ethidium bromide staining. All primers for RT-PCR spanned at least 2 exons to avoid contamination by genomic DNA.

#### Breast cancer cell line and hormonal stimulation experiments

The breast cancer cell line BT-474 was purchased from the American Type Culture Collection (ATCC), Rockville, MD. Cells were cultured in RPMI media (Gibco BRL, Gaithersburg, MD) supplemented with glutamine (200 mmol/L), bovine insulin (10 mg/L), fetal bovine serum (10%), antibiotics and antimycotics, in plastic flasks, to near confluency. The cells were then aliquoted into 24-well tissue culture plates and cultured to 50% confluency. 24 hours before the experiments, the culture media were changed into phenol red-free media containing 10% charcoal-stripped fetal bovine serum. For stimulation experiments, various steroid hormones dissolved in 100% ethanol were added into the culture media, at a final concentration of 10<sup>-8</sup> M. Cells stimulated with 100% ethanol were included as controls. The cells were cultured for 24 hours, then harvested for total RNA extraction by the Trizol method (Gibco-BRL). cDNA was prepared and amplified as described above. Control genes (PSA, pS2, and actin) were amplified as previously described herein.

#### Cloning and sequencing of the PCR products.

To verify the identity of the PCR products, they were cloned into the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The inserts were sequenced from both directions using vector-specific primers, with an automated DNA sequencer.

#### Identification of positive PAC and BAC genomic clones from human genomic DNA libraries

The PCR product generated with primer set Z1S and Z1AS (Table 14, SEQ.ID.NOS. 27-42), was purified and then labeled with <sup>32</sup>P by the random primer method (Sambrook, supra) and used as a probe to screen a human genomic DNA BAC library, spotted in duplicate on nylon membranes, for identification of positive clones. The filters were hybridized in 15% formamide, 500 mM Na<sub>2</sub>HPO<sub>4</sub>, 7% SDS, 1% BSA (w/v) at 65°C overnight, then washed sequentially with 2X SSC, 1X SSC, 0.2X SSC, containing 0.1% SDS

at 65°C, and then exposed to X-ray film as described (Sambrook, supra). Positive clones were obtained, plated on selective LB medium, and then a single colony was transferred into LB broth for overnight cultures. A PAC clone positive for NES1 was identified by a similar methodology as described elsewhere (14). PAC and BAC libraries were constructed by de Jong and associates (42). Purification of BAC and  
5 PAC DNA was done by a rapid alkaline lysis miniprep method, which is a modification of the standard Qiagen-Tip method. Positive clones were further confirmed by Southern blot analysis as described (Sambrook, supra).

#### **5' Rapid amplification of cDNA ends (5' RACE)**

According to the EST sequences and the computer-predicted structure of the KLK-L3 gene, two  
10 gene specific primers were designed. Two rounds of RACE reactions (nested PCR) were performed with 5µl Marathon Ready™ cDNA of human testis (Clontech) as a template. The reaction mix and PCR conditions were selected according to the manufacturer's recommendations. Positive bands were gel-purified using Qiagen Gel Purification kits according to manufacturer's recommendations.

#### **Gene-specific amplification of other genes from genomic DNA**

According to the published sequence of prostatic specific antigen (PSA), human renal kallikrein  
15 (KLK1), human glandular kallikrein (KLK2), normal epithelial cell-specific 1 gene (NES1), KLK-L1, KLK-L2 and zyme genes, gene-specific primers were designed for each of these genes (Table 14) and developed polymerase chain reaction (PCR)-based amplification protocols which allowed us to generate specific PCR products with genomic DNA as a template. The PCR reactions were carried out as described above but by  
20 using an annealing/extension temperature of 65°C.

#### **Structure analysis studies.**

Multiple alignment was performed using the clustal X software package available at:  
[ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/\(clustalx1.64b.msw.exe\)](ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/(clustalx1.64b.msw.exe)) and the multiple alignment  
program available from the Baylor College of Medicine (BCM) search launcher  
25 ([kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html](http://kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html)). Phylogenetic studies were performed using the Phylip software package available from: (<http://evolution.genetics.washington.edu/phylip/getme.html>). Distance matrix analysis was performed using the "Neighbor-Joining/UPGMA" program and parsimony analysis was done using the "Promoters" program. Hydrophobicity study was performed using the BCM  
search launcher programs (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>). Signal  
30 peptide was predicted using the SignalP WWW server (<http://www.cbs.dtu.dk/services/signal>). Protein structure analysis was performed by SAPS (structural analysis of protein sequence) program (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>).

#### **Results:**

#### **Construction of a contiguous map of the human kallikrein locus on chromosome 19q13.3-q13.4**

35 Sequence information around the human chromosome 19q13.3-q13.4 locus (the proposed kallikrein locus) is available at the Lawrence Livermore National Laboratory web site. Sequences of approximately 300 kb in length were obtained. These sequences were in the form of contigs of different lengths. A restriction analysis study of the contigs was performed using various computer programs. With

the aid of the EcoRI restriction map of this area which is also available at the LLNL web site, the relative positions of these contigs was defined in relation to each other. Some contigs were overlapping, enabling construction of a contiguous segment; however, three gaps were present. <BLAST> analysis of these segments against the GenBank database (37) enabled the precise location of two classical kallikreins, namely PSA and KLK2 to be defined. Other newly discovered serine proteases were localized which are homologous with the kallikrein genes, namely protease M/zyme/neurosin (10, 11, 12), human stratum corneum chymotryptic enzyme ( HSCCE) (55), neuropsin (28), normal epithelial cell-specific 1 gene ( NES1) (13), trypsin-like serine protease ( TLSP) (GenBank accession # AF164623), KLK-L1 (SEQ.ID.NO. 1) and KLK-L2 (SEQ.ID.NO. 13). The gaps in the 300 kb genomic sequence were partially filled as follows:

- (a) The margins of the first gap were found to contain the 5' and 3' ends of the KLK2 gene; this gap was filled with the genomic structure of the KLK2 gene (GenBank Accession # M18157).
- (b) The margins of the third gap (gaps are numbered from centromere to telomere) were found to have the 3' and 5' ends of the zyme gene mRNA sequence; thus, a radiolabeled probe specific for the zyme gene was used to screen a human BAC library and two positive clones were obtained. Restriction analysis was performed, followed by Southern blotting and a fragment containing the zyme gene was obtained and sequenced, thus filling this gap.
- (c) The second gap (between KLK-L1 and KLK-L2 genes) still exists and the EcoRI restriction map of this area was used to approximately define its length (Figure 14).

Further support for the relative locations of these genes was obtained by performing PCR reactions with gene-specific primers to screen genomic DNA clones. The most centromeric group of genes (PSA, KLK2, KLK-L1, KLK-L2 and zyme ) were found to be clustered in one genomic BAC clone, and the next group (HSCCE, neuropsin, KLK-L3 and NES1) were found to be clustered together in another clone, as expected from the data of Figure 14.

#### Cloning of the KLK-L3 gene

A putative new gene, formed of three exons, was predicted by computer analysis of the genomic sequence. The predicted exons were subjected to sequence homology search against the human EST database (dbEST) and revealed an EST clone (GenBank accession # AA583908) which exhibited 99% homology with the putative gene. This EST was obtained, purified and sequenced and the sequence was aligned by BLAST software (37) against the genomic area that contains the putative gene. An additional exon, downstream of the predicted structure, was identified. The 3' end of the gene was verified by: (a) The presence of the serine residue (S) of the catalytic triad in a well-conserved region. This highly conserved motif (GDSGGP) always occurs at the beginning of the last exon in all known kallikreins. (b) The presence of a stop codon that is in frame with the predicted amino acid sequence. (c) The presence of a 19-poly A stretch at the end of the EST that was not found in the genomic sequence.

To verify the accuracy of the cDNA sequence of the gene, PCR reactions were performed using gene-specific primers for the first and last exons of the predicted structure of the gene (L3-F2 and L3-R1) with cDNA isolated from different human tissues as putative templates. A positive band of the expected

size was isolated from testis cDNA and fully sequenced. Its sequence was aligned by BLAST against the genomic sequence to unequivocally define the exon/intron boundaries. For further characterization of the 5' end of the gene, 5'RACE reaction was performed using Marathon Ready cDNA from testis as a template. This allowed identification of an additional exon that contains the start codon and 5' untranslated region. The full sequence of the gene is shown in SEQ. ID. NO. 21 (GenBank Accession # AF135026) and the amino acid sequences of KLK-L3 proteins are shown in SEQ. ID. Nos. 22 and 23.

#### Structural characterization of the KLK-L3 gene:

As shown in Figure 15, the KLK-L3 gene is formed of 5 coding exons and 4 intervening introns, although, as with other kallikreins, the presence of further upstream untranslated exon(s) could not be ruled out (14, 28). All of the exon /intron splice sites conform to the consensus sequence for eukaryotic splice sites (32). The gene further follows strictly the common structural features of the human kallikrein multigene family, as described below.

The predicted protein-coding region of the gene is formed of 753 bp, encoding a deduced amino acid polypeptide with a predicted molecular weight of 27.5 kDa. A potential translation initiation codon is found at position 28 of the predicted first exon (numbers of nucleotides refer to SEQ. ID. NO. 21 or GenBank Accession # AF135026. This codon does not match well with the consensus Kozak sequence (33), however, it has a purine at position (-3) which occurs in 97% of vertebrate mRNAs (43), and it is almost identical to the sequence of the zyme gene flanking the start codon. It should also be noted that most kallikreins do not have the consensus G nucleotide in position (+4).

Nucleotides 6803-6808 (AGTAAA) closely resemble a consensus polyadenylation signal (44) and are followed by a stretch of 19 poly A nucleotides not found in genomic DNA, after a space of 14 nucleotides. No other potential polyadenylation signals were discernable in the 3' untranslated region, suggesting that the above motif is indeed the polyadenylation signal. The same polyadenylation signal motif was predicted for the KLK1 and KLK2 genes.

Although the KLK-L3 protein sequence is unique, comparative analysis revealed that it is highly homologous to other members of the kallikrein multigene family. KLK-L3 shows 40% protein identity with the TLSP gene product and 38% and 33% identity with the KLK-L2 and KLK1 proteins, respectively. Hydrophobicity analysis revealed that the amino-terminal region is quite hydrophobic (Figure 16), consistent with the possibility that this region may harbor a signal sequence, analogous to other serine proteases. Computer analysis of the amino acid sequence of KLK-L3 predicted a cleavage site between amino acids 19 and 20 (GWA-DT). Sequence alignment (Figure 17) also revealed a potential cleavage site (Arg<sup>22</sup>), at a site homologous to other serine proteases (lysine (K) or arginine (R) is present in most cases). Several evenly distributed hydrophobic regions throughout the KLK-L3 polypeptide are consistent with a globular protein, similar to other kallikreins and serine proteases. The dotted region in Figure 17 indicates an 11-amino acid loop characteristic of the classical kallikreins (PSA, KLK1, and KLK2) but not found in KLK-L3 or other members of the kallikrein multi-gene family (11, 41).

Twenty nine "invariant" amino acids surrounding the active site of serine proteases have been described. Of these, twenty-six are conserved in KLK-L3. One of the unconserved amino acids (Ser<sup>168</sup>

instead of Pro) is also found in prostase, KLK-L2 and enamel matrix serine proteinase (EMSP1) genes. The second (Leu<sup>58</sup> instead of Val) is also found in TLSP and KLK-L2 genes, and the third is Ala<sup>26</sup> instead of Gly. According to protein evolution studies, each of these changed amino acids represents a conserved evolutionary change to a protein of the same group (45). Twelve cysteine residues are present in the putative mature KLK-L3 protein, ten of them are conserved in all the serine proteases that are aligned in Figure 17, and would be expected to form disulphide bridges. The other two (C<sup>136</sup> and C<sup>238</sup>) are not found in PSA, KLK1, KLK2 or trypsinogen; however, they are found in similar positions in prostase, HSCCE, zyme neuropsin, and TLSP genes and are expected to form an additional disulphide bond.

To predict the phylogenetic relatedness of the KLK-L3 gene with other serine proteases, the amino acid sequences of the kallikrein genes were aligned together using the "Clustal X" multiple alignment program and a distance matrix tree was predicted using the Neighbor-joining/UPGMA method (Figure 18). Phylogenetic analysis separated the classical kallikreins (KLK1, KLK2, and PSA) and grouped KLK-L3 with TLSP, neuropsin, zyme, HSCCE and prostase/KLK-L1, consistent with previously published studies (11, 41).

#### **Tissue expression and hormonal regulation of the KLK-L3 gene**

As shown in Figure 19, the KLK-L3 gene is primarily expressed in thymus, testis, spinal cord, cerebellum, trachea, mammary gland, prostate, brain, salivary gland, ovary and skin (the latter two tissues are not shown in the figure). Lower levels of expression are seen in fetal brain, stomach, lung, thyroid, placenta, liver, small intestine, and bone marrow. No expression was seen in uterus, heart, fetal liver, adrenal gland, colon, spleen, skeletal muscle, pancreas, and kidney. In order to verify the RT-PCR specificity, representative PCR products were cloned and sequenced. Figure 20 shows that KLK-L3 gene is regulated by steroid hormones in the human breast cancer cell line BT-474.

#### **DISCUSSION**

A human kallikrein gene locus has been defined, and the first detailed map describing the relative positions of the kallikreins and other kallikrein-like genes has been constructed (Figure 14). This map is consistent with previous reports on the localization of the classical kallikreins and the approximate mapping of some new kallikreins by radiation hybrid and FISH techniques (9, 14, 67). It should be noted, however, that the lengths of certain segments of this map (as depicted in Figure 14) are dependent on the EcoRI restriction map of the area and are measured in terms of approximate kb units. Also, the measure of intervals between genes may change slightly in the future, since some kallikreins may have extra 5'-exon(s) that have not as yet been identified. Kallikreins with verified 5'-untranslated exons include NES1 (14), zyme, and neuropsin (35). This map is also directional; it indicates that PSA and KLK2 genes are transcribed in the same direction (centromere to telomere) and that the rest of the kallikrein-like genes are transcribed in the reverse direction (Figure 14).

An early report indicated that KLK1 is located approximately 31 kb centromeric to PSA (9). The map described extends only 24 kb centromeric to PSA, and for this reason, KLK1 was not precisely localized. Thus, the exact location of the KLK1 gene is still to be defined from linear chromosome 19 sequencing data. The possibility still exists that this locus is extended further, and that other kallikrein-like

genes may be located upstream of PSA or downstream from TLSP.

Traditionally, kallikreins are characterized by their ability to liberate lysyl-bradykinin (kallidin) from kininogen (2). In humans, only KLK1 meets this "functional" definition. KLK2 and KLK3 are assigned to the same family based on the strong structural similarities of the genes and proteins and the close localization of these genes on the same chromosomal region (20). More recently, a new structural concept has emerged to describe kallikreins. Richards and co-workers introduced the concept of a "kallikrein multigene family" in mice, to refer to these genes (20, 21). This definition is not based much on the specific enzymatic function of the gene product, but more on its sequence homology and its close linkage on mouse chromosome 7.

Irwin et al. (27) proposed that the serine protease genes can be classified into five different groups according to intron position as discussed above. The results indicate the presence of some more common structural features that are found in all kallikreins (including the newly identified KLK-L3 gene): (1) All genes are formed of 5 coding exons and 4 intervening introns (with the possibility that some genes may have extra 5' untranslated exon(s) (24, 31, 35) (Figure 21). (2) The exon lengths are usually comparable. (3) The intron phases are always conserved (I-II-I-0) (see Figure 21 for description of intron phases). (4) These genes are clustered in the same chromosomal region, apparently without any intervening non kallikrein-like genes (Figure 14). Thus, all the recently identified serine proteases that are present in this region (zyme, HSCCE, neuropsin, NES1, prostase/KLK-L1, KLK-L2 and TLSP), together with the newly identified kallikrein-like gene (KLK-L3), could be considered members of the expanded human kallikrein multigene family.

The chromosomal band 19q13 is nonrandomly rearranged in a variety of human solid tumors including ovarian cancers (46), and the currently available data indicate that the kallikrein gene locus is related to many malignancies. At least three kallikrein genes (PSA, zyme and NES1) are down regulated in breast cancer (10, 13, 47, 48), and NES1 appears to be a novel tumor suppressor gene (29). Furthermore, PSA exhibits potent antiangiogenic activity (49). It is possible that some of these kallikreins are involved in a cascade pathway, similar to the coagulation or apoptotic process, whereby pro-forms of proteolytic enzymes are activated and then act upon downstream substrates. Such activity was found for the KLK2 gene product which acts upon and activates pro PSA (50, 51).

The expanded human kallikrein gene family has similar number of members as the rodent family of genes. Some new compelling data have raised the possibility that at least some of these genes behave as tumor suppressors (29), as negative regulators of cell growth (52), as antiangiogenic (49) and apoptotic (53) molecules. The paramount diagnostic value of some members is also well-established (24, 54). For these reasons, it is important to check all members of this family of genes as potential diagnostic or prognostic markers or as candidate therapeutic targets.

The newly identified KLK-L3 gene is expressed in many tissues, including skin, thymus, central nervous system, breast, prostate, and testis. The wide range of tissue expression of KLK-L3 should not be surprising since, by using the more sensitive RT-PCR technique, many kallikrein genes were found to be expressed in a wide variety of tissues. For example, PSA, KLK2, prostase/KLK-L1, and KLK-L2 are now

known to be expressed in breast and many other tissues (30, 54).

Like many other kallikreins, KLK-L3 is regulated by steroid hormones but in a more complex fashion than PSA and KLK2 which are up-regulated by androgens and progestins (71). In the breast carcinoma cell line studied, KLK-L3 appears to be up-regulated by progestins > estrogens > androgens (Figure 20).

#### Example 5

##### Materials and Methods

##### DNA sequence on chromosome 19 and prediction of new genes

Sequencing data of approximately 300Kb of nucleotides, around chromosome 19q13.3-q13.4, was obtained from the web site of the Lawrence Livermore National Laboratory (LLNL) (<http://www.llnl.gov/genome/genome.html>) and an almost contiguous stretch of genomic sequences was constructed. A number of computer programs were used to predict the presence of putative new genes in this genomic area.

##### Expressed sequence tag (EST) searching

The predicted exons of the putative new gene were subjected to homology search using the BLASTN algorithm (37) on the National Center for Biotechnology Information web server (<http://www.ncbi.nlm.nih.gov/BLAST/>) against the human EST database (dbEST). Clones with > 95% homology were obtained from the I.M.A.G.E. consortium (38) through Research Genetics Inc, Huntsville, AL. The clones were propagated, purified and sequenced from both directions with an automated sequencer, using insert-flanking vector primers.

##### Rapid amplification of cDNA ends (3' RACE)

According to the EST sequence data and the predicted structure of the gene, two gene-specific primers were designed and two rounds of RACE reactions (nested PCR) were performed with 5µl Marathon Ready™ cDNA of human testis (Clontech, Palo Alto, CA, USA) as a template. The reaction mix and PCR conditions used were according to the manufacturer's recommendations.

##### Tissue expression

Total RNA isolated from 26 different human tissues was purchased from Clontech. cDNA was prepared as described below, and used for PCR reactions with different sets of primers (Table 15, SEQ.ID.NO.s. 46-55, 9-12). Tissue cDNAs were amplified at various dilutions.

##### Breast cancer cell line and hormonal stimulation experiments

The breast cancer cell line BT-474 was purchased from the American Type Culture Collection (ATCC), Rockville, MD. Cells were cultured in RPMI media (Gibco BRL, Gaithersburg, MD) supplemented with glutamine (200 mmol/L), bovine insulin (10 mg/L), fetal bovine serum (10%), antibiotics and antimycotics, in plastic flasks, to near confluency. The cells were then aliquoted into 24-well tissue culture plates and cultured to 50% confluency. 24 hours before the experiments, the culture media were changed into phenol red-free media containing 10% charcoal-stripped fetal bovine serum. For stimulation experiments, various steroid hormones dissolved in 100% ethanol were added into the culture media, at a final concentration of 10<sup>-8</sup> M. Cells stimulated with 100% ethanol were included as controls.

The cells were cultured for 24 hours, then harvested for mRNA extraction.

#### Reverse transcriptase polymerase chain reaction

Total RNA was extracted from the breast cancer tissues and cell lines using Trizol™ reagent (Gibco BRL) following the manufacturer's instructions. RNA concentration was determined spectrophotometrically. 2 µg of total RNA was reverse-transcribed into first strand cDNA using the Superscript™ preamplification system (Gibco BRL). The final volume was 20 µl. Based on the combined information obtained from the predicted genomic structure of the new gene and the EST sequences, two gene-specific primers were designed (L4-F1 and L4-R1, see Table 15, SEQ.ID.NO.s. 46 and 47) and PCR was carried out in a reaction mixture containing 1 µl of cDNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200 µM dNTPs (deoxynucleoside triphosphates), 150 ng of primers and 2.5 units of AmpliTaq Gold DNA polymerase (Roche Molecular Systems, Branchburg, NJ, USA) on a Perkin-Elmer 9600 thermal cycler. The cycling conditions were 94°C for 9 minutes to activate the Taq Gold DNA polymerase, followed by 43 cycles of 94°C for 30 s, 63°C for 1 minute and a final extension at 63°C for 10 min. Equal amounts of PCR products were electrophoresed on 2% agarose gels and visualized by ethidium bromide staining. All primers for RT-PCR spanned at least 2 exons to avoid contamination by genomic DNA.

To verify the identity of the PCR products, they were cloned into the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The inserts were sequenced from both directions using vector-specific primers, with an automated DNA sequencer.

#### Normal and malignant breast tissues

Normal breast tissues were obtained from women undergoing reduction mammoplasties. Breast tumor tissues were obtained from female patients at participating hospitals of the Ontario Provincial Steroid Hormone Receptor Program. The normal and tumor tissues were immediately frozen in liquid nitrogen after surgical resection and stored in this manner until extracted. The tissues were pulverized with a hammer at dry ice temperature and RNA was extracted as described above, using Trizol reagent.

#### Structure analysis

Multiple alignment was performed using the Clustal X software package available at: [ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/\[clustalx1.64b.msw.exe\]](ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/[clustalx1.64b.msw.exe]) and the multiple alignment program available from the Baylor College of Medicine (BCM), Houston, TX, USA [kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html]. Phylogenetic studies were performed using the Phylip software package available at: <http://evolution.genetics.washington.edu/phylip/getme.html>. Distance matrix analysis was performed using the "Neighbor-Joining/UPGMA" program and parsimony analysis was done using the "Protpars" program. Hydrophobicity study was performed using the BCM search launcher programs [<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>]. Signal peptide was predicted using the "SignalP" server [<http://www.cbs.dtu.dk/services/signal>]. Protein structure analysis was performed by the "SAPS" (structural analysis of protein sequence) program [<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>].

#### Results

### Cloning of the KLK-L4 gene

Computer analysis of the genomic sequence around chromosome 19 q13.3-q13.4 predicted a putative new gene formed of at least 3 exons. To experimentally verify the existence of this gene, the putative exons were subjected to sequence homology search against the human expressed sequence tag (EST) database (dbEST), and four EST clones with > 97% homology were identified (Table 16). All ESTs were cloned from testicular tissue. These clones were obtained and inserts were sequenced from both directions. Sequences were then compared with the computer-predicted structure and final selection of the intron/exon splice sites was made according to the EST sequences.

As shown in Figure 22, three ESTs match almost perfectly with the predicted 3 exons (exons 3, 4, 5) of the gene and one EST matches perfectly with predicted exons 3 and 5. However, each of the ESTs extends further upstream with different exonic patterns, suggesting the presence of different splice variants.

Attempts to translate these clone sequences demonstrated the presence, in some ESTs, of interrupting stop codons in all three possible reading frames. A homology search of the three common exons against the GenBank database revealed a cDNA sequence from the German Human Genome Project. This clone has an identical exon 2 as the long form of KLK-L4 gene [this form will be described below] but has an extended exon 3 that ends with a stop codon (Figure 22). This clone was isolated from uterine tissue and is translated by software into a truncated protein product of 196 amino acids which is followed by a 3' untranslated region [GenBank accession # AL050220].

Screening of cDNAs from 26 different tissues by RT-PCR, using gene-specific primers for exons 3 and 5 [L4-F1 and L4-R1] (Table 15 & Figure 22) revealed that this gene is expressed in many tissues. Four tissues that show the highest level of expression [salivary gland, mammary gland, prostate, and testis] (Figure 23) and uterus [the EST clone AL050220 was isolated from this tissue] were selected for identification of the full structure of the gene. Different PCR reactions were performed using one reverse primer (L4-R1) together with each of the forward primers located in upstream exons that were found in the different EST clones [primers L4-B, L4-D, L4-E] (Table 15 & Figure 22). The PCR reactions were performed under different experimental conditions, using the EST clones as positive controls, and the PCR products were sequenced. None of these forms were found in any of the tissues, except in testis where all three forms were found.

By RT-PCR of the KLK-L4 gene using primers L4-R1 and L4-F1, it was found that the gene is expressed in a wide variety of tissues (Figure 23). In order to obtain the structural forms that exist in these tissues, a homology study was performed. Aligning the predicted polypeptide of the KLK-L4 gene with all other kallikreins and kallikrein-like genes, suggested, by homology, that at least two more exons should be present upstream of the predicted three exons. The genomic fragment upstream of the third exon was subjected to further computer analysis for gene prediction, and exon 2 was identified based on: a) a consensus exon/intron splice site b) preservation intron phase II after this exon, in agreement to intron phases of all other known kallikreins c) presence of the histidine residue of the catalytic triad (H<sup>76</sup>) surrounded by a well-conserved peptide motif [see below] just before the end of this exon d) comparable exon length to other kallikrein genes. A potential first exon was also predicted from the upstream genomic

sequence, based on the preserved intron phase (phase I), and the existence of an in-frame start codon that is located at a comparable distance [in relation to other kallikreins] from the end of this exon. To verify this predicted structure, a PCR reaction was performed using one reverse primer (L4-R1) together with another forward primer that is located in the predicted first exon (primer L4-X1) (Table 15 & Figure 22).

5 Two main PCR bands were obtained from the tissues examined; the expected 819 bp band (predominant) and an additional minor band of about 650 bp (Figure 24). Cloning and sequencing of these two bands revealed that the gene exists in two main forms in these tissues; the long form [SEQ. ID. No. 43 or GenBank Accession No. AF135024] and another form [referred to as the short KLK-L4 variant] that utilizes an upstream alternative splice donor site, located inside exon 3, thus creating an mRNA product

10 that that is 214 bp shorter. This alternative splice site causes frame-shifting of the coding region that will generate a predicted stop codon at the beginning of exon 4, giving rise to a truncated protein product that does not contain the serine residue of the catalytic triad (Figures 24 and 25).

Aligning the long KLK-L4 form with the ESTs (Figure 22) demonstrated that all ESTs utilize a different splice donor site located 80 bp downstream from the end of exon 3. These additional 80 bp

15 contain an in-frame stop codon at nucleotide position 5505 which will lead to the formation of a shorter polypeptide product. They also utilize an alternative polyadenylation signal located at position 8706 [numbers refer to SEQ. ID. No. 43 or GenBank Accession No. AF135024]. The clone from the German Genome Project utilizes another splice donor site that is located further downstream, inside intron 3, and ends up with a poly A tail without having a fourth or fifth exon. The same stop codon (position 5505) will

20 be in-frame, and therefore, a truncated protein product is predicted to be formed (Figure 22).

In order to obtain the 3' end of the gene, a 3'RACE reaction was performed, and an additional 375 bp fragment of 3' untranslated region, downstream from PCR primer L4-R1, was obtained. This fragment was further confirmed to be present in all tissues tested, by performing a PCR reaction using primers L4-F1 and L4-R3 (Table 15 & Figure 22). This fragment ends with a putative polyadenylation signal variant

25 (TATAAA).

#### **Structural characterization of the KLK-L4 gene and its protein product**

The long form of the KLK-L4 gene is presented in Figure 25 (SEQ.ID.NO. 43). KLK-L4 is formed of five coding exons and four intervening introns, spanning an area of 8,905 bp of genomic sequence on chromosome 19q13.3-q13.4. The lengths of the coding regions are 52, 187, 269, 137 and 189

30 bp, respectively. The predicted protein coding region of the gene is formed of 831 bp, encoding a deduced 277-amino acid protein with a predicted molecular mass of 30.6 kDa (Figure 25). The intron/exon splice sites (mGT....AGm, where m is any base) and their flanking sequences are in agreement with the consensus splice site sequence. A potential translation initiation codon is present at position 45 of the predicted first exon [numbers refer to SEQ. ID. No. 43]. The cDNA extends at least 382 bp further downstream from the

35 stop codon and a putative polyadenylation signal (TATAAA) is present at the end of this region (Figure 25).

Hydrophobicity analysis revealed that the amino-terminal region is quite hydrophobic (Figure 26), consistent with the possibility that this region may harbor a signal sequence, analogous to other serine

proteases. Figure 26 also shows the presence of several evenly distributed hydrophobic regions throughout the KLK-L4 polypeptide, which are consistent with a globular protein, similar to other serine proteases (13). Computer analysis of the amino acid sequence of KLK-L4 predicted a cleavage site between amino acids 20 and 21 (GVS-QE). Sequence homology with other serine proteases (Figure 27) predicted another potential cleavage site (Lys25) in close proximity. Most other kallikreins are activated by cleavage after arginine or lysine. Thus, the protein product is very likely to be a secreted protein. The dotted region in Figure 27 indicates an 11-amino acid loop characteristic of the classical kallikreins (PSA, KLK1, and KLK2) which is not found in KLK-L4 or other members of the kallikrein multi-gene family (11,13, 35).

Amino acid sequences for KLK-L4 proteins are shown in SEQ.ID.NO. 44 and 45.

Sequence analysis of eukaryotic serine proteases indicates the presence of twenty nine invariant amino acids (39). Twenty eight of them are conserved in the KLK-L4 protein and the remaining amino acid (Q182 instead of P) is not conserved among all other kallikreins (Figure 27). Ten cysteine residues are present in the putative mature KLK-L4 protein. These are conserved in all the serine proteases that are aligned in Figure 27, and would be expected to form disulphide bridges. The presence of aspartate (D) in position 239 suggests that KLK-L4 will possess a trypsin-like cleavage pattern, similarly to most of the other kallikreins [e.g., KLK1, KLK2, TLSP, neuropsin, zyme, prostase, and EMSP] but different from PSA which has a serine (S) residue in the corresponding position, and is known to have chymotrypsin like activity (Figure 27) (2,40).

#### Mapping and chromosomal localization of the KLK-L4 gene

Alignment of the KLK-L4 gene and the sequences of other known kallikrein genes within the 300 Kb area of interest [the human kallikrein gene family locus], enabled precise localization of all known genes and to determine the direction of transcription, as shown by the arrows in Figure 28. The PSA gene lies between KLK1 and KLK2 genes and is separated by 13, 319 base pairs (bp) from KLK2, and both genes are transcribed in the same direction (centromere to telomere). All other kallikrein-like genes are transcribed in the opposite direction. KLK-L4 is 13 kb centromeric from KLK-L6 [SEQ.ID.NO. 65], and 21 kb more telomeric to KLK-L5 [SEQ. ID. NO. 56].

#### Homology with the kallikrein multi-gene family

Alignment of the amino acid sequence of the KLK-L4 protein (long form) against the GenBank database and the known kallikreins, using the BLAST algorithm (37), indicated that KLK-L4 has 51% amino acid sequence identity with the TLSP and zyme genes, 49% identity with KLK-L2 and 47% and 45% identity with PSA and KLK2 genes, respectively. Multiple alignment study shows that the typical catalytic triad of serine proteases is conserved in the KLK-L4 gene (H<sup>108</sup>, D<sup>153</sup>, and S<sup>245</sup>) and, as is the case with all other kallikreins, a well conserved peptide motif is found around the amino acid residues of the catalytic triad [i.e. histidine (WLLTAAHC), serine (GDSGGP), and aspartate (DLMLI)] (Figure 27) (1, 11, 13, 35).

In addition, several other residues were found to be fully or partially conserved among the human kallikrein gene family, as further shown in Figure 27. To predict the phylogenetic relatedness of the KLK-L4 gene with other serine proteases, the amino acid sequences of the kallikrein genes were aligned together using the "Clustal X" multiple alignment program and a distance matrix tree was predicted using the Neighbor-

joining/UPGMA method (Figure 29). Phylogenetic analysis separated the classical kallikreins (KLK1, KLK2, and PSA) and grouped KLK-L4 with zyme, TLSP, KLK-L3, neuropsin, and NES1 genes, consistent with previously published studies (41) and indicating that this group of genes probably arose from a common ancestral gene by duplication.

#### 5 Tissue expression and hormonal regulation of the KLK-L4 gene

As shown in Figure 23, the KLK-L4 gene is primarily expressed in mammary gland, prostate, salivary gland and testis, but, as is the case with other kallikreins, lower levels of expression are found in many other tissues. In order to verify the RT-PCR specificity, the PCR products were cloned and sequenced.

10 A steroid hormone receptor-positive breast cancer cell line (BT-474) was used as a model, to verify whether the KLK-L4 gene is under steroid hormone regulation. PSA was used as a control gene, known to be up-regulated by androgens and progestins and pS2 as an estrogen up-regulated control gene in the same cell line. Preliminary results indicate that KLK-L4 is up-regulated by progestins and androgens and to a lower extent by estrogens (Figure 30).

#### 15 Expression of KLK-L4 in breast cancer tissues and cell lines

To characterize the extent and frequency of expression of the KLK-L4 gene in breast tumors, cDNA derived from 3 normal and 19 malignant breast tissues and 3 breast cancer cell lines was used. The data were interpreted by comparison of band intensities. Out of the 19 tumors, KLK-L4 gene expression was undetectable in 7, lower than normal tissues in 9, comparable to the normal tissues in 1, and higher than  
20 normal tissues in 2 tumors. Without hormonal stimulation, the BT-474 and T-47D cell lines had no detectable KLK-L4 mRNA, while the MCF-7 cell line was positive. These preliminary results suggest that this gene is down-regulated in the majority (16/19) of breast tumors.

#### Discussion

The established kallikreins (KLK1, KLK2, and PSA), trypsinogen and chymotrypsinogen belong  
25 to a group that has: (1) an intron just downstream from the codon for the active site histidine residue, (2) a second intron downstream from the exon containing the codon for the active site aspartic acid residue, and (3) a third intron just upstream from the exon containing the codon for the active site serine residue. Figure 31 shows that KLK-L4 meets the above mentioned criteria; moreover, is located in close proximity to other kallikrein genes on the chromosomal locus 19q13.3-q13.4 (Figure 28).

30 The preliminary findings, supporting that the KLK-L4 gene may be down-regulated in a subset of breast cancers, is not surprising. There is now growing evidence that many of the kallikreins and kallikrein-like genes that are clustered in the same chromosomal region (Figure 28) are related to malignancy. PSA is the best marker for prostate cancer so far (24). A recent report provided evidence that PSA has antiangiogenic activity, and that this activity may be related to its function as a serine protease  
35 (49). This study suggested that other serine proteases, including new members of the kallikrein multigene family of enzymes, should also be evaluated for potential antiangiogenic activity (49). Recent reports suggest that hK2 (encoded by the KLK2 gene) could be another useful diagnostic marker for prostate cancer (57, 58). NES1 appears to be a tumor suppressor gene (29). The protease M gene was shown to

be differentially expressed in primary breast and ovarian tumors (10), and the human stratum corneum chymotryptic enzyme has been shown to be expressed at abnormally high levels in ovarian cancer (59).

Another recently identified kallikrein-like gene, located close to KLK-L4 and tentatively named tumor-associated, differentially expressed gene-14 (TADG14) [an alternatively spliced form of neuropsin, see  
5 Figure 28] was found to be overexpressed in about 60% of ovarian cancer tissues (59). Also, prostate/KLK-L1, another newly discovered kallikrein-like gene, is speculated to be linked to prostate cancer (41). Thus, extensive new literature suggests multiple connections of many kallikrein genes to various forms of human cancer.

The removal of intervening RNA sequences (introns) from the pre-messenger RNA in eukaryotic  
10 nuclei is a major step in the regulation of gene expression (60). RNA splicing provides a mechanism whereby protein isoform diversity can be generated and the expression of particular proteins with specialized functions can be restricted to certain cell or tissue types during development (60). The sequence elements in the pre-mRNA at the 5' and 3' splice sites in metazoans have very loose consensus sequence; only the first and the last two bases (GT..AG) of the introns are highly conserved (Sambrook,  
15 supra). These sequences cannot be the sole determinants of splice site selection, since identical, but not ordinarily active, consensus sequences can be found within both exons and introns of many eukaryotic genes. Other protein factors and sequences downstream of the splice sites are also involved.

The existence of multiple splice forms is frequent among kallikreins. Distinct RNA species are transcribed from the PSA gene, in addition to the major 1.6-kb transcript (61). Several distinct PSA  
20 transcripts have been described by Reigman et al (7). Interestingly, one of these clones lacks the 3' untranslated region and the first 373 nucleotides of the open reading frame, and has an extended exon that contains a stop codon, a pattern that is comparable with some alternative forms of the KLK-L4 cDNA, as described here (Figure 22). Heuze et al., reported the cloning of a full-length cDNA corresponding to a 2.1 kb PSA mRNA. This form results from the alternative splicing of intron 4 and lacks the serine residue  
25 that is essential for catalytic activity (61). Also, Reigman et al reported the identification of two alternatively spliced forms of the human glandular kallikrein 2 (KLK2) gene (62). A novel transcript of the tissue kallikrein gene (KLK1) was also isolated from the colon (63). Interestingly, this transcript lacks the first two exons of the tissue kallikrein gene, but the last three exons were fully conserved, a pattern that is similar to the findings with some ESTs containing parts of the KLK-L4 gene (Figure 22). Neuropsin,  
30 a recently identified kallikrein-like gene, was found to have two alternatively spliced forms, in addition to the major form (59, 64). Here, the cloning of the KLK-L4 gene is described and the identification of a number of alternative mRNA forms. These forms may result from alternative splicing (Sambrook, supra), retained intronic segment (7), or from the utilization of an alternative transcription initiation site (63). Because the long form of KLK-L4 and the major alternative splice variant [short KLK-L4 variant] (Figure  
35 24) have an identical 5' sequence required for translation, secretion and activation, it is possible to assume that both mRNAs encode for a secreted protein (61).

In order to investigate the relative predominance of the long KLK-L4 and related forms, cDNA from various tissues was amplified by PCR. Although, in general, it is difficult to use PCR for quantitative

comparisons between mRNA species, in this experiment, [mRNAs of comparable sizes, using one set of primers under identical conditions], such a comparison is reasonable (62). In all five normal tissues examined [breast, prostate, testis, salivary gland and uterus] the long form of KLK-L4 was the predominant, with minimal level of expression of the short form (Figure 24).

5       The presence of alternatively spliced forms may be related to malignancy. Recent literature suggests that distinct molecular forms of PSA could be expressed differently by malignant versus benign prostate epithelium (65). Aberrant PSA mRNA splicing in benign prostatic hyperplasia, as opposed to prostate cancer, has been described by Henttu et al (66). In addition, it has been recently postulated that different prostatic tissues potentially harboring occult cancer could account for the presence of various  
10       forms of PSA (65).

### Example 6

#### Materials and Methods

##### DNA sequence on chromosome 19

Sequencing data of approximately 300Kb of nucleotides on chromosome 19q13.3-q13.4 was  
15       obtained from the web site of the Lawrence Livermore National Laboratory (LLNL) (<http://www-bio.llnl.gov/genome/genome.html>). This sequence was in the form of 9 contigs of different lengths. Restriction enzyme analysis, long PCR strategies, followed by DNA sequencing, BAC and PAC library screening and end sequencing of selected clones, were used to construct a contiguous genomic region, representing the complete human kallikrein gene locus.

##### 20       New gene identification

A number of computer programs were used to predict the presence of putative new genes within the contiguous genomic area of interest. The ability of these programs for predicting new genes was first examined by using the genomic sequences of the known kallikreins as testing parameters. The most reliable computer programs; GeneBuilder (gene prediction) (<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html>),  
25       GeneBuilder (exon prediction) (<http://l25.itba.mi.cnr.it/~webgene/genebuilder.html>), Grail 2 (<http://compbio.ornl.gov>), and GENEID-3 (<http://apollo.imim.es/geneid.html>) were selected for further use.

##### Expressed sequence tag (EST) searching

The predicted exons of the putative new gene were subjected to homology search using the BLASTN algorithm (37) on the National Center for Biotechnology Information web server (<http://www-ncbi.nlm.nih.gov/BLAST/>) against the human EST database (dbEST). A clone with > 95% homology was  
30       obtained from the I.M.A.G.E. consortium (38) through Research Genetics Inc, Huntsville, AL. This clone was propagated, purified and sequenced from both directions with an automated sequencer, using insert-flanking vector primers.

##### Rapid amplification of cDNA ends (RACE)

35       According to the EST sequence and the predicted structure of the gene, two sets of gene-specific primers were designed for 5' and 3' RACE reactions. Two rounds of RACE reactions (nested PCR) were performed for each type of RACE with 5µl Marathon Ready™ cDNA of human testis and prostate (Clontech, Palo Alto, CA, USA) as templates. The reaction mix and PCR conditions were selected

- 55 -

according to the manufacturer's recommendations. In brief, the initial denaturation was for 5 min at 94°C, followed by 94°C for 5 s and 72°C for 2 min, for 5 cycles; then, 94°C for 5 s and 70°C for 2 min, for 5 cycles; then, 94°C for 5 s and 65°C for 2 min for 30 cycles for the first reaction and 25 cycles for the nested PCR reaction.

## 5 Tissue expression

Total RNA isolated from 26 different human tissues was purchased from Clontech, Palo Alto, CA. cDNA was prepared as described below for the tissue culture experiments and used for PCR reactions. After aligning all known kallikrein genes, two primers (KLK-L5-R1 and KLK-L5-F1) (Table 17, SEQ.ID.NO. 61-64, 9-12, and Figure 32) were designed from areas with relatively low homology. Tissue cDNAs were amplified at various dilutions. Due to the high degree of homology between kallikreins, and to exclude non-specific amplification, PCR products were cloned and sequenced.

### Normal and malignant breast tissues

Normal breast tissues were obtained from women undergoing reduction mammoplasties. Breast tumor tissues were obtained from female patients at participating hospitals of the Ontario Provincial Steroid Hormone Receptor Program. The normal and tumor tissues were immediately frozen in liquid nitrogen after surgical resection and stored in this manner until extracted. The tissues were pulverized with a hammer under liquid nitrogen and RNA was extracted as described below, using Trizol reagent.

### Breast and prostate cancer cell lines and hormonal stimulation experiments

The breast cancer cell lines BT-474 and T-47D, and the LNCaP prostate cancer cell line were purchased from the American Type Culture Collection (ATCC), Rockville, MD. Cells were cultured in RPMI media (Gibco BRL, Gaithersburg, MD) supplemented with glutamine (200 mmol/L), bovine insulin (10 mg/L), fetal bovine serum (10%), antibiotics and antimycotics, in plastic flasks, to near confluency. The cells were then aliquoted into 24-well tissue culture plates and cultured to 50% confluency. 24 hours before the experiments, the culture media were changed into phenol red-free media containing 10% charcoal-stripped fetal bovine serum. For stimulation experiments, various steroid hormones dissolved in 100% ethanol were added into the culture media at a final concentration of  $10^{-8}$  M. Cells stimulated with 100% ethanol were included as controls. The cells were cultured for 24 hours, then harvested for mRNA extraction.

### Reverse transcriptase polymerase chain reaction (RT-PCR)

Total RNA was extracted from the cell lines or tissues using Trizol reagent (Gibco BRL) following the manufacturer's instructions. RNA concentration was determined spectrophotometrically. 2 µg of total RNA was reverse-transcribed into first strand cDNA using the Superscript™ preamplification system (Gibco BRL). The final volume was 20 µl. Based on the combined information obtained from the predicted genomic structure of the new gene and the EST sequences, two gene-specific primers were designed (KLK-L5-F1 and KLK-L5-R1) (Table 17) and PCR was carried out in a reaction mixture containing 1 µl of cDNA, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200 µM dNTPs (deoxynucleoside triphosphates), 150 ng of primers and 2.5 units of AmpliTaq Gold DNA polymerase (Roche Molecular Systems, Branchburg, NJ, USA) on a Perkin-Elmer 9600 thermal cycler. The cycling

conditions were 94°C for 9 minutes to activate the Taq Gold DNA polymerase, followed by 43 cycles of 94°C for 30 s, 63°C for 1 minute and a final extension step at 63°C for 10 min. Equal amounts of PCR products were electrophoresed on 2% agarose gels and visualized by ethidium bromide staining. All primers for RT-PCR spanned at least 2 exons to avoid contamination by genomic DNA.

- 5 To verify the identity of the PCR products, they were cloned into the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The inserts were sequenced from both directions using vector-specific primers, with an automated DNA sequencer.

#### Structure analysis

- 10 Multiple alignment was performed using the Clustal X software package available at: [ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/ \(clustalx1.64b.msw.exe\)](ftp://ftp.ebi.ac.uk/pub/software/dos/clustalw/clustalx/ (clustalx1.64b.msw.exe)) and the multiple alignment program available from the Baylor College of Medicine (BCM), Houston, TX, USA ([kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html](http://kiwi.imgen.bcm.tmc.edu:8808/search-launcher/launcher/html)). Phylogenetic studies were performed using the Phylip software package available at: <http://evolution.genetics.washington.edu/phylip/getme.html>. Distance matrix analysis was performed using the "Neighbor-Joining/UPGMA" program and parsimony analysis was done using
- 15 the "Protpars" program. Hydrophobicity study was performed using the BCM search launcher programs (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>). Signal peptide was predicted using the "SignalP" server (<http://www.cbs.dtu.dk/services/signal>). Protein structure analysis was performed by "SAPS" (structural analysis of protein sequence) program (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/struc-predict.html>).

## 20 RESULTS

### Identification of the KLK-L5 gene

- Computer analysis of the genomic area of interest (300Kb around chromosome 19q13.3-q13.4) predicted a putative gene comprised of at least three exons. Screening of the human expressed sequence tag (EST) database revealed an EST clone (GenBank Accession #394679) with 99% homology with the
- 25 predicted exons. This clone was obtained, purified, and sequenced. The full-length sequence of the EST was compared with the genomic area containing the putative new gene and showed 100% homology with certain areas (exons), which were separated by introns. This alignment indicated that the new gene was comprised of 7 exons. Sequence homology comparisons and phylogenetic analysis revealed that this new gene is structurally similar to known kallikreins and other serine proteases (see below). Since four other
- 30 new kallikrein-like genes were discovered in this area, this gene was named KLK-L5 (for kallikrein-like gene 5). Attempts to translate the coding region in all three possible reading frames indicated that only one reading frame will produce a full-length polypeptide chain without interrupting in-frame stop codons. Further support for the correctness of this reading frame was obtained by demonstrating that only this frame will preserve the three amino acid residues necessary for serine protease activity (catalytic triad) and the
- 35 conserved motifs around them. An in-frame methionine start codon was found in the second exon. This start codon falls within a typical consensus Kozak sequence (CCACCATGG) (33). Thus, the gene will have at least one 5' untranslated exon, similarly to other kallikrein-like genes [e.g. zyme, the normal epithelial cell-specific 1 gene (NES1) (14), and neuropsin (35)]. 5' and 3' RACE reactions were performed in order to

obtain the 5' and 3' ends of the gene. No more sequence was obtained by 5' RACE. However, 3' RACE enabled identification of the 3' untranslated region of the gene. The additional sequence ends with a poly-A stretch that does not exist in the genomic structure, thus marking the 3' end of the gene and the start of the poly-A tail.

## 5 Splice variants of the KLK-L5 gene

Homology analysis of the KLK-L5 gene with other kallikreins revealed the presence of an additional 3' exon, an observation that has not been reported for any other member of the kallikrein multigene family. Furthermore, two different PCR bands were obtained with the 3' RACE. Sequencing of these bands revealed that this gene has at least two splice forms at its 3' end; one form in which the last exon is a single continuous fragment, and another form in which the last exon is split into two exons, with an intervening intron. In order to identify the full structure of other possible splice variants of the gene, PCR was performed using two primers (L5-F2 and L5-R2) (Table 17 and Figure 32). cDNA from 26 different tissues were used as templates and the reaction was performed under different experimental conditions (annealing temperature, MgCl<sub>2</sub> concentration). Three distinct bands were observed in many tissues. These bands were excised, gel-purified, and sequenced. As shown in Figure 32, the KLK-L5 gene was found to have 3 molecular forms:

1) One form (referred to, from now on, as the "classical" form) represents a typical kallikrein-like serine protease with five coding exons and four intervening introns (Figure 32). As is the case with some other kallikreins, a 5' untranslated exon is also present, and the possibility of further upstream untranslated exon(s) could not be excluded. Exons 1, 2 and 3 were present at the aforementioned EST. The start codon is present in the second exon (numbers refer to SEQ.ID.NO. 56 or GenBank Accession # AF135025). The stop codon is located in the sixth exon, followed by a 3' untranslated region, and a typical polyadenylation signal (AATAAA) is located 16 bp before the poly-A tail (Figure 33). This form of KLK-L5 spans a genomic length of 5,801 bp on chromosome 19q13.3-q13.4. The lengths of the coding regions of the exons are 37, 160, 260, 134, and 156 bp, respectively (Figures 33 and 34). The predicted protein-coding region is formed of 747 bp, encoding a deduced 248-amino acid protein with a predicted molecular mass of 26.7 kDa. The intron/exon splice sites (GT....AG) and their flanking sequences are in agreement with the consensus splice site sequence.

2) The second mRNA form, encoding the KLK-L5-related protein-1, is an alternatively spliced form in which the last exon is split into two separate exons with an additional intervening intron (Figure 32). This splitting of the last exon results in the utilization of another stop codon at position 9,478, thus creating a deduced 254-amino acid protein that is 6 amino acids longer than the "classical" KLK-L5 form and its carboxyterminal end is different in sequence by 19 amino acids (Figure 32). This variant has a predicted molecular mass of 27.1 kDa (for base numbering please see SEQ.ID.NO. 56 and GenBank Accession # AF135025).

3) The third mRNA form, encoding for KLK-L5-related protein-2, is similar to the classical form except that the fourth exon is missing (Figure 32). This leads to frame shifting of the coding region, and an earlier in-frame stop codon will be encountered at position 9,180. The protein-coding region of this form

consists of 336 bp, encoding for a predicted 111-amino acid protein with a molecular mass of 12 kDa. This protein will lack both the serine and aspartate residues characteristic of serine proteases.

Amino acid sequences for KLK-L5 proteins are shown in SEQ. ID. NOs. 57 to 60.

#### Structural analysis of the classical KLK-L5 gene

5 Figure 35 shows a comparative hydrophobicity analysis of the KLK-L5 and the prostate-specific antigen (PSA) proteins. The amino terminal regions of both genes are quite hydrophobic, indicating that this region of KLK-L5 is possibly harboring a signal peptide analogous to PSA. Figure 35 also shows the presence of several evenly distributed hydrophobic regions throughout the KLK-L5 polypeptide, which are consistent with a globular protein, similar to other serine proteases (13). Figure 36 shows the alignment of  
10 KLK-L5 protein with another 10 members of the same gene family. The dotted region in Figure 36 indicates an 11-amino acid loop characteristic of the classical human kallikreins (PSA, hK1 and hK2) but not found in KLK-L5 protein or other members of the kallikrein multigene family (11, 13, 35). Sequence analysis of eukaryotic serine proteases indicates the presence of twenty nine invariant amino acids (39). Twenty eight of them are conserved in the KLK-L5 polypeptide and the remaining amino acid (S156  
15 instead of P) is not conserved among all other kallikreins (Figure 36). Twelve cysteine residues are present in the putative mature KLK-L5 protein, ten of them are conserved in all kallikreins, and the remaining two (C133, and C235 ) are present in most of the other kallikrein-like proteins but not in the classical kallikreins and they are expected to form an additional disulphide bridge (Figure 36).

The presence of aspartate (D) at position 194 suggests that KLK-L5 will possess a trypsin-like  
20 cleavage pattern, similarly to most of the other kallikreins (e.g., hK1, hK2, TLSP, neuropsin, zyme, prostase, and EMSP) but different from PSA, which has a serine (S) residue in the corresponding position, and is known to have chymotrypsin like activity (Figure 36) (54).

#### Homology with other members of the kallikrein multigene family

Although the protein encoded by the KLK-L5 gene is unique, it has a high degree of homology  
25 with the other kallikrein-like genes. The KLK-L5 protein (classical form) has 48% amino acid sequence identity and 57% overall similarity with neuropsin, 46% identity with the normal epithelial cell-specific 1 gene product (NES1) and 38% identity with both PSA and hK2 proteins. Multiple alignment shows that the typical catalytic triad of serine proteases is conserved in the KLK-L5 protein (H<sup>62</sup>, D<sup>108</sup>, and S<sup>200</sup>) (Figures 33 and 36). In addition, a well-conserved peptide motif is found around the amino acid residues  
30 of the catalytic triad as is the case with other serine proteases [i.e., histidine (VLTA~~A~~HC), serine (GDSG~~G~~P), and aspartate (DLRLL)] (11, 12) (Figure 36). Figure 36 also shows other amino acid residues that are completely conserved between kallikreins and kallikrein-like proteins. To predict the phylogenetic relatedness of the KLK-L5 protein with other serine proteases, the amino acid sequences of the kallikrein proteins were aligned together using the "Clustal X" multiple alignment program and a distance matrix tree  
35 was predicted using the Neighbor-joining/UPGMA and Protpars parsimony methods. Figure 37 shows the phylogenetic analysis which separated the classical kallikreins (hK1, hK2, and PSA) and clustered KLK-L5 with NES1 and neuropsin proteins in a separate group away from other serine proteases, consistent with previously published studies (27, 41) and indicating that this group of genes probably arose from a common

ancestral gene, by gene duplication.

#### **Tissue expression and hormonal regulation of the KLK-L5 gene**

As shown in Figure 38, the KLK-L5 gene is primarily expressed in the salivary gland, stomach, uterus, trachea, prostate, thymus, lung, colon, brain, breast and thyroid gland, but, as is the case with other kallikreins, lower levels of expression are found in some other tissues (testis, pancreas, small intestine, spinal cord). In order to verify the RT-PCR specificity, the PCR products were cloned and sequenced. The three splice forms of the gene were expressed in most of these tissues. However, the relative abundance of each form was different among tissues (Figure 38).

In order to investigate whether the KLK-L5 gene is under steroid hormone regulation, two breast cancer cell lines (BT-474 and T-47D) and a prostate cancer cell line (LNCaP) were used as models. In LNCaP cells, the gene was up-regulated only by androgen and progestin. Only in this cell line all 3 isoforms were expressed. In BT-474 cells, KLK-L5 was found to be up-regulated, at the mRNA level, by estrogen and androgen, and to a lesser extent by the progestin. The rank of potency was estrogen>androgen>progestin. However, the rank of potency for the T-47D cell line was androgen>progestin>estrogen. In both of these cell lines, only the short isoform (related protein-2) was present (Figure 39).

#### **KLK-L5 is down regulated in breast cancer**

Expression of the KLK-L5 gene, at the mRNA level, was compared between seventeen breast cancer tissues and one normal breast tissue, by RT-PCR. The gene is not expressed at all in 12 tumors (Figure 40). In all breast tissues (normal and malignant) the short isoform (related protein-2) was predominant, with the exception of one tumor, which expressed only the classical form (Figure 40, lane 8). These results should be interpreted as preliminary, since the number of tumors and normal tissues tested is relatively small.

#### **Mapping and chromosomal localization of the KLK-L5 gene**

The knowledge of extensive genomic sequence on chromosome 19q13.3-q13.4 enabled the precise localization of 14 kallikrein-like genes and determination of the direction of transcription, as shown by the arrows in Figure 28. Only PSA and KLK2 transcribe from centromere to telomere; the rest of the genes are transcribed in the reverse direction. The KLK1 gene was found to be the most centromeric, and the KLK-L6 gene the most telomeric (KLK-L6; SEQ.ID. NO.65). KLK-L5 is 21.3 Kb centromeric to KLK-L4 (SEQ.ID.NO. 43) and 1.6 kb more telomeric to the trypsin-like serine protease gene (TLSP) (Figure 28).

#### **DISCUSSION**

As shown in Figure 34, kallikreins are characterized by the following common structural features: (a) All genes are formed of 5 coding exons and 4 intervening introns [some genes may have extra 5' untranslated exon(s)] (14, 35). (b) The exon lengths are usually comparable. (c) The intron phases are always conserved (I-II-I-0) (see legend of Figure 34 for definition of intron phases). (d) These genes are clustered in the same chromosomal region, without any intervening non-kallikrein-like genes (Figure 28). (e) The histidine residue of the catalytic triad of serine proteases is located near the end of the second coding exon; the aspartate residue in the middle of the third coding exon; and serine at the beginning of the

fifth coding exon. As shown in Figure 34, all these criteria apply to the newly identified KLK-L5 gene. Thus, KLK-L5 should be considered a new member of the kallikrein multigene family.

Serine proteases and kallikreins are synthesized as "preproenzymes" that contain an N-terminal signal peptide (pre-zymogen), followed by a short activation peptide and the enzymatic domain (41, 54). PreproPSA has 24 additional residues that constitute the pre-region (signal peptide, 17 residues), and the propeptide (7 residues) (67). The signal peptide directs the protein to and across the endoplasmic reticulum (ER). In the ER, the signal peptide is removed and the resulting proPSA is transported to the plasma membrane, where it is secreted. The hydrophobicity study (Figure 35) indicates that the amino terminal region of the KLK-L5 protein is harboring a signal peptide. Also, computer analysis of the amino acid sequence of KLK-L5 predicted a cleavage site between amino acids 17 and 18 (SQA-AT). Thus, the protein product is very likely a secreted protein.

The presence of alternatively spliced forms is not a unique feature of the KLK-L5 gene; several other kallikreins are known to be expressed in various alternatively spliced forms. In addition to the major 1.6-kb transcript, several RNA species are transcribed from the PSA gene (61). Furthermore, others (69, 70) have described several PSA isoforms. Retained introns and loss of complete exons have been reported in some of these forms. In addition, Riegman et al. reported the identification of two alternatively spliced forms of the human glandular kallikrein (KLK2) gene (62) and Liu et al. isolated three alternative forms of the same gene (68). A novel transcript of the tissue kallikrein gene was isolated from the colon (63). Neuropsin, a recently identified kallikrein-like gene, was found to have two alternatively spliced forms in addition to the major form (35, 64). Here, the cloning of the classical kallikrein form and two unique splice forms of the KLK-L5 gene are described. Because the classical form and the splice forms all have the same 5' sequence required for translation, secretion and activation as do other kallikreins, i.e. a 5' leader sequence, a signal peptide, and a proregion, it is reasonable to assume that all three mRNA forms should produce a secreted protein. Preliminary findings identifying forms of KLK-L5 predominant in certain tissues are presented in Figures 35 and 40.

The preliminary results indicate that KLK-L5 is up-regulated by steroid hormones in breast and prostate cancer cell lines (Figure 39). These results are not surprising, since many other kallikrein genes are also regulated by steroid hormones. The differences in the rank of potency of steroid hormones among different cell lines could be attributed to differences in the abundance of the steroid hormone receptors between them as described elsewhere.

In conclusion, a new member of the human kallikrein gene family, KLK-L5, has been characterized which maps to the human kallikrein locus (chromosome 19q13.3-q13.4). This gene has two related splice forms in addition to the main form. KLK-L5 is expressed in a variety of tissues, appears to be down-regulated in breast cancer and its expression is influenced by steroid hormones. Since a few other kallikreins are already used as valuable tumour markers, KLK-L5 may also find a similar clinical application.

#### Example 7

Using the Materials and Methods substantially as set out in Example 6, the present inventors

identified another novel gene of the kallikrein multigen family referred to as KLK-L6. The full structure of the KLK-L6 gene is shown in Figure 41. Exons 1, 2, 3, 4, 5, 6, and 7 are at nucleic acids 1172-1281; 2561-2695; 2781-2842, 3714-3885; 5715-5968; 6466-6602; and 7258-7520. The nucleic acid sequence of the KLK-L6 gene is also shown in SEQ.ID.NO. 65 and amino acid sequences for the KLK-L6 protein are shown in SEQ.ID. Nos. 66 and 67. (See also GenBank Accession # AF161221).

Figure 42 shows a comparative hydrophobicity analysis of KLK-L6 and the prostate-specific antigen (PSA). The amino terminal regions of both genes are quite hydrophobic indicating that this region of KLK-L6 is possibly harboring a signal peptide analogous to PSA

Multiple alignment of KLK-L6 was carried out using the Clustal X software program as described herein (Figure 43).

To predict the phylogenetic relatedness of the KLK-L6 protein with other serine proteases, the amino acid sequences of the kallikrein proteins were aligned together using the "Clustal X" multiple alignment program and a distance matrix tree was predicted using the Neighbor-joining/UPGMA and Protpars parsimony methods. Figure 44 shows the phylogenetic analysis which separated the classical kallikreins (hK1, hK2, and PSA) and placed KLK-L6 in a separate group.

Having illustrated and described the principles of the invention in a preferred embodiment, it should be appreciated to those skilled in the art that the invention can be modified in arrangement and detail without departure from such principles. All modifications coming within the scope of the following claims are claimed.

All publications, patents and patent applications referred to herein are incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

**FULL CITATIONS FOR REFERENCES REFERRED TO IN THE SPECIFICATION**

1. Evans BAE, Yun ZX, Close JA, Tregear GW, Kitamura N, Nakanishi S, et al. Structure and chromosomal localization of the human renal kallikrein gene. *Biochemistry* 1988;27:3124-3129.
2. Clements JA. The glandular kallikrein family of enzymes: Tissue-specific and hormonal regulation. *Endocr Rev* 1989;10:393-419.
3. Evans BA, Drinkwater CC, Richards RI. Mouse glandular kallikrein genes: Structure and partial sequence analysis of the kallikrein gene locus. *J Biol Chem* 1987;262:8027-8034.
4. Drinkwater CC, Evans BA, Richards RI. Kallikreins, kinins and growth factor biosynthesis. *Trends Biochem Sci* 1988b;13:169-172.
5. Ashley PL, MacDonald RJ. Tissue-specific expression of kallikrein-related genes in the rat. *Biochemistry* 1985;24:5420-5427.
6. Gerald WL, Chao J, Chao L. Sex dimorphism and hormonal regulation of rat tissue kallikrein mRNA. *Biochim Biophys Acta* 1986;867:16-23.
7. Riegman PHJ, Vlietstra RJ, van der Korput JAGM, Romijn JC, Trapman J. Characterization of the prostate-specific antigen gene: A novel human kallikrein-like gene. *Biochem Biophys Res Commun* 1989;159:95-102.
8. Schedlich LJ, Bennetts BH, Morris BJ. Primary structure of a human glandular kallikrein gene. *DNA* 1987;6:429-437.
9. Riegman PH, Vlietstra RJ, Suurmeijer L, Cleutjens CBJM, Trapman J. Characterization of the human kallikrein locus. *Genomics* 1992;14:6-11.
10. Anisowicz A, Sotiropoulou G, Stenman G, Mok SC, Sager R. A novel protease homolog differentially expressed in breast and ovarian cancer. *Mol Med* 1996;2:624-636.
11. Little SP, Dixon EP, Norris F, Buckley W, Becker GW, Johnson M, et al. Zyme, a novel and potentially amyloidogenic enzyme cDNA isolated from Alzheimer's disease brain. *J Biol Chem* 1997;272:25135-25142.
12. Yamashiro K, Tsuruoka N, Kodama S, Tsujimoto M, Yamamura Y, Tanaka T, et al. Molecular cloning of a novel trypsin-like serine protease (neurosin) preferentially expressed in brain. *Biochim Biophys Acta* 1997;1350:11-14.
13. Liu XL, Wazer DE, Watanabe K, Band V. Identification of a novel serine protease-like gene, the expression of which is down-regulated during breast cancer progression. *Cancer Res* 1996;56:3371-3379.
14. Luo L, Herbrick J-A, Scherer SW, Beatty B, Squire J, Diamandis EP. Structural characterization and mapping of the normal epithelial cell-specific 1 gene. *Biochem Biophys Res Commun* 1998;247:580-586.
15. Milanesi L, Kolchanov N, Rogozin I, Kel A, Titov I. Sequence functional inference. In: "Guide to human genome computing", ed. M.J. Bishop, Academic Press, Cambridge, 1994, 249-312.
16. Burset M, Guigo R. Evaluation of gene structure prediction programs. *Genomics* 1996;34:353-367.

17. Nadeau J, Grant P, Kosowsky M. Mouse and human homology map. *Mouse Genome* 1991;89:31-36.
18. Schachter M. Kallikreins (kininogenases) - a group of serine proteases with bioregulatory actions. *Pharmacol Rev* 1980;31:1-17.
- 5 19. Morris BJ, Catanzaro DF, Richards RI, Mason AJ, Shine J. Kallikrein and renin: Molecular biology and biosynthesis. *Clin Sci* 1981;61:351s-353s.
20. Richards RI, Catanzaro DF, Mason AJ, Morris BJ, Baxter JD, Shine J. Mouse glandular kallikrein genes. Nucleotide sequence of cloned cDNA coding for a member of the kallikrein arginyl ester-peptidase group of serine proteases. *J Biol Chem* 1982;257:2758-2761.
- 10 21. Van Leeuwen BH, Evans BA, Tregear GW, Richards RI. Mouse glandular kallikrein genes. Identification, structure and expression of the renal kallikrein gene. *J Biol Chem* 1986; 261:5529-5535.
22. Evans BA, Richards RI. Genes for the  $\alpha$  and  $\gamma$  subunits of mouse nerve growth factor. *EMBO J* 1985; 4:133-138.
- 15 23. Rogozin IB, Milanesi L, Kolchanov NA. Gene structure prediction using information on homologous protein sequence. *Comput Applic Biosci* 1996;12:161-170.
24. Diamandis, E.P. Prostate specific antigen-its usefulness in clinical medicine. *Trends Endocrinol. Metab.*, 9: 310-316, 1998.
25. Diamandis, E. P., Yu H., and Sutherland, D.J. Detection of prostate-specific antigen immunoreactivity in breast tumours. *Breast Cancer Res.Treat.*, 32: 301-310, 1994
- 20 26. Ishikawa, T., Kashiwagi, H., Iwakami, Y., et al. Expression of alpha-fetoprotein and prostate specific antigen genes in several tissues and detection of mRNAs in normal circulating blood by reverse transcriptase-polymerase chain reaction. *Jpn. J. Oncol.*, 28:723-728, 1998.
- 25 27. Irwin, D.M., Robertson, K.A., and MacGillivray, R.T. *J.Mol.Biol.*212:31-45, 1988.
28. Yoshida, S., Taniguchi, M., Hirata, A., and Shiosaka, S. Sequence bovine prothrombin gene. *J. Mol. Biol.*, 212: 31-45, 1988.
29. Goyal, J., Smith, K.M., Cowan, J.M., et al. The role of NES1 serine protease as a novel tumor suppressor. *Cancer Res.*, 58: 4782-4786, 1998.
- 30 30. Diamandis, E.P., and Yu, H. New biological functions of prostate specific antigen? *J. Clin. Endocrinol. Metab.*, 80 : 1515-1517, 1995.
31. Reifenberger, J., reifenberger, G., Liu, L., James, C.D. et al. Molecular genetic analysis of oligodendroglial tumors shows preferential allelic deletions on 19q and 1p.*Am. J. Pathol.*,145: 1175-90, 1994.
- 35 32. Iida, Y. (1990). Quantification analysis of 5'-splice signal sequence in mRNA precursors. Mutations in 5'-splice signal sequence of human  $\beta$ -globin gene and  $\beta$ -thalassemia. *J. Theor. Biol.* 145: 523-533.
33. Kozak, M. (1991). An analysis of vertebrate mRNA sequences: Intimations of translational

- control. *J. Cell Biol.* 115: 887-892.
34. Clements, J. (1997). The molecular biology of the kallikreins and their roles in inflammation. *In*: S. Farmer (ed.), *The kinin system*, pp. 71-97. New York: Academic Press.
  35. Yoshida, S., Taniguchi, M., Hirata, A., and Shiosaka, S. (1998). Sequence analysis and expression of human neuropsin cDNA and gene. *Gene* 213 :9-16.
  36. Takayama, T. K., Fujikawa, K., Davie, E. W. (1997). Characterization of the precursor of prostate-specific antigen. Activation by trypsin and by human glandular kallikrein. *J. Biol. Chem.* 272: 21582-21588.
  37. Altschul, S.F. et al., *Nucleic Acids Res.* 25: 3389-3402, 1997.
  38. Lennon, G. et al, *Genomics* 33: 151-152, 1996.
  39. Dayhoff, M. O., *Natl. Biomed. Res. Found.* 5 (Suppl 3) 79-81, 1998.
  40. Simmer, J.P., et al, *J. Dent. Res.* 77:377-386, 1998.
  41. Nelson, P.S. et al, *PNAS* 96: 3114-3119, 1999.
  42. Osoegawa, K. et al, *Genomics* 52: 1-8, 1999.
  43. Kozak, M., *Nucleic Acid Res.* 15: 8125-8148, 1987.
  44. Proudfoot, N.J. and Brownlee, C.G., *Nature* 263: 211-214, 1976.
  45. Miyata, T. et al, *J. Mol. Evol.* 12: 219-236, 1979.
  46. Mitelman, F., *Catalog of Chromosome Aberrations in Cancer*, 5<sup>th</sup> ed. Wiley-Liss, New York, pp. 3067-3198.
  47. Yu, H. et al, *Clin. Cancer res.* 4: 1489-1497, 1998.
  48. Sauter, E.R., *Cancer Epidemiol. Biomarkers Prev.* 5:967-970, 1996.
  49. Fortier, A.H. et al, *J. Natl. Cancer Inst.* 91: 1635-1640, 1999.
  50. Kumar, A., 1998, *Cancer res.* 57: 3111-3114, 1997.
  51. Lovgren, J. *Biochem. Biophys. Res. Commun.* 238: 549-555, 1987.
  52. Lai, L.C. et al, *Int. J. Cancer* 66: 743-746, 1996.
  53. Balbay, M.D. et al, *Proc. Amer. Assoc. Cancer Res.* 40: 225-226, 1999.
  54. Rittenhouse, H.G. et al *Crit. Rev. Clin. Lab. Sci.* 35: 275-368, 1998.
  55. Hansson, L. et al, *J. Biol. Chem.* 269: 19420-19426, 1994.
  56. Stephenson, S. et al *J. Biol. Chem.* 27: 23210-23214, 1999.
  57. Stenman, U.H. *Clin. Chem.* 45: 753-754, 1999.
  58. Black, M.H., *Clin. Chem.* 45: 790-799, 1999.
  59. Underwood, L.J. et al, *Cancer Res.* 59:4435-4439, 1999.
  60. Adams, M.D. et al, *Curr. Opin. Cell Biol.* 8:331-339, 1996.
  61. Heuze, N. et al, *cancer Res.* 59: 2820-2824, 1999.
  62. Riegman, P.H., *Mol. Cell Endocrinol.* 76: 181-190.
  63. Chen, L.M. *Braz. J. Med. Biol Res.* 27: 1829-1838, 1994.
  64. Mitsui, S. et al, *Eur. J. Biochem.* 260:627-634.
  65. Baffa, R., *Urology*, 47:795-800,1996.

66. Henttu, P. et al, *Int. J. Cancer* 45: 654-660, 1990.
67. McCormack, R.T. et al, *Urology* 45: 729-744, 1995.
68. Liu, X.F., et al, *Biochem. Biophys. Res. Commun.* 264:833-839, 1999.
69. Riegman, P.H. et al, *Biochem. Biophys. Res. Commun.* 155: 181-188, 1988.
- 5 70. Lundwall, A. and H. Lilja, *FEBS Lett.* 214: 317-322, 1987.
71. Zarghami, N. et al, *Br. J. Cancer* 75: 579-88, 1997.

TABLE 1. Predicted exons of the putative gene KLK-L1. The translated protein sequences of each exon (open reading frame) are shown.

Exon No. <sup>1</sup>	Putative coding region <sup>2</sup>	No. of bases	Translated protein sequence	EST match <sup>3</sup>	Intron phase <sup>4</sup>	Stop codon <sup>5</sup>	Catalytic triad <sup>6</sup>	Exon prediction program <sup>7</sup>
2	From(bp) 2263 To(bp) 2425	163	SLVSGSCQIINGEDCSPHSQPWQAALVMENELFCGV LVHPQWVLSAAHCFQ	+	II	-	H	A,B,D
3	2847 3109	263	NSYTGGLHLSLEADQEPGSMVEASLSVRHPEYNRPL LANDLMILKLDSESVSESDTIRISIASQCPTAGNSCLVSG WGLLANGELT	+	I	-	D	A,B,C,D
4	3180 3317	137	GRMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGG GQDQKDCN	+	0	-		A,B,C,D
5	4588 4737	150	GDGGGPLICNGYLQGLVSFGKAPCGQGVGPVYTNLG KFTIEWIEKTVQAS	+	-	+	S	A,B,C

1. Conventional numbering of exons in comparison to the five coding exons of PSA, as described in Ref.14.

2. Nucleotide numbers refer to the related contig

3. (+) = >95% homology with published human EST sequences.

4. Intron phase: 0 = the intron occurs between codons; I = the intron occurs after the first nucleotide of the codon;

II = the intron occurs after the second nucleotide of the codon.

5. (+) denotes the exon containing the stop codon.

6. H = histidine, D = aspartic acid, S = serine. The amino acids of the catalytic triad are bold and underlined.

7. A = GeneBuilder (gene analysis), B = GeneBuilder (exon analysis), C = Grail 2,

D = GENEID-3

TABLE 2. Predicted exons of the putative gene KLK-L2. The translated protein sequences of each exon (open reading frame) are shown.\*

Exon No. <sup>1</sup>	Putative coding sequence <sup>2</sup> From(bp) To(bp)	No. of bases	Translated protein sequence	EST match <sup>3</sup>	Intron phase <sup>4</sup>	Stop codon <sup>5</sup>	Catalytic triad <sup>6</sup>	Exon prediction program <sup>7</sup>
1	15,361 15,433	73	MATARPPWMWVLCALITALLGVT	+	I	-	-	-
2	17,904 18,165	262	EHVLANNVSDHPSNTVPSGNSQDLGAGAGEDARSDSSRIIN GSDCDMHTQPWQAALLLRPNQLYCGA VLHPQWLLTAAHCRK K	+	II	-	H	A,B,C,D
3	18,903 19,159	257	VPRVLGHYSLSPVYESGQQMFQGVKSIHPGYSHPGHSNDLMLI KLNRRIRPTKDYRPINVSCHCPSAGTKCLVSGWGTTKSPQ	+	I	-	D	C,D
4	19,245 19,378	134	VHFPKVLQCLNISVLSQKRCEDAYPRQIDDTMFCAGDKAGRDC Q	+	0	-	-	B,C
5	24,232 24,384	153	GDGGPVVVCNGSLQGLVSWGDYPCARPNRPGVYTNLCKFTKWI QETIQANS	+	-	+	S	A,B,C

\* All footnotes same as Table 1

TABLE 3. Predicted exons of the putative gene KLK-L3. The translated protein sequences of each exon (open reading frame) are shown

Exon No. <sup>1</sup>	Putative coding region <sup>2</sup> From(bp) To(bp)	No. of bases	Translated protein sequence	EST match <sup>3</sup>	Intron phase <sup>4</sup>	Stop codon <sup>5</sup>	Catalytic triad <sup>6</sup>	Exon prediction program <sup>7</sup>
1	70,473 70,584	112	VHFTPINHRGGPMEEEGDGMAYHKEALDAGCTFQDP	-	I	-	-	A,B,C,D
2	70,764 70,962	199	ACSSLTPLSLIPTPGHGWADTRAIGAEECRPN SQPWQAGLF HLTRLFCGATLISDRWLLTAAHCRK	+	II	-	H	A,B,C,D
3	73,395 73,687	293	PLTSEACFSRYLVWRLGEHLLWKWEGPEQLFRVTDFFPH GFNKDLSANDHNDIMLIRLPRQARLSPAVQPLNLSQTCV SPGMQCLISGWGAVSPK	+	I	-	D	A,B,C,D
4	76,305 76,441	137	ALFPVTLQCANISILENKLCHWAYPGHISDSMLCAGLWEG GRGSCQ	+	0	-	-	A,B,C,D
5	76,884 77,633	749	GDGGPLVCNGTLAGVVGGAEPCSRPRRPAVYTSVCHYL DWIQEIMEN	-	-	+	S	A,B

\* All footnotes same as Table 1

TABLE 4. Predicted exons of the putative gene KLK-L4. The translated protein sequences of each exon (open reading frame) are shown

Exon No. <sup>1</sup>	Putative coding region <sup>2</sup> From(bp) To(bp)	No. of bases	Translated protein sequence	EST match <sup>3</sup>	Intron phase <sup>4</sup>	Stop codon <sup>5</sup>	Catalytic triad <sup>6</sup>	Exon prediction program <sup>7</sup>
2	24,945 25,120	176	ESSKVLNTNGTSGFLPGGYTCFPHSQPWQAALLVQGRLL CGGVLVHPKWVLTAAHCLKE	+	II	-	H	C
3	25,460 25,728	269	GLKVYLGKHALGRVEAGEQVREVVHSIPHEYYRRSPTHL NHDHIMLLELQSPVQLTGYYIQLPLSHNNRLTPGTTTCRV SGWGTTTSPQ	+	I	-	D	A,B,C,D
4	26,879 27,015	137	VNYPKTLQCANIQLRSDEECRQVYPGKITDNMLCAGTKE GGKDSCE	+	0	-	-	A,B,C,D
5	28,778 28,963	189	GDSGGPLVCNRTLYGIVSWGDFPCGQFDRPGVYTRVSRV VLWIRETIRKYETQQQKWLKGPQ	+	-	+	S	A,B,C

\* All footnotes same as Table 1

TABLE 5. Predicted exons of the putative gene KLK-L5. The translated protein sequences of each exon (open reading frame) are shown

Exon No. <sup>1</sup>	Putative coding region <sup>2</sup> From(bp) To(bp)	No. of bases	Translated protein sequence	EST match <sup>3</sup>	Intron phase <sup>4</sup>	Stop codon <sup>5</sup>	Catalytic triad <sup>6</sup>	Exon prediction program <sup>7</sup>
2	1588 1747	160	LSQAATPKIFNGTECGRNSQPWQVGLFEGTSLRCGGV LIDHRWVL TAAHCSG	-	II	-	H	A,B,C
3	3592 3851	260	SRYWVRLGEHSLQLDWTEQIRHSGFSVTHPGYLGAS TSHEHDLRLRLRPVRTSSVQPLPLPNDCATAGTEC HVSQGWGITNHPR	+	I	-	D	A,B,C,D
4	4806 4939	134	NPPFDLLQCLNLSIVSHATCHGVYPORITSNMVCAGG VPGQDACQ	+	0	-	-	A,B,C,D

\* All footnotes same as Table 1

TABLE 6. Homology between the predicted amino acid sequences of the newly identified putative genes and protein sequences deposited in Genbank

No.	Gene identity	Homologous known protein	Identity% (number of amino acids)
1	KLK-L1	• Human stratum corneum chymotryptic enzyme	44(101/227)
		• Rat kallikrein	40( 96/237)
		• Mouse glandular kallikrein K22	39( 94/236)
		• Human glandular kallikrein	38( 93/241)
		• Human prostatic specific antigen	37( 91/241)
		• Human protease M	37( 87/229)
2	KLK-L2	• Human neuropsin	48(106/219)
		• Human stratum corneum chymotryptic enzyme	47(103/216)
		• Human protease M	45( 99/219)
		• Human trypsinogen I	45(100/221)
		• Rat trypsinogen	44( 98/220)
3	KLK-L3	• Human neuropsin	44(109/244)
		• Rat trypsinogen 4	39( 95/241)
		• Human protease M	38( 98/253)
		• Human glandular kallikrein	37( 94/248)
		• Human prostatic specific antigen	36( 89/242)
4	KLK-L4	• Human protease M	52(118/225)
		• Human neuropsin	51(116/225)
		• Mouse neuropsin	51(116/226)
		• Human glandular kallikrein	48(113/234)
		• Human prostatic specific antigen	47(108/227)
5	KLK-L5	• Human neuropsin	44( 81/184)
		• Rat trypsinogen I	42( 76/178)
		• Rat trypsinogen II	42( 75/178)
		• Human protease M	41( 73/178)

- 72 -

**Table 7.** Expressed sequence tags with >95% homology to exons of the prostate/KLK-L1 gene.

GenBank #	Source	Tissue	homologous exons
AA551449	I.M.A.G.E.	prostate	3,4,5
AA533140	I.M.A.G.E.	prostate	4,5
AA503963	I.M.A.G.E.	prostate	5
AA569484	I.M.A.G.E.	prostate	5
AA336074	TIGR	endometrium	2,3

**Table 8.** Primers used for reverse transcription-polymerase chain reaction (RT-PCR) analysis of various genes.

Gene	Primer name	Sequence <sup>1</sup>	Product size (base pairs)
Protease (KLK-L1)	RS	TGACCCGCTGTACCACCCCA	278
	RAS	GAATTCCTTCCGCAGGATGT	
pS2	PS2S	GGTGATCTGCGCCCTGGTCCT	328
	PS2AS	AGGTGTCCGGTGGAGGTGGCA	
PSA	PSAS	TGCGCAAGTTCACCCTCA	754
	PSAAS	CCCTCTCCTTACTTCATCC	
Actin	ACTINS	ACAATGAGCTGCGTGTGGCT	372
	ACTINAS	TCTCCTTAATGTCACGCACGA	

1. All nucleotide sequences are given in the 5'→3' orientation.

Table 9. Tissue expression of prostate/CLK-L1 by RT-PCR analysis

Expression level			
High	medium	low	No Expression
Prostate	Mammary gland	Salivary glands	Stomach
Testis	Colon	Lung	Heart
Adrenals	Spinal cord	Brain	Spleen
Uterus		Bone marrow	Placenta
Thyroid		Thymus	Liver
		Trachea	Pancreas
		Cerebellum	Kidney
			Fetal brain
			Fetal liver
			Skeletal muscle
			Small intestine

**Table 10.** EST clones with >95% homology to exons of KLK-L2

GENBANK #	Tissue of Origin	I.M.A.G.E. ID	Homologous exons
W73140	Fetal heart	344588	4,5
W73168	Fetal heart	344588	3,4,5
AA862032	Squamous cell carcinoma	1485736	4,5
AI002163	Testis	1619481	3,4,5
N80762	Fetal lung	300611	5
W68361	Fetal heart	342591	5
W68496	Fetal heart	342591	5
AA292366	Ovarian tumor	725905	1,2
AA394040	Ovarian tumor	726001	5

**Table 11.** Primers used for reverse transcription polymerase chain reaction (RT-PCR) analysis.

Gene	Primer name	Sequence <sup>1</sup>	Product size (base pairs)
KLK-L2	KS	GGATGCTTACCCGAGACAGA	342
	KAS	GCTGGAGAGATGAACATTCT	
pS2	PS2S	GGTGATCTGCGCCCTGGTCCT	328
	PS2AS	AGGTGTCCGGTGGAGGTGGCA	
PSA	PSAS	TGCGCAAGTTCACCCTCA	754
	PSAAS	CCCTCTCCTTACTTCATCC	
Actin	ACTINS	ACAATGAGCTGCGTGTGGCT	372
	ACTINAS	TCTCCTTAATGTCACGCACGA	
KLK-L2	R1	CCGAGACGGACTCTGAAAACTTTCTTCC	
	R2	TGAAAACTTTCTTCTCCTGCAGTGGGCGGC	

1. All nucleotide sequence are given in the 5' 3' orientation.

**Table 12.** Tissue expression of KLK-L2 by RT-PCR analysis.

<b>sion level</b>			
<b>high</b>	<b>Medium</b>	<b>low</b>	<b>No Expression</b>
Brain	Salivary gland	Uterus	Stomach
Mammary gland	Fetal brain	Lung	Adrenal gland
Testis	Thymus	Heart	Colon
	Prostate	Fetal liver	Skeletal muscle
	Thyroid	Spleen	
	Trachea	Placenta	
	Cerebellum	Liver	
	Spinal cord	Pancreas	
		Small intestine	
		Kidney	
		Bone marrow	

TABLE 13. Primers used for reverse transcription polymerase chain reaction (RT-PCR) analysis.

Gene	Primer name	Sequence <sup>1</sup>
KLK-L3	L3-F1	CATGCAGTGTCTCATCTCAG
	L3-F2	CATGGAGGAGGAAGGAGATG
	L3-R1	CTTCGGCCTCTCTTGGTCTT
PSA	PSAS	TGCGCAAGTTCACCCTCA
	PSAAS	CCCTCTCCTTACTTCATCC
Actin	ACTINS	ACAATGAGCTGCGTGTGGCT
	ACTINAS	TCTCCTTAATGTCACGCACGA

1. All nucleotide sequence are given in the 5' → 3' orientation.

TABLE 14. Primers used for gene-specific PCR amplification of the kallikrein genes using DNA as a template.

Primer name	Sequence <sup>1</sup>	Coordinates	GenBank accession #	Gene name
ZIS	GACCCTGACATGTGACATCTA	979-999	U62801	Zyme
ZIAS	GCCACTGCCTGATGGAGACTG	1422-1402		
GL3-F1	AACATCAGCATCCTGGAGAA	7324-7343	AF135026	KLK-L3
LL3-R1	CTTCGGCCTCTCTTGGTCTT	8051-8060		
L2-1	GGGTCAGAGCTGCAGAGAAG	11104-11123	AF135028	KLK-L2
L2-2	GGGCCTGTCGTCTGCAATGG	11522-11541		
KLK-L1	ATGGCCACAGCAGGAAATCC	1411-1430	AF113141	KLK-L1
	GGTCACTTGTCTGCGCAGAC	1990-2019		
PS	CCCAACCCTGTGTTTTTCTC	3634-3653	M33105	PSA <sup>2</sup>
PAS	GGCCCTCCTCCCTCAGA	4143-4118		
K1S	ATCCCTCCATTCCCATCTTT	2-22	M18157	KLK1 <sup>3</sup>
K1AS	CACATACAATTCTCTGGTTC	324-30		
K2S	AGTGACACTGTCTCAGAAAT	131-150	AF024605	KLK2 <sup>4</sup>
K2AS	CCCAATCTCACCAGTGCAC	580-561		
NS	GCTTCCCTACCGCTGTGCT	552-570	AF055481	NES1 <sup>5</sup>
NAS	CACTCTGGCAAGGGTCCTG	763-744		

1. all nucleotide sequences are given in the 5' → 3' orientation
2. prostate specific antigen
3. human renal kallikrein
4. human glandular kallikrein
5. normal epithelia cell-specific 1 gene.

TABLE 15. Primers used for reverse transcription polymerase chain reaction (RT-PCR) analysis.

Gene	Primer name	Sequence <sup>1</sup>
KLK-L4	L4-F1	AACTCTACAATGTGCCAACA
	L4-R1	TTATTGTGGGCCCTTCAACC
	L4-R3	GGATGGTCCATTTATAGGAC
	L4-A	AGGCTGCCCTACTAGTGCAA
	L4-B	ATATTGCCTAGGTGGATGTG
	L4-D	AAGACTTCAAGGAGCCAAGC
	L4-E	GACCCTTCACCTCCCAAAT
	L4-X1	CTAGTGATCGCCTCCCTGAC
	PS2S	GGTGATCTGCGCCCTGGTCCT
	PS2AS	AGGTGTCCGGTGGAGGTGGCA
PSA	PSAS	TGCGCAAGTTCACCCTCA
	PSAAS	CCCTCTCCTTACTTCATCC
Actin	ACTINS	ACAATGAGCTGCGTGTGGCT
	ACTINAS	TCTCCTTAATGTCACGCACGA

1. All nucleotide sequence are given in the 5'→3' orientation.

TABLE 16. EST clones with &gt;95% homology to exons of KLK-L4

GenBank #	Tissue of origin	I.M.A.G.E. ID
AA399955 AA401397	Testis	743113
AA846771	Testis	1392889
AI002101	Testis	1619045
AI032327	Testis	1644236

TABLE 17. Primers used for reverse transcription polymerase chain reaction (RT-PCR) analysis.

Gene	Primer name	Sequence <sup>1</sup>
KLK-L5	KLK-L5-F1	TCAGCCAGGCAGCCACACCG
	KLK-L5-R1	TTGGTGATGCCCCAGCCTGA
	L5-F2	CCACACCGAAGATTTTCAAT
	L5-R2	GCCCCTCCTTCATTTATA
PSA	PSAS	TGCGCAAGTTCACCCTCA
	PSAAS	CCCTCTCCTTACTTCATCC
Actin	ACTINS	ACAATGAGCTGCGTGTGGCT
	ACTINAS	TCTCCTTAATGTCACGCACGA

1. All nucleotide sequence are given in the 5'→3' orientation.

We Claim:

1. An isolated KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 nucleic acid molecule of at least 30 nucleotides which hybridizes to SEQ ID NO. 1, 13, 21, 43, 56, or 65, respectively, or the complement of SEQ ID NO. 1, 13, 21, 43, 56, or 65, under stringent hybridization conditions
2. An isolated nucleic acid molecule which comprises:
  - (i) a nucleic acid sequence encoding a protein having substantial sequence identity with an amino acid sequence of a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively;
  - (ii) a nucleic acid sequence encoding a protein comprising an amino acid sequence of a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively;
  - (iii) nucleic acid sequences complementary to (i);
  - (iv) a degenerate form of a nucleic acid sequence of (i);
  - (v) a nucleic acid sequence capable of hybridizing under stringent conditions to a nucleic acid sequence in (i), (ii) or (iii);
  - (vi) a nucleic acid sequence encoding a truncation, an analog, an allelic or species variation of a protein comprising an amino acid sequence of a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein as shown in SEQ.ID.NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67, respectively; or
  - (vii) a fragment, or allelic or species variation of (i), (ii) or (iii).
3. A purified and isolated nucleic acid molecule of the invention comprises:
  - (i) a nucleic acid sequence comprising the sequence of SEQ.ID.NO. 1, 13, 21, 43, 56, or 65 wherein T can also be U;
  - (ii) nucleic acid sequences complementary to (i), preferably complementary to the full nucleic acid sequence of SEQ.ID.NO. 1, 13, 21, 43, 56, or 65;
  - (iii) a nucleic acid capable of hybridizing under stringent conditions to a nucleic acid of (i) or (ii) and preferably having at least 18 nucleotides; or
  - (iv) a nucleic acid molecule differing from any of the nucleic acids of (i) to (iii) in codon sequences due to the degeneracy of the genetic code.
4. An isolated nucleic acid molecule which encodes a protein which binds an antibody of a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein.
5. A regulatory sequence of an isolated nucleic acid molecule as claimed in any of the preceding claims fused to a nucleic acid which encodes a heterologous protein.
6. A vector comprising a nucleic acid molecule of any of the preceding claims.
7. A host cell comprising a nucleic acid molecule of any of the preceding claims.

8. An isolated KLK-L1 protein comprising an amino acid sequence of SEQ. ID. NO. 2 or 3.
9. An isolated KLK-L2 protein comprising an amino acid sequence of SEQ. ID. NO. 14.
10. An isolated KLK-L3 protein comprising an amino acid sequence of SEQ. ID. NO. 22 or 23.
11. An isolated KLK-L4 protein comprising an amino acid sequence of SEQ. ID. NO. 44 or 45.
- 5 12. An isolated KLK-L5 protein comprising an amino acid sequence of SEQ. ID. NO. 57, 58, 59, or 60.
13. An isolated KLK-L6 protein comprising an amino acid sequence of SEQ. ID. NO. 66 or 67.
14. An isolated protein having at least 65% amino acid sequence identity to an amino acid sequence of SEQ. ID. NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67.
15. A method for preparing a protein as claimed in any of the preceding claims comprising:
  - 10 (a) transferring a vector as claimed in claim 6 into a host cell;
  - (b) selecting transformed host cells from untransformed host cells;
  - (c) culturing a selected transformed host cell under conditions which allow expression of the protein; and
  - (e) isolating the protein.
- 15 16. A protein prepared in accordance with the method of claim 15.
17. An antibody having specificity against an epitope of a polypeptide as claimed in claim 8, 9, 10, 11, 12, or 13.
18. An antibody as claimed in claim 17 labeled with a detectable substance and used to detect the protein in biological samples, tissues, and cells.
- 20 19. A probe comprising a sequence encoding a protein as claimed in claim 8, 9, 10, 11, 12, or 13, or a part thereof.
20. A method of diagnosing and monitoring conditions mediated by a protein as claimed in claim 8, 9, 10, 11, 12, or 13, by determining the presence of a nucleic acid molecule encoding the protein as claimed in any of the preceding claims or determining the presence of the protein.
- 25 21. A method as claimed in claim 20 wherein the condition is cancer.
22. A method for identifying a substance which associates with a protein as claimed in claim 8, 9, 10, 11, 12, or 13 comprising (a) reacting the protein with at least one substance which potentially can associate with the protein, under conditions which permit the association between the substance and protein, and (b) removing or detecting protein associated with the substance, wherein detection of associated protein and substance indicates the substance associates with the protein.
- 30 23. A method for evaluating a compound for its ability to modulate the biological activity of a protein as claimed in claim 8, 9, 10, 11, 12, or 13 comprising providing a known concentration of the protein with a substance which associates with the protein and a test compound under conditions which permit the formation of complexes between the substance and protein, and removing and/or detecting complexes.
- 35 24. A method for detecting a nucleic acid molecule encoding a protein comprising an amino acid sequence of SEQ. ID. NO. 2, 3, 14, 22, 23, 44, 45, 57, 58, 59, 60, 66, or 67 in a biological sample comprising the steps of:

- 85 -

- (a) hybridizing a nucleic acid molecule of claim 2 to nucleic acids of the biological sample, thereby forming a hybridization complex; and
- (b) detecting the hybridization complex wherein the presence of the hybridization complex correlates with the presence of a nucleic acid molecule encoding the protein in the biological sample.
- 5
25. A method as claimed in claim 24 wherein nucleic acids of the biological sample are amplified by the polymerase chain reaction prior to the hybridizing step.
26. A method for treating a condition mediated by a protein as claimed in claim 8, 9, 10, 11, 12, or 13 comprising administering an effective amount of an antibody as claimed in claim 17 or a substance or compound identified in accordance with a method claimed in claim 22 or 23.
- 10
27. A method as claimed in claim 26 wherein the condition is cancer.
28. A composition comprising one or more of a nucleic acid molecule or protein claimed in any of the preceding claims, or a substance or compound identified using a method as claimed in any of the preceding claims, and a pharmaceutically acceptable carrier, excipient or diluent.
- 15
29. Use of one or more of a nucleic acid molecule or protein claimed in any of the preceding claims, or a substance or compound identified using a method as claimed in any of the preceding claims in the preparation of a pharmaceutical composition for treating a condition mediated by a protein as claimed in any of the preceding claims.
30. A transgenic non-human mammal which does not express a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein as claimed in claim 8, 9, 10, 11, 12, or 13, respectively, resulting in a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein associated pathology, respectively.
- 20
31. A transgenic animal assay system which provides a model system for testing for an agent that reduces or inhibits a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein associated pathology comprising
- 25
- (a) administering the agent to a transgenic non-human animal as claimed in claim 26; and
- (b) determining whether said agent reduces or inhibits a KLK-L1, KLK-L2, KLK-L3, KLK-L4, KLK-L5, or KLK-L6 protein associated pathology in the transgenic non-human animal relative to a transgenic non-human animal of step (a) which has not been administered the agent.
- 30

FIGURE 1






FIGURE 2



3/51


FIGURE 3



**FIGURE 4**


TGACCCGCTG TACCACCCCA GCATGTTCTG CGCCGGCGGA GGGCAAGACC  
AGAAGGACTC CTGCAACGGT GACTCTGGGG GGCCCCTGAT CTGCAACGGG  
TACTTGCAGG GCCTTGTGTC TTTCGGAAAA GCCCCGTGTG GCCAAGTTGG  
CGTGCCAGGT GCCTACACCA ACCTCTGCAA ATTCACTGAG TGGATAGAGA  
AAACCGTCCA GGCCAGTTAA CTCTGGGGAC TGGGAACCCA TGAAATTGAC  
CCCCAAATAC ATCCTGCGGA AGGAATTC

# FIGURE 5



6/51


FIGURE 6



## FIGURE 7

(ATG)GCTACAGCAAGACCCCCCTGGATGTGGGTGCTCTGTGCTCTGATCACAGCCT  
 M A T A R P P W M W V L C A L I T A  
 TGCTTCTGGGGGTCACAG[gt]aaccaga ----- intron 1----- tccc[ag]  
 L L L G V T  
 AGCATGTTCTCGCCAACAATGATGTTTCCTGTGACCACCCCTCTAACACCGTGCC  
 E H V L A N N D V S C D H P S N T V P  
 TCTGGGAGCAACCAGGACCTGGGAGCTGGGGCCGGGAAGACGCCCGGTCCGAT  
 S G S N Q D L G A G A G E D A R S D  
 GACAGCAGCAGCCGCATCATCAATGGATCCGACTGCGATATGCACACCCAGCCGT  
 D S S S R I I N G S D C D M H T Q P  
 GGCAGGCCGCGCTGTTGCTAAGGCCCAACCAGCTCTACTGCGGGGCGGTGTTGGT  
 W Q A A L L R P N Q L Y C G A V L V  
 GCATCCACAGTGGCTGCTCACGGCCGCCCACTGCAGGAAGAA[gt]agtgga-----  
 H P Q W L L T A A [H] C R K K  
 ----- intron 2----- tcttctc[ag]AGTTTTTCAGAGTCCGTCT  
 V F R V R L  
 CGGCCACTACTCCCTGTCAACAGTTTATGAATCTGGGCAGCAGATGTTCCAGGGG  
 G H Y S L S P V Y E S G Q Q M F Q G  
 GTCAAATCCATCCCCACCCCTGGCTACTCCACCCCTGGCCACTCTAACGACCTCAT  
 V K S I P H P G Y S H P G H S N [D] L M  
 GCTCATCAAAGTGAACAGAAGAATTTCGTCCTAAAGATGTCAGACCCATCAAC  
 L I K L N R R I R P T K D V R P I N  
 GTCTCCTCTCATTGTCCCTCTGCTGGGACAAAGTGCTTGCTGCTGGCTGGGGGAC  
 V S S H C P S A G T K C L V S G W G T  
 AACCAAGAGCCCCCAAG[gt]agtgagagag[gt]----- intron 3----- tgac[ag]  
 T K S P Q  
 TGCATTCCCTAAGGTCTCCAGTGCTTGAATATCAGCGTGCTAAGTCAGAAAAG  
 V H F P K V L Q C L N I S V L S Q K R  
 GTGCGAGGATGCTTACCCGAGACAGATAGATGACACCATGTTCTGCGCCGGTGAC  
 C E D A Y P R Q I D D T M F C A G D  
 AAAGCAGGTAGAGACTCCTGCCAG[gt]agagacacc----- intron 4-----  
 ag  
 K A G R D S C Q  
 GGTGATTCTGGGGGCGCTGTGGTCTGCAATGGCTCCCTGCAGGGACTCGTGTCCCT  
 G D [S] G G P V V C N G S L Q G L V S  
 GGGGAGATTACCCTTGTGCCCCGCCCAACAGACCGGTGTCTACACGAACCTCTG  
 W G D Y P C A R P N R P G V Y T N L C  
 CAAGTTCACCAAGTGGATCCAGGAAACCATCCAGGCCAACTCCTGAGTCATCC  
 CA  
 K F T K W I Q E T I Q A N S  
 GGACTCAGCACACCGGCATCCCCACCTGCTGCAGGGACAGCCCTGACACTCCTTT  
 CAGACCCTCATTCCTTCCCAGAGATGTTGAGAATGTTTCATCTCTCCAGCCCCCTGAC  
 CCCATGTCTCCTGGACTCAGGGTCTGCTTCCCCACATTGGGGCTGACCGTGTCTCT  
 CTAGTTGAACCTGGGAACAATTTCCAAAACCTGTCCAGGGCGGGGGTTGCGTCTC  
 AATCTCCCTGGGGCACTTTCATCCTCAAGCTCAGGGCCCATCCCTTCTCTGCAGCT  
 CTGACCCAAATTTAGTCCCAGAAATAAACTGAGAAG

FIGURE 8




## FIGURE 9

protease	MATAGNPWGWLFG----	YLILGVAGSLVSG-----	26
EMSP	MATAGNPWGWLFG----	YLILGVAGSLVSG-----	26
KLK-L2	MATARPPWMVLCALITALLGVTEHVLANNVSCDHPSNTVPSGSNQDLGAGAGEDARS		60
zyme	-----MKKLM-----	VVLSLIAAAWA-----	16
neuropsin	-MGRPRPRAAKTW----	MFLLLLGAWAGH-----	S 26
TLSP	-----MRILQ-----	LILLALATGLVG-----	17
PSA	-----MWVPVF-----	LTLSTWIGAAPL-----	20
KLK2	-----MWDLVLS-----	IALSVGCTGAVPL-----	20
KLK1	-----MWFLVLC-----	LALSLGGTGAAPP-----	20
trypsinogen	-----MNPLLI-----	LTFVAAALAAPFD-----	19
+			
protease	--SCSQIINGEDCSPHSQPWQAALVM--	ENELFCSGVLVHPQVWLSAAHCFQNSYITIGLGL	83
EMSP	--SCSQIINGEDCSPHSQPWQAALVM--	ENELFCSGVLVHPQVWLSAAHCFQNSYITIGLGL	83
KLK-L2	DDSSSRIINGSDCDMHTQWPQAALLLRPNQLYCGAVLVHPQVWLTAAHCKKPVFRVRLGH		120
zyme	--EEQNKLVHGGPCDKTSHPYQAALYT--	SGHLLCGGVLIHPLVWLTAAHCKKPNLQVFLGK	74
neuropsin	RAQEDKVLGGHECQPHSQPWQAALFE--	GOQLCGGVLVGGVWLTAAHCKKPKYIVHLGQ	85
TLSP	--GETRIKGFCEKPHSQPWQAALFQ--	KTRLLCGATLIAPVWLTAAHCKKPKYIVHLGQ	74
PSA	--ILSRIVGGWECEKHSQPWQVLVAS--	RGRAVCGGVLVHPQVWLTAAHCKKPNLQVFLGK	77
KLK2	--IQSRIVGGWECEKHSQPWQVAVYS--	HGWAHCGGVLVHPQVWLTAAHCKKPNLQVFLGK	77
KLK1	--IQSRIVGGWECEKHSQPWQAALYH--	FSTFQCGGILVHRQVWLTAAHCKKPNLQVFLGK	77
trypsinogen	--DDDKIVGGYNCEENSVPYQVLSNS--	GYHFCGGSLINEQVWLSAGHCYKSRIQVRLGE	75
+			
protease	HSLEADQEPGSMVEASLSVRHPEYN----	RP-----LLANDLMLIKLDESVS--ESDT	131
EMSP	HSLEADQEPGSMVEASLSVRHPEYN----	RP-----LLANDLMLIKLDESVS--ESDT	131
KLK-L2	YSLSPVYESGQMFQGVKSIHPHGYG----	HP-----GHSNDLMLIKLNRIR--PTKD	168
zyme	HNLRLQ--RESSQEQSSVVRAVTHPDY----	DAA-----SHDQDIMLIRLARPAK--LSEL	121
neuropsin	HSLQN--KDGPEQEIPVQSIHPHCYN--SSDVE----	DHNHDLMLIKLQRDQAS--LGSK	135
TLSP	HNLQK--EEGCEQTRTATESFPHPGFNNSLPNK-----	DHRNDIMLVKMASPVS--ITWA	125
PSA	HSLFH--PEDTGQVFQVSHSFPHPLYNMSLLKHQSLRPEDSSHDLMMLIRLSEPAE--LTDV		135
KLK2	HNLFE--PEDTGQVFPVSHSFPHPLYNMSLLKHQSLRPEDSSHDLMMLIRLSEPAE--LTDV		135
KLK1	HNLFD--DENTAQFVHVSESFPHPGFNMSLLENHTRQADEYSHDLMMLIRLSEPAE--LTDV		136
trypsinogen	HNIEV--LEGNEQFINAAKIRHPQYDRKTLNN-----	DIMLIRLSSRAV--INAR	122
+			
protease	IRISIASQCPTAGNSCLVSGWGLLANG--	RMPTVLQCVNVSVVSEEVCSKLYDPLYHPS	189
EMSP	IRISIASQCPTAGNSCLVSGWGLLANG--	RMPTVLQCVNVSVVSEEVCSKLYDPLYHPS	189
KLK-L2	VRPINVSSHCPSAGTKCLVSGWGTTSKSPQVHFPKVLQCLNISVLQKRCEDAYPRQIDDT		228
zyme	IQPLPLERDCSANTTSCHILGWGKTADG--	DFPDTIQAYIHLVSREECEHAYPGQITQ	179
neuropsin	VKPIISLADHCTQPGQKCTVSGWGTVTSPRENFPTLNCAEVKIFPQKKCEDAYPGQITDG		195
TLSP	VRPLTLSSRCVTAGTSCLSISGWGSTSSPQLRLPHTLRANITIEHQKCEAYPGNITDT		185
PSA	VKVMDLPTQEPALGTTTCYASGWGSIEPEEFLTPKKLQCVDLHVISNDVCAQVHPQKVTKF		195
KLK2	VKVLGLPTQEPALGTTTCYASGWGSIEPEEFLTPKKLQCVDLHVISNDVCAQVHPQKVTKF		195
KLK1	VKVVELPTEEPVGSTCLASGWGSIEPENFSFPDDLQCVDLKILPENDECKKAHVQKVTFD		196
trypsinogen	VSTISLPTAPPATGKCLISGWGNTASSGADYPDELQCLDAPVLSQAKCEASYPGKITSN		182
protease	MFCAGGGHDQKDCSCNGDSGGFLICNGYLQGLVSFGKAPCGQVGPVGYTNLCKFTIEWIEK		249
EMSP	MFCAGGGHDQKDCSCNGDSGGFLICNGYLQGLVSFGKAPCGQVGPVGYTNLCKFTIEWIEK		249
KLK-L2	MFCAG--DKAGRDSCQGDSDGGFLVVCNGSLQGLVSWGDYPCARPNRPVGYTNLCKFTKWIQE		287
zyme	MLCAGDEKYKDCSCQGDSDGGFLVCGDHLRGLVSWGNIPCGSKEKPGVYTNVCRYTNWIKK		239
neuropsin	MVCAGSSK--GADTCQGDSDGGFLVCDGALQGITSWGSDPCGRSDKPGVYTNICRYLDWIKK		254
TLSP	MVCASVQEGGKDCSCQGDSDGGFLVCNQLQGIISWGQDPCAITRKPGVYTKVCKYVDWIKK		245
PSA	MLCAGRWGKSTCSGDSDGGFLVCNQLQGITSWGSEPCALPERPSLYTKVHYRKWIKD		255
KLK2	MLCAGRWGKSTCSGDSDGGFLVCNQLQGITSWGSEPCALPERPSLYTKVHYRKWIKD		255
KLK1	MLCVGHLEGGKDTCTVGDSDGGFLMCDGVLVQGVTSWGVPCGTPNKPVSVAVRVLSYVKWIED		256
trypsinogen	MFCVGFLEGGKDCSCQGDSDGGFLVVCNGQLQGVVSWG--DGCAQKNKPGVYTKVYNYVKWIKN		241
O     +			


10/51

FIGURE 10

(A)




(B)




11/51

FIGURE 11



12/51

**FIGURE 12**

# FIGURE 13





FIGURE 14



## FIGURE 15

CACTGGACGGGTGCACGTTTCAGGATCCAGGTGCCAGGGGTC(ATG)AAG CTG GGA  
 CTC  
 M K L G L  
 CTC TGT GCT CTG CTC TCT CTG CTG GCA Ggtga... intron 1 ..ccag GG CAT  
 GGC  
 L C A L L S L L A G H  
 TGG GCA GAC ACC CGT GCC ATC GGG GCC GAG GAA TGT CGC CCC AAC TCC  
 CAG  
 W A D T R A I G A E E C R C N S  
 Q  
 CCT TGG CAG GCC GGC CTC TTC CAC CTT ACT CGG CTC TTC TGT GGG GCG  
 ACC  
 P W Q A G L F H L T R L F C G A T  
 CTC ATC AGT GAC CGC TGG CTG CTC ACA GCT GCC CAC TGC CGC AAG  
 CCgtga.....  
 L I S D R W L L T A A H C R K P  
 intron 2 .....gcag G TAT CTG TGG GTC CGC CTT GGA GAG CAC CAC CTC TGG AAA  
 Y L W V R L G E H H L W K  
 TGG GAG GGT CCG GAG CAG CTG TTC CGG GTT ACG GAC TTC TTC CCC CAC  
 CCT  
 W E G P E Q L F R V T D F F P H P  
 GGC TTC AAC AAG GAC CTC AGC GCC AAT GAC CAC AAT GAT GAC ATC ATG  
 CTG  
 G F N K D L S A N D H N D D I M L  
 ATC CGC CTG CCC AGG CAG GCA CGT CTG AGT CCT GCT GTG CAG CCC CTC  
 AAC  
 I R L P R Q A R L S P A V Q P L N  
 CTC AGC CAG ACC TGT GTC TCC CCA GGC ATG CAG TGT CTC ATC TCA GGC  
 TGG  
 L S Q T C V S P G M Q C L I S G W  
 GGG GCC GTG TCC AGC CCC AAG Ggtat..... intron ..acag CG CTG TTT CCA  
 GTC  
 G A V S S P K A L F P  
 V  
 ACA CTG CAG TGT GCC AAC ATC AGC ATC CTG GAG AAC AAA CTC TGT CAC  
 TGG  
 T L Q C A N I S I L E N K L C H W  
 GCA TAC CCT GGA CAC ATC TCG GAC AGC ATG CTC TGT GCG GGC CTG TGG  
 GAG  
 A Y P G H I S D S M L C A G L W E  
 GGG GGC CGA GGT TCC TGC CAGgtga..... intron ..acag GGT GAC TCT GGG  
 GGC  
 G G R G S C Q G D S G  
 G  
 CCC CTG GTT TGC AAT GGA ACC TTG GCA GGC GTG GTG TCT GGG GGT GCT  
 GAG  
 P L V C N G T L A G V V S G G A E

## FIGURE 15 (CONT'D)


CCC TGC TCC AGA CCC CGG CGC CCC GCA GTC TAC ACC AGC GTA TGC CAC  
TAC

P C S R P R R P A V Y T S V C H Y

CTT GAC TGG ATC CAA GAA ATC ATG GAG AAC (TGA)  
L D W I Q E I M E N

GCCCCGCGCCACGGGGGCACCTTGGAAGACCAAGAGAGGCCGAAGGGCACGGGGTA  
GGGGGTTCTCGTAGGGTCCCAGCCTCAATGGTTCCCGCCCTGGACCTCCAGCTGCCCTG  
ACTCCCCTCTGGACACTAAGACTCCGCCCCCTGAGGCTCCGCCCCCTCACGGGTCAAGCA  
AGACACAGTCGCGCCCCCTCGGAACGGAGCAGGGACACGCCCTTCAGAGCCGTCTCTAT  
GACGTCACCGACAGCCATCACCTCCTTCTTGGAACAGCACAGCCTGTGGCTCCGCCCCA  
AGGAACCACTTACACAAAATAGCTCCGCCCCCTCGGAACCTTTGCCAGTGGGACTTCCCC  
TCGGGACTCCACCCCTTGTGGCCCCGCTCCTTCACCAGAGATCTCGCCCCCTCGTGATGT  
CAGGGGCGCAGTAGCTCCGCCCCAGTGGAGCTCGGGCGGTGTAGAGCTCAGCCCTTGTG  
GCCCCGTCTGGGCGTGTGCTGGGTTTGAATCCTGGCGGAGACCTGGGGGGAATTGAG  
GGAGGGTCTGGATACCTTTAGAGCCAATGCAACGGATGATTTTTTCAGTAAACGGGGAAA  
CCTCA

FIGURE 16




## FIGURE 17

PSA	-----MWVPVFLTSLVWTWIGAAPLI-LSRIVGGWECEKHSQWPQVLVASRGRAVC
KLK2	-----MWDLVLSIALSVGCTGAVPLI-QSRIVGGWECEKHSQWPQVAVYSHGWAHC
KLK1	-----MWFLVLCALSLGGTGAAPPI-QSRIVGGWECEQHSQWPQAALYHFSTFQC
trypsinogen	-----MNPLLILTFVAAALAAFFDD-DDKIVGGYNCEENSVPYQVSLNS-GYHFC
KLK-L3	-----MKLGLLCCALLSLLAGHGWA--DTRAIGAEECRPNSQPWQAGLFHLTRLFC
TLSP	-----MRI-LQLILLALATGLVGG--ETRIIKGFECKPHSQWPQAALFEKTRLLC
neuropsin	-MGRPRPRAAKTWMFLLLGGAWAGHSRAQ-EDKVLGGHECQPHSQWPQAALFQGGQLLC
zyme	-----MKK--LMVVLSLIAAAWAE--QNKLVHGGPCDKTSHPYQAALYTSGLLCC
HSCCE	---MARSLLLPLQILLLSLALETAGEEAQG--DKIIDGAPCARGSHPWQVALLSGNQLHC
protease	---MA-TAGNPWGWFGLYLILGVAGSLVSGSCSIINGEDCSPHSQWPQAALVMENELFC
♦	
PSA	GGVLVHPQVLTAAHCIRNKSVILLGRHSLFHPEDT-GQVFQVSHSFPHPLYDMSLLKNR
KLK2	GGVLVHPQVLTAAHCLKNSQVWLGRHNLFEEDT-GQVFPVSHSFPHPLYNMSLLKHQ
KLK1	GGILVHRQVLTAAHCISDNYQLWLGRHNLFDENT-AQFVHVSESFPHPGFNMSLLENH
trypsinogen	GGSLINEQVVSAGHCYKSRIOVRLGEHNIIEVLEGN-EQFINAAKIIIRHPQYDRKTLNN-
KLK-L3	GATLISDRALLTAAHCRKPYLVVRLGEHHLWKWEGP-EQLFRVTDFFPHPGFNKDLNAN-
TLSP	GATLIAPRLLTAAHCLKPRYIVHLGQHNLQKEEGC-EQTRTATESFPHPGFNNSLPNK-
neuropsin	GGVLVGGNVLTAACHCKPKYTVRLGDHSLQNKDGP-EQEIPVVQSIHPPCYNSSD-VE-
zyme	GGVLIHPLVLTAAHCKKPNLQVFLGKHNLQRESS-QEQSSVRAVIHPDYDAAS----
HSCCE	GGVLVNERVLTAAHCKMNEYTVHLGSDTLGDRR---AQRIKASKSFRHPGYSTQT----
protease	SGVLVHPQVLTAAHCFQNSYITIGLGLHSLQEPGSGMVEASLSVRHPEYNRPLLAN-
♦ ♦ ♦ ♦ ♦	
PSA	FLRPGDDSSHDMLRLSEPAAE-LTDAVKVMDLPTQEPALGTTTCYASGWGSIPEEFLTP
KLK2	SLRPDESSHDMLRLSEPAK-ITDVVKVLGLPTQEPALGTTTCYASGWGSIPEEFLRP
KLK1	TRQADEYSHDMLRLTEPADTITDAVKVVELPTEEPVSTCLASGWGSIPEENFSFP
trypsinogen	-----DIMLIKLSRAV-INARVSTISLPTAPPATGKCLISGWGNTASSGADYP
KLK-L3	-----DHNDIMLIRLPRQAR-LSPAVQPLNLSQTCVSPGMQCLISGWGAVSSPKALFP
TLSP	-----DHRNDIMLVKMASPVS-ITWAVRPLTLSSRCVTAGTSCLISGWGSTSSPQLRLP
neuropsin	-----DHNHDLMLQLRDQAS-LGSKVKPISLADHCTQPGQKCTVSGWGTVTSPRENFP
zyme	-----HDQDIMLRLRPAK-LSELIQPLPLERDCSANTTSCHILGWGKTADG--DFP
HSCCE	-----HVNDLMLVKNLSQAR-LSSMVKKVRLPSRCEPPGTCTCTVSGWGTTSPTDVTFP
protease	-----DLMLIKLDESVS-ESDTIRSIASQCPTAGNSCLVSGWGLLANG--RMP
♦ ♦ ♦ ♦ ♦	
PSA	KKLQCVDLHVISNDVCAQVHPQKVTKFMLCAGRWTGGKSTCSGDSGGGLVCNGVLQGITS
KLK2	RSIQCVSLHLLSNDMCARAYSEKVFMLCAGLWTGKDTCCGDSGGGLVCNGVLQGITS
KLK1	DDLQCVDLKILPNDECKKAHVQKVTDFMLCVGHLEGGKDTCCVGGDSGGGLMCDGVLQGVTS
trypsinogen	DELQCLDAPVLSQAKCEASYPGKITSNMFCVGFLEGGKDSQCGDSGGFPVVCNGQLQGVVS
KLK-L3	VTLQCANISILENKLCHWAYPGHISDSMLCAGLWEGGRGSCQCGDSGGGLVCNGTLAGVVS
TLSP	HTLRCANITIIHQKCEANAYPGNITDTMVCASVQEGGKDSQCGDSGGGLVCNQSLQGIIS
neuropsin	DTLNCAEVKIFPQKKCEDAYPGQITDGMVCAGSSKG-ADTCQCGDSGGGLVCDGALQGITS
zyme	DTIQCAYIHLVSREECEHAYPGQITQNMLCAGDEKYGKDSQCGDSGGGLVCGDHLRGLVS
HSCCE	SDLMCVDVKLISPQDCTKVYKDLLENSMLCAGIPDSKKNACNGDSGGGLVCRGTQGLVVS
protease	TVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGGHDQKDSNCNDSGGGLICNGYLQGLVS
♦ ♦ ♦ ♦ ♦	
PSA	WGSEPCALPERPSLYTKVVHYRKWKIDTIVANP
KLK2	WGPEPCALPEKPAVYTKVVHYRKWKIDTIAANP
KLK1	WGYVPCGTPNKPSAVRVL SYVKWIEDTIAENS
trypsinogen	WG-DGCAQKNKPGVYTKVYNYVKWIKNTIAANS
KLK-L3	GGAEPCSRPRRPAVYTSVCHYLDWIEIMEN--
TLSP	WGQDPCAITRKPGVYTKVCKYVDWIQETMKN-
neuropsin	WGSDDPCGRSDKPGVYTNICRYLDWIKKIIGSKG
zyme	WGNIPCGSKEKPGVYTNVCRYTNWIQKTIQAK-
HSCCE	WGTFCGQPNDPGVYTVQCKFTKWINDTMKKHR
protease	FGKAPCGQVGPVYTNLCKFTIEWIEKTVQAS-


19/51

## FIGURE 18



20/51

FIGURE 19




**FIGURE 20**

FIGURE 21







FIGURE 22




24/51

FIGURE 23



25/51

FIGURE 24



SUBSTITUTE SHEET (RULE 26)

**FIGURE 25**


TCAGGCCCCGCCCCGCCCTGCCCTCCCCTCCCGATCCCGGAGCC (ATG) TGG CCC CTG GCC  
M W P L A  
CTA GTG ATC GCC TCC CTG ACC TTG GCC TTG TCA GGA G...gtaaga.... intron 1 ..... ttaccag  
L V I A S L T L A L S G  
GT GTC TCC CAG GAG TCT TCC AAG GTT CTC AAC ACC AAT GGG ACC AGT GGG TTT  
G V S Q E S S K V L N T N G T S G F  
CTC CCA GGT GGC TAC ACC TGC TTC CCC CAC TCT CAG CCC TGG CAG GCT GCC  
L P G G Y T C F P H S Q P W Q A A  
CTA CTA GTG CAA GGG CGG CTA CTC TGT GGG GGA GTC CTG GTC CAC CCC AAA  
L L V Q G R L L C G G V L V H P K  
TGG GTC CTC ACT GCC GCA CAC TGT CTA AAG GA gtatgt ..... intron 2..... cacag G GGG  
W V L T A A H C L K E G  
CTC AAA GTT TAC CTA GGC AAG CAC GCC CTA GGG CGT GTG GAA GCT GGT GAG  
L K V Y L G K H A L G R V E A G E  
CAG GTG AGG GAA GTT GTC CAC TCT ATC CCC CAC CCT GAA TAC CGG AGA AGC  
Q V R E V V H S I P H P E Y R R S  
CCC ACC CAC CTG AAC CAC GAC CAT GAC ATC ATG CTT CTG GAG CTG CAG TCC  
P T H L N H D H D I M L L E L Q S  
CCG GTC CAG CTC ACA GGC TAC ATC CAA ACC CTG CCC CTT TCC CAC AAC AAC CGC  
P V Q L T G Y I Q T L P L S H N N R  
CTA ACC CCT GGC ACC ACC TGT CGG GTG TCT GGC TGG GGC ACC ACC ACC AGC  
L T P G T T C R V S G W G T T T S  
CCC CAG G gtatgcac... intron 3..... tcgcc ag TG AAT TAC CCC AAA ACT CTA CAA TGT GCC  
P Q V N Y P K T L Q C A  
AAC ATC CAA CTT CGC TCA GAT GAG GAG TGT CGT CAA GTC TAC CCA GGA AAG  
N I Q L R S D E E C R Q V Y P G K  
ATC ACT GAC AAC ATG TTG TGT GCC GGC ACA AAA GAG GGT GGC AAA GAC TCC  
I T D N M L C A G T K E G G K D S  
TGT GAG gtatgca... intron 4..... aactcag GGT GAC TCT GGG GGC CCC CTG GTC TGT AAC  
C E G D S G G P L V C N  
AGA ACA CTG TAT GGC ATC GTC TCC TGG GGA GAC TTC CCA TGT GGG CAA CCT  
R T L Y G I V S W G D F P C G Q P  
GAC CGG CCT GGT GTC TAC ACC CGT GTC TCA AGA TAC GTC CTG TGG ATC CGT  
D R P G V Y T R V S R Y V L W I R

## FIGURE 25 (CONT'D)

GAA ACA ATC CGA AAA TAT GAA ACC CAG CAG CAA AAA TGG TTG AAG GGC CCA  
E T I R K Y E T Q Q Q K W L K G P

CAA (TAA) AAGTTGAGAAATGTACCGGCTTCCATCCTGTCACCATGACTTCCTCAC  
Q

ATGGTCTGCTTAGCCCTTCTCTGCTCCTTATTCCCAGTGTTCCATTTGAACCAGTGATCCATGTC  
CTGAAAAATGCTCAATCTCAGCTAACATTCCATGTTTCAGAAGCATTGAGGCACTGCCAGGCT  
TGCAGTCTCCCAGATGTTGCATCCCTGAAACATCTCAACAACCTGAATGTCCCAACCCAGACA  
ATGGCCCAGGTCTCTCAACTTCATCAGTGTGGCTTCTATGAGCCCAGATCACCACCTGAACGT  
TCTGTCTGTGGCACATTCTTAAATATTTCCATCAGCCCATCTCAACAATATATGTCCTATAAAT  
GGACCATCCTTGACA


**FIGURE 26**

1	KLK-L1/protease	15 16	30 31	45 46	60 61	75 76	90
2	EMSP	---MATAGN	PWGFLG---	YLIL	GVAG---	SLVSG---	INGEDCSPHSQPW
3	KLK-L2	---MATAGP	PWGFLG---	YLIL	GVAG---	SLVSG---	INGEDCSPHSQPW
4	PSA	---MATARP	PMMVLCALITALL	GYTERH	LVANDVSCD	HPSNTVPSGSNQDLG	AGAGEDARSDDSSR
5	KLK2	---MATVPVF---	LTLSV	TWIG	---	AAPLI---	IVGWCEKHSQPW
6	KLK1	---MMDLVLS---	IALLS	GCTG	---	AVPLI---	IVGWCEKHSQPW
7	trypsinogen	---MFLVLTC---	LALLS	GCTG	---	AAPPI---	IVGWCEKHSQPW
8	zyme/protease M	---MKPLLI---	LTFFA	AALA	---	APFDD---	IVGVNCEENSVPYQ
9	KLK-L4	---MKLMV---	VLSL	IAAA	---	WAEBQNK---	LHVGPCDKTSHIPYQ
10	TLSP	---MW	PLALVTA	---SLTL	ALSG---	GVGSSESKV---	LPGGYTCFPHSQPW
11	neuropsin	---MR	ILQILLALATGLVG	GETR	---	I---	IKGFECKPHSQPW
12	NES1	---MGRPRPRA	ATAKMFELLGGLGAWA	GHSR	---	AQEDK---	VLGHEQCQPHSQPW
		MRAPHLHL	SAASGAR	AKTWILLPMLLAQLWA	AEAA---	LIPQN---	EAYGAPCARGSQPW
91	1	105 106	120 121	135 136	150 151	165 166	180
1	protease	AALVN-ENELFC	SGV LVHPQ---	NSYITGILGLH	SLEAD	QEPGSMVEASLSVR	HPEVNRPLAN---
2	EMSP	AALVN-ENELFC	SGV LVHPQ---	NSYITGILGLH	SLEAD	QEPGSMVEASLSVR	HPEVNRPLAN---
3	KLK-L2	AALLRN-ENELFC	SGV LVHPQ---	NSYITGILGLH	SLEAD	QEPGSMVEASLSVR	HPEVNRPLAN---
4	PSA	VLVAS-RGRV	CGGV LVHPQ---	NKSVIRLGRH	SLSPV	YEDGQMFQGVKSIP	HPLYDMSLLKNRFLR
5	KLK2	VAVYS-HGMV	CGGV LVHPQ---	NKSVIRLGRH	SLSPV	YEDGQMFQGVKSIP	HPLYDMSLLKNRFLR
6	KLK1	AALVH-FSTFC	CGG LVHPQ---	IS	DNQMLGRHNLFPD	EDTGRVPSVHSFP	HPLVNMSLLKHQSLR
7	trypsinogen	VSLNS--GYHPC	CGGS LINEQ---	VS	GRYK	SRTQVRLEHNI	EVL
8	zyme	AALYT-SGHL	CGGV LHPL---	QVGLK	PNLQVTLGK	HNLR	---
9	KLK-L4	AALLV-QGRLL	CGGV LVHPQ---	LNK	EGKVVYLG	RHAGLGRV	---
10	TLSP	AALFE-KTRULL	CGAT LIAPR---	PL	PKYIVH	LGHOHLOKE	---
11	neuropsin	AALFQ-QOQL	CGGV LVGGN---	QVGLK	PKYTVR	GLDHSLOK	---
12	NES1	VSLFN-GLSP	HCAG LVDSQ---	LNK	KPLVAR	VGDHLL	---
181	1	195 196	210 211	225 226	240 241	255 256	270
1	protease	ESVS-ESDTR	IRTSI ASOCTP	AG-NSCLVS	GWGLLANG--	RMPTV	LQCVNVSVVSEVCS
2	EMSP	ESVS-ESDTR	IRTSI ASOCTP	AG-NSCLVS	GWGLLANG--	RMPTV	LQCVNVSVVSEVCS
3	KLK-L2	RRIR-PTKDV	RTINV SSHCP	SPAG-TKCLVS	GWGTSIEP	EEFLP	PKV
4	PSA	EPAE-LTDAV	KVMDL PTOEP	PALG-TTCYAS	GWGTSIEP	EEFLP	PKV
5	KLK2	EPAK-ITDV	KVYGL PTOEP	PALG-TTCYAS	GWGTSIEP	EEFLP	PKV
6	KLK1	EPADITDAV	KVVEL PTEBE	VEG-STCLAS	GWGTSIEP	EEFLP	PKV
7	trypsinogen	SRAV-INAR	VTSI PTAPP	ATG-TKCLVS	GWGTSIEP	EEFLP	PKV
8	zyme	RPAK-LESEL	IOPLP ERDCS	ANT-TSCHL	GWGKTADG--	DEPDT	IQCAIYHLV
9	KLK-L4	SPVQ-LTGY	IQIOTPL SHNRN	TPGTCRVS	GWGTTSPQ	VNYPKPT	LQCANITLR
10	TLSP	SPVS-ITWAV	PLTL SCRVCT	AG-TSCLS	GWGTTSSP	QLRLP	PKT
11	neuropsin	DQAS-LGSK	VXPISL ADHCT	QPG-QKTVS	GWGTTS	SPRENF	PDT
12	NES1	RPVVP-GRV	RVALQL PYRCA	QPG-DQCQVA	GWGTTA	ARRVYNK	G


FIGURE 27 (CONT'D)

1 prostate	271	285	286	300	301	315	316	330	331
2 EMSP	271	285	286	300	301	315	316	330	331
3 KLK-L2	271	285	286	300	301	315	316	330	331
4 PSA	271	285	286	300	301	315	316	330	331
5 KLK2	271	285	286	300	301	315	316	330	331
6 KLK1	271	285	286	300	301	315	316	330	331
7 trypsinogen	271	285	286	300	301	315	316	330	331
8 zyme	271	285	286	300	301	315	316	330	331
9 KLK-L4	271	285	286	300	301	315	316	330	331
10 TLSP	271	285	286	300	301	315	316	330	331
11 neuropsin	271	285	286	300	301	315	316	330	331
12 NES1	271	285	286	300	301	315	316	330	331

FIGURE 28



## FIGURE 29




33/51  
**FIGURE 30**

FIGURE 31






FIGURE 32



## FIGURE 33

GCAGGTAGGTGGACGGAGAGATAGCAGCGACGAGGACAGGCCAAACAGTGACAGCCACG  
 TAGAGGATCTGGCAGACAAAGAGACAAGGTGAGAAGGAG gtagg.....Intron 1.....  
 .....tgacactccccag ACTTTGGAAGTGACCCACC (ATG)  
 M  
 GGG CTC AGC ATC TTT TTG CTC CTG TGT GTT CTT G gtgagttctcccg  
 G L S I F L L L C V L  
 gagcagggagagggca..... Intron 2 .....cctgtctgtctccag GG CTC  
 G L  
 AGC CAG GCA GCC ACA CCG AAG ATT TTC AAT GGC ACT GAG TGT GGG  
 S Q A A T P K I F N G T E C G  
 CGT AAC TCA CAG CCG TGG CAG GTG GGG CTG TTT GAG GGC ACC AGC  
 R N S Q P W Q V G L F E G T S  
 CTG CGC TGC GGG GGT GTC CTT ATT GAC CAC AGG TGG GTC CTC ACA  
 L R C G G V L I D H R W V L T  
 GCG GCT CAC TGC AGC GGC AG gtaagtcccttcc.....intron3.....  
 A A (H) C S G S  
 .ccgtcgccaccggcag C AGG TAC TGG GTG CGC CTG GGG GAA CAC AGC  
 R Y W V R L G E H S  
 CTC AGC CAG CTC GAC TGG ACC GAG CAG ATC CGG CAC AGC GGC TTC  
 L S Q L D W T E Q I R H S G F  
 TCT GTG ACC CAT CCC GGC TAC CTG GGA GCC TCG ACG AGC CAC GAG  
 S V T H P G Y L G A S T S H E  
 CAC GAC CTC CGG CTG CTG CGG CTG CGC CTG CCC GTC CGC GTA ACC  
 H (D) L R L L R L R L P V R V T  
 AGC AGC GTT CAA CCC CTG CCC CTG CCC AAT GAC TGT GCA ACC GCT  
 S S V Q P L P L P N D C A T A  
 GGC ACC GAG TGC CAC GTC TCA GGC TGG GGC ATC ACC AAC CAC CCA  
 G T E C H V S G W G I T N H P  
 CGG A gtaaggggcccagggccaggg.....intron 4 .....  
 R  
 .gaccctgcagcacgcatgttctctctccag AC CCA TTC CCG GAT CTG CTC  
 N P F P D L L  
 CAG TGC CTC AAC CTC TCC ATC GTC TCC CAT GCC ACC TGC CAT GGT  
 Q C L N L S I V S H A T C H G  
 GTG TAT CCC GGG AGA ATC ACG AGC AAC ATG GTG TGT GCA GGC GGC  
 V Y P G R I T S N M V C A G G  
 GTC CCG GGG CAG GAT GCC TGC CAG gtgagcc..... Intron 5 .....  
 V P G Q D A C Q  
 .aaaacagaaataagatgtctccctgttccagacagtacttctcttccctccag GGT  
 G  
 GAT TCT GGG GGC CCC CTG GTG TGT GGG GGA GTC CTT CAA GGT CTG  
 D (S) G G P L V C G G V L Q G L  
 GTG TCC TGG GGG TCT GTG GGG CCC TGT GGA CAA GAT GGC ATC CCT  
 V S W G S V G P C G Q D G I P  
 GGA GTC TAC ACC TAT ATT TGC AA(G TAT GTG GAC TGG ATC CCG ATG  
 G V Y T Y I C K Y V D W I R M  
 ATC ATG AGG AAC AAC (TGA) CCTGTTTCCTCCACCTCCACCCCCACCCCTTAACCT  
 I M R N N  
 GGGTACCCCTCTGGCCCTCAGAGCACCATAATCTCCTCCATCACTTCCCTAG) CTCCAC  
 TCTTGTGGCCTGGGAACCTTCTTGGAACTTTAACTCCTGCCAGCCCTTC (TAA) GACCCACG  
 AGCGGGGTGAGAGAAGTGTGCAATAGTCTGGAATAAATATAAATGAAGGAGGGGC

FIGURE 34





**FIGURE 35**



Figure 36 cont'd


1	PSA	271	285	286	300	301	261
2	hk2	SLYTKVWHYRKWKD	TIVANP	-----	-----	-----	261
3	hk1	AVYTKVWHYRKWKD	TIAANP	-----	-----	-----	262
4	protease	SVAVRVLSYVKWIED	TIAENS	-----	-----	-----	254
5	zyme	GVYTNLCCKFTENIEK	TVQAS	-----	-----	-----	244
6	TLSP	GVYTNVCRYTNNIOK	TIQAK	-----	-----	-----	250
7	KLK-L4	GVYTKVCKYVDWIOE	TMKNN	-----	-----	-----	277
8	NES1	GVYTRYSRYVLWIRE	TIRKYETQQQKWLKG	PQ	-----	-----	248
9	KLK-L5	AVYQJICKYMSWINK	VIRSN	-----	-----	-----	260
10	neuropsin	GVYTIICKYVDWIRM	IMRNN	-----	-----	-----	
		GVYTNICRYLDWIKK	IIGSKG	-----	-----	-----	

**FIGURE 37**




42/51

# FIGURE 38




43/51

**FIGURE 39**



44/51

# FIGURE 40



SUBSTITUTE SHEET (RULE 26)

45/51  
FIGURE 41

ATCGTGTAAT CACCGCCACA TCCAGTGCAA AGCTGATTCG TCACCACAGA GCAGCTCCCT  
 CCTGCCACCC CATCCCTGGG TCCCAAGAGA ACCCTTTCTT AAAAGAGGGA GTTCTTGACG  
 GGTGTGGTGG CTCATGCCTG TAATCCTTGC ACTTTGGGAG GCCAAGGAGG GTGGATCATT  
 TGAGGTCAGG AGTTTGAGAC CAGACTGGCC AACATGGTGA AACCTGTCT TACTAAAAA  
 TACAAAAA TGAGCGGGG ATGGTGGTGG GTGCCATAG CCCAGCTAC TCAGGAGGCT  
 GAGGCAGGAG AATCGCTTGA ACCCAGGAGG CAGAGGTGC AGTGAGCCGA GATTGAGCCA  
 CTGCACTCCA GCCGGGGCTA AAGAGTGAGA CTCTGTCTCA AAAAAA AAAAGAAAAA  
 AAAAAAGAA AAAAAATAA AATAAATAA TAAATAAAT AAATTTAAAA ATTTAAAAAT  
 AAAGAGGGGG TTCTTGTGTT GATGCCGAGC CTGAACCAAG GCAGAGGAGG CCGGGAAGGC  
 TTCCAAGGC CTTACAGCTCA AAGCAGGGAG GCCCATAGTT AAACAGAAAC AGTTCAGGAA  
 TCACAGAAAG GCACCTGGGG AGAGATGGGT GTGTGGCTCC AGATGCAGGT CCCAGACAG  
 TGCGTCCCCA GGTGTACAGA CAGACCCAGG CCAAGCTCCA GCTCAAAGAG CCAGCCTAGG  
 GGGGTGCCGA GGTGGAGGGA GGCTGAGTCA GGCTGAGGCC GGGGAACAGT TGGGGTAGCC  
 AAGGGAGGCA AGCAGCCTCC TGAGTCACCA CGTGGTCCAG GTACGGGGCT GCCCAGGCCC  
 AGAGACGGAC ACAAGCACTG GGAATTTAA GGGGCTAGGG GAGGGGCTGA GGAGGGTAGG  
 CCCTCCCCCA AATGAGGATG GAACCCCCC AACTCCAGAA CCCCCCTGCA GGCTGGCCAG  
 AATCCTTCCC CATCTCATTC ACTCTGTCTC TCCTGTCTCT TGCCGTCTCC TATTTTGAAT  
 TTCCAACCCC GTCTGTTAAG ACTGTCTCTC TGTCTCTGAA TCTCTGTCCC CTCTCTTTTC  
 TGGGTCTCTC TCCTCTCTCC TCTGGGTCTC TGTCCCCCTC TCTGGGTCTC TGTCACCTCTC  
 TCTTTCATC TCCAGCTCTC ACTTTGTCTC **TGCACCTAGC AGATCCCAAG CTGGGGAATG**  
**CCAGTTCTGG CACCAACCTT CCTGCTCCCT GCTGGGGCCT CTGCTCCCC ATCTCTCAGG**  
**AGTCGAAAGT GAGAAAGCAA GGTGGGCAGC TCTGCTCCAG GTCCAGGTAT CTCCGCCCCA**  
 CCTCTGCCC GTCCTCTATC CCACCCCTCC TCTCCATCTC TCCCTGGCGC TGCCATCTCT  
 CATCTAGGCC TCCGTCTCCT CTGTCAATTGT CCCCATCCCC TGTAGGTGCC CATCCTTCCC  
 GTCTCCCCCTC TGCCATCGGC CTGCCGTGCC CATCCTCTTT CTCCACCAT GTCCGTTCT  
 CTTCCACGTC TCATGCCCGC ACTGCCCTCA TCATCATCGC TGTGTGTTCTG TGTGTGTTTG  
 TGGTGAGTGC CGCATGGTGG GGGCGTCTCG GCCTCTCTCC TCTCTCTCCA CTGTTTTCTC  
 TTTCTGTGTG TCTGTTTTCA TTCTATCTCC ACCTTCTTCC CTCCGTCTTT TGCTTTTCTA  
 TCTCCACTTC TCCACACCCC TCTCTCCCTG CGTCTCTGTG TCTCCCTCTT CCTCTGTCTT  
 GTTTTTTTC CACCGTCTGC CTCTTCTGTT CCCTGTCACA TCCAACCTCC ACCGGTTTCT  
 CCAGCTCTCT CCTCAGTTCC TTCTCTCATG AGCACACCTG CCTCTGTGCT CGTATTCTCTG  
 GACTCCTCTC TCTCCACTGT CATATCTTCT CATTCATTTT CCCAGTCTCT CTCTGTCTCT  
 TGCTCTCCCC CTCTCTGTCA CTCTGTCTCT GTCTCTCTCT TTCTCTCTCT CTCTCTGTGT  
 CTCTCTGTCT GGCTCTCTCT CTGTCTCTCT CTCCATCTCT CTCTCTCTCT CCCCCCGTC  
 ACCCTGTCTC TGTCTCTCTC TGTCTGTGTG TCTCTCTGTC TTTCTCTCTC TCCATCTCTC  
 TCTGTCTCTC TCTCTCTCTC TCTCTCTCTC CCTCTCTCCC TCCTCCCGTG ACTCCCTCTC  
 TCAGTCCATC TCTTCTCTCC TCTCTCAGCC CCTTCGTGCC CTTTCTCTG ACACTCCCCA  
 CCCTGGTTTC CTGACTCCAC CACTAGATCC ACCACCTCCA GCAACTGGGA ACCCTCCCTT  
 GCCCACCTG CCCTGGGGTC CCTCCCAGG ATTCCTTCTA GATTATAGCA TCTTCCCTGG  
 GCGGTTCTC ATGAACAATT GTGGCTGCTT TTTTGCCAG ACAGGGGAGG GAGGGGATGG  
 GATCAGGGAG TCTTGAATG GGAAGTAGGC AATAAAAAA AAAAAATGTC AGAAGCAGGG  
 CGGCGGAGG TGGGGGAGG GCCAGCTGTC CTTACCAGGG ATAAAAGGCT TTGCCAGTGT  
 GACTAGGAAG AGAGACACCT CCCCTCCTTC CTTTCATCAAG **ACATCAAGGA GGGACCTGTG**  
**CCCTGCTCCA CATCTCCCA CCTGCCGCC GCAGAGCCTG CAGGCCCGC CCCCCTCGTC**  
**TCTGGTCCCT ACCTCTCTGC TGTGCTTCA TGTCCCTGAG GGTCTTGGGC TCTGGGTAAG**  
 TGCCCCCTGC TGTCTCTGCC TCTCAGCCCC CGGTTCTGTT GAAGGTTCTT TCTCTCTCAC  
 TTTTCTCTG CATTTGACAG **GACCTGGCCC TCAGCCCTA AAATGTTCTT CCTGCTGACA**  
**GCACTTCAAG TCCTGGCTAT AGGTAAGAGA** ACGGTGCGGT ATGACACAAG GGGGTCCCTT  
 GGAGACTCTG AGAAGAGATG GGGATGGGTC CTTGGGGCCC CTGGATGCTC ATGGTGACCT  
 CATAAGAAAG AGCAGGGAGT GGTGTTGGGG TCATGGTGGG GGAACGTGCT GGAGGCCTAA  
 ATTCCTAGTT GTGGAGGTGC TAGGGAATTG TGGGGCCGGG GAGAGAGGTG TTTATAAGGT  
 CTGGTGCAAA ATACATAAGG AATCTTAGGG AACTATTAGG TCCTGAGTGG GTCATAGCAG  
 AAAGATCACG GGGCTCTACC TGACTGTGTT AGGAAAGAAA CAATGTCAGA AAGATGTTTT  
 GTTGTACAGG GGAAGGTGGA GAAGGATGAT GGGATGGCGG GATCGTGGCA TGGGTGGCG  
 GGATCGTGGC ATGGGTGTGT GAGGTGGATG GGGCAAGTG TGGGGCAAGA GATGGCGGAT  
 CCTTGGGGTC CCACTGAGTG GGAACGTTGG GGAGGAGACA GGGAGTCTCT TGAATGTGTT  
 GGGGAAGGAC TCATTGGGGG GAAATGTGGC ATATTTCCAG AAGTGATCAC AGAAATTATG  
 GGAGCATAGA GCTAAGGGTC GTAGATGTAG CAAGGCCCTG GATAAGGTGG CCACGGCACA  
 AAATAAGAGA TGCTACGGAG GTGACTTGGG AGGTGAGTCA GAAAGCTCTC CGTGTGGGG


## FIGURE 41 (CONT'D)

CAATAACGGG	GTCAATATTG	GGCATGTCTC	ACCCTGGGTG	GGACAGATAG	AGGCGGGCAG
TTTAGGGGTT	AGACCAAAAAG	GAAGGGGATT	TGTCAGTTTT	GGAATCCTAC	AAACTTGTGG
AGTGGAGAGT	GTTTGCTCAT	CTACTTTCCC	CACCCAATCC	TGTCCACTCC	<u>TAGCCATGAC</u>
<u>ACAGAGCCAA</u>	<u>GAGGATGAGA</u>	<u>ACAAGATAAT</u>	<u>TGGTGGCCAT</u>	<u>ACGTGCACCC</u>	<u>GGAGCTCCCA</u>
<u>GCCGTGGCAG</u>	<u>GCGGCCCTGC</u>	<u>TGGCGGGTCC</u>	<u>CAGGCGCCGC</u>	<u>TTCCTCTGCG</u>	<u>GAGGCGCCCT</u>
<u>GCTTTCAGGC</u>	<u>CAGTGGGTCA</u>	<u>TCACTGCTGC</u>	<u>TCACTGCGGC</u>	<u>CGCCCGTAAG</u>	TGACCCCTC
CCCTGTCCCT	GTACCTAGTG	AATTCCAGAG	TCTAAAGCCC	TAGAGCTGAG	CTGAGAACCT
GGATCTCTGT	ATAGAACCCA	ATGTAGTGGC	TGGCTCCTGG	TTTGAGGTCT	AGAGAAGAGC
CTGGAACAAA	AACACAGCTC	GGGATGTGGG	CTCCTCCATA	AATCTCGAAC	TCAGCATAGG
TTCTGAAAGC	AGATGGGCAG	CTTGGAACCC	ATGGACCTGC	TGAGAACCGA	ACATCTGATC
CAGTGATTCT	TCCAGAGGCC	ACACATTACA	TCGAGACCAA	GCTTAGCCCA	TTCAGATTG
GTGGCTGAAT	TCAGGACCCC	GTCTACATTC	AGAAACTCAG	GACACTACGT	AGAACTCAGA
GCCCAGTTCA	GGAGCTGCAG	TCTAGCCATA	AATCCAGAAC	TAGAACGCTG	CTCACAGCTG
GAACATACAA	CTCTAAGAAAT	AGAGGCAAAA	CCTGGAGGCT	GTTTCACACC	CAAGGTTTAG
TTCAGAGTCT	AGTCTATAGC	TCCGCTATGA	GCAGACTTCA	ACCCAGTGTT	TGAATCCAG
AATGTGGCGG	GTGCGGTGGC	TCATGCCTAT	AATCCTAGCA	CTTTGGGATG	CTGAGGCAGG
CAGATCACCT	GAGGTCAGGA	GTTTCGAGACC	AGCCTGAGCA	ACATAGAGAA	ACCCTGTCTC
TACTAAAAAT	GCAAAATTAG	CCAGGCATGG	TGGCACATGC	CTGTAATCCC	AGCCACTCGG
GAGGCTGAGG	CAGGAGAATC	ACTTGAACCT	GGGAGGCGGA	GTTTGCAAGT	AGTCAAGATC
GCACCATTGC	ACTCCAGGCT	AGGCAACAAG	AGCGAAACTC	CATATCAATC	AATCAATCAA
TAAATCCCAG	AATGCAGATC	CTAATCAGAA	GCCCCATATA	AAACCTAGAC	CCCTCCTAAA
TTCTAGATCT	GAACCTACAA	CCCAGACCCC	AGCCAAGAGG	TCAAAATGCC	TATAAGCCAT
ATCTATGCCA	TAAACAGGTC	AGTCTAGAAC	CTAGAGATCA	AAGCTCAGGC	CAGAGTCTAG
AATATAAAGG	CCAGAATGCA	AACCAGACTC	TAGAATCTTG	GATCCGGGCC	ATAACCTAGA
GCTCCAACCTA	GAACCCAGAG	CCCAACCTGA	GGTCAAGGGC	TAGGGCCAGA	GTCCAGAACC
AAGAGCCCTA	TAATCCAATA	TGAAACAGAC	CTGTAGAGGC	TGGGTGCGGT	GGCTCACGCC
TGTAATCCCA	GCACCTTGGG	AGGCTGAGGC	GGGAGAATCA	CTTGAACCTG	GAGTTGGAGG
TCGAGAGTGA	CTGAGATCG	TGCCACTGCA	CTCCAGCCTA	GGTGACAGAG	CGAGACTCCA
TCACAAAAAA	AAAAATAATA	AATAAATCAA	GTCAATAATC	AGGTTGATC	TAGAATCCTG
ATCTTAGCAT	AGAGTCAAAA	GTTTAAGATG	TCTAGAACTC	AGAACCCAGG	CTAGAAACAG
AATGGTGCCT	ACTCCGGAAT	ATCAGTTCCG	ATTTAGAGCC	TAGACTCATA	ACGCAGTTTC
GCTTAGGACT	CAATGCACCG	AGCCCAGCAC	AGACCCTGGC	ACGGAGCCAA	GCTCTCCCAA
TCATCACCTT	CTTCCCAGC	CAGGAGCTGG	AGCCCAGCCC	AAGAGCGGAA	GGAGAGGCAG
CTGGGGCTGG	GCCGAGAGAA	TGCCCTGGCC	ATGGGGAAGG	GCACAGGAGG	CCAAGAATGC
TCGGCCTGCA	GTTAGTGAGA	AGCAGGCTAG	ACCTCGGGGA	AGACTCGTCA	CCCGGCCCAGG
GAACCGGGCT	GGAGGGTGGG	GAGGAGTCTC	TGGCTCAGAC	CCTGAGCAGC	GCTTCTCTTG
GGGGTCGTGG	CCAGGATCCT	<u>TCAGGTTGCC</u>	<u>CTGGGCAAGC</u>	<u>ACAACCTGAG</u>	<u>GAGGTGGGAG</u>
<u>GCCACCCAGC</u>	<u>AGGTGCTGCG</u>	<u>CGTGGTTCCG</u>	<u>CAGGTGACGC</u>	<u>ACCCCAACTA</u>	<u>CAACTCCCGG</u>
<u>ACCCACGACA</u>	<u>ACGACCTCAT</u>	<u>GCTGCTGCAG</u>	<u>CTACAGCAGC</u>	<u>CCGCACGGAT</u>	<u>CGGGAGGGCA</u>
<u>GTCAGGCCCA</u>	<u>TTGAGGTCAC</u>	<u>CCAGGCCTGT</u>	<u>GCCAGCCCCG</u>	<u>GGACCTCCTG</u>	<u>CCGAGTGTC</u>
<u>GGCTGGGGAA</u>	<u>CTATATCCAG</u>	<u>CCCCATCGGT</u>	GAGGACTCCT	GCGTCTTGGA	AAGCAGGGGA
CTGGGCCTGG	GCTCCTGGGT	CTCCAGGAGG	TGGAGCTGGG	GGGACTGGGG	CTCCTGGGTC
TGAGGGAGGA	GGGGCTGGGC	CTGGACTCCT	GGGTCTGAGG	GAGGAGGGGG	CTGAGGCCTG
GACTCCTGGG	TCTCAAGGAG	GAGGAGCTGG	GCCTGGACTC	ATACGTCTGA	GGGAGGAGGG
GCTGGAGCCT	GGACTCCTGG	GTCTCAAGGA	GGAGGGGCTG	GGCCTGGACT	TCTGGGTCTG
AGGGAGGAGG	GGCTGGGGAC	CTGGACTCCC	GGGTCTGAGG	GAGGAGGGAC	TGGGGGTCTG
GACTCCTGGG	TCTGAGGGAG	GAGGGGCTGG	GGGCCTGGAC	TCCTGGGTCT	GAGGGAGGAG
GTGCTGGGGC	TGGACTCCTG	GGTCGGAAGG	AGGAGGGGCT	GGGGGCTTGG	ACCCTTGGGT
CTTATGGGAG	GGTAGACCCA	GTTATAACCC	TGCAGTGTCC	CCCAGCCAGG	<u>TACCCGCGCT</u>
<u>CTCTGCAATG</u>	<u>CGTGAACATC</u>	<u>AACATCTCCC</u>	<u>CGGATGAGGT</u>	<u>GTGCCAGAAG</u>	<u>GCCTATCTCA</u>
<u>GAACCATCAC</u>	<u>GCCCTGGCATG</u>	<u>GTCTGTGCAG</u>	<u>GAGTTCCCCA</u>	<u>GGGCGGGAAG</u>	<u>GACTCTTGTC</u>
AGGTAAGGCC	CAGGATGGGA	GCTGTGGTAG	GGATTATTG	GGACTGGGAT	TTAAGCAAAT
GATGTCAGGA	GCATGGAAGT	CTGCAGAGGT	CTTCAGAAGA	GAGTGAACCG	CAGGCACAGA
GAGATTCCGA	TAGCCAGGCC	ACCCTGCTTC	CTAGCCCTGT	GCCCCCTGGG	TAATGGACTC
AGAGCATTTCA	TGCCTCAGTT	TCCTCATCTG	TCAGGTGGGA	GTAACCCCTT	TAGGGTAGTT
GGTGGAATGG	GATGAGGCAG	GTTGGGGAAA	GATCGCAGAG	TGGCCTCTGC	TCATATGGGT

## FIGURE 41 (CONT'D)

CTGGGAAAGG CTGTGCTGAG GCTTCTAGAA ATCTTAATGC ATCCTTGAGG GAGGCAGAGA  
 TGGGGAAATA GAAAAAGAGA GACACACAAA TGTTCTACAG TTGGAGCGAA CAGAGAGGGG  
 CCTGGTGAGA TTCAAGGGAC AGGCAGGTGC ACACAGAGAC AGAGCCAGAC CCAGCGGAGA  
 GGGAAGGAAG TGCCCCGACC TCCGGGGCTG AGACCTCAGA GCTGGGGCAG GACTGTGTCC  
 CTAAGTGTCC ACCAGTGTCT CTGCCTGTCT CCCTGTGTCT GCTTCTCGGG TTCTCTGTGC  
 CATGGTGGCT CTGGCTACCT GTCCATCAGT GTCTCCATTT CTGTTCTCTCC CCCTCAGGGT  
**GACTCTGGGG GACCCCTGGT GTGCAGAGGA CAGCTCCAGG GCCTCGTGTC TTGGGGAATG**  
**GAGCGCTGCG CCCTGCCTGG CTACCCCGGT GTCTACACCA ACCTGTGCAA GTACAGAAGC**  
**TGGATTGAGG AAACGATGCG GGACAAATGA TGGTCTTCAC GGTGGGATGG ACCTCGTCA**  
**CTGCCCAGGC CCTCCTCTCT CTACTCAGGA CCCAGGAGTC CAGGCCCCAG CCCCTCCTCC**  
**CTCAGACCCA GGAGTCCAGG CCCCCAGCCC CTCCCTCCTC AGACCCGGGA GTCCAGGCCC**  
 CCAGCCCCCTC CTCCCTCAGA CCCAGGAGTC CAGGCCCCAG CCCCTCCTCC CTCAGACCCG  
 GGAGTCCAGG CCCCCAGCCC CTCCCTCCTC AGACCCAGGA GTCCAGGCCC CAGTCCCTCC  
 TCCCTCAGAC CCAGGAGTCC AGGCCCCCAG CCCCTCCTCC CTCAGACCCA GGAATCCAGG  
 CCCAGCCCCCT CCTCCCTCAG ACCCAGGAGC CCCAGTCCCC CAGCCCCCTCC TCCTTGAGAC  
 CCAGGAGTCC AGGCCCAGCC CCTCCTCCCT CAGACCCAGG AGCCCCAGTC CCCAGCATCC  
 TGATCTTTAC TCCGGCTCTG ATCTCTCCTT TCCAGAGCA GTGCTTCAG GCGTTTTCTC  
 CCCACCAAGC CCCACCCCTT GCTGTGTAC CATCACTACT CAAGACCGGA GGCACAGAGG  
 GCAGGAGCAC AGACCCCTTA AACCAGCATT GTATTCCAAA GACGACAATT TTAAACACGC  
 TTAGTGTCTC TAAAAACCGA **ATAAATAATG ACAATAAATA** TGGAATCATC CTAAATTGTA  
 TTCATTCATC CATGTGTTTA CTTTTTATTT TTTGAGACAA GGTCTTGCTC AGTCTCCTGG  
 TGAATGCTG TAACGCAATC ATAGCTCACT GCAACCGTGA CCTCCTGGGC TCCAGTGATC  
 CTCTTACCTC AGCCTCCCGA GTAGCTGGGA CCACAGGTGC CCGTCACCAT GCCCCGCTAC  
 TTTTAAATTT TTGTGTAGAG ATGAGGTTTC CCTGTGTTGC TCAGGCTGGT CTCGAACACC  
 TGACCCCAAG CAATCCGCCT ACGTCGGTTT CCCAAAGTGC CGGGATTGCA GCGGTGAGCT  
 GCCGCGCCCA GCCTTATCCA TCCAATTAAT GACTTCAAGA AACATGTACA CAGTGGCCCC  
 ACCATGCCAA GCCAGGAGCT GTGTACTGAC AAGTGGCTGC CTCCCTCTTT GCGTGTTTTT  
 CCTTGGGAGT CCCCCGTCCA CCCCCTGTA TCAGGTTTCT AGACGGAAC ACCTCAGCCC  
 TGCAGAGTGA CCTTGAGCAT GACTGCCTTC TACCAGCCTC CTCCCTGGAG CCCCTGTGGT  
 CCAGGGTAGG GAACTAAGTG CCTTGTTCCT TGGAAAATTC TATGCAAATG AAGATGTCTT  
 CATTTTCCTA ATCAGATCTC AGGTGAGGAG AGTTGAGTTA ATCACAGGCT TCAGTCTCTG  
 CCCAGGCAAA GCCCTTCTCT CATTTTATTA ATTTATTTCC ACTCTTCATC TCTGGCTCTG  
 CTCCCCTCCC TCCCCACAGG CACCGACATA AATGGCTTTG AGTGCCCTGC ATCCTTGGA  
 AACAAGGCAG TGTCACAGTG TACTGTTTCT AATTTACATG AAACCATGTG GTTAGGAATC  
 TCATTCTCTT TCTTACTTTT ACTCATCAAC AGCTATTGAG CACCTACTAC GGGCCAGGCA  
 TTGGTCTATT TATTAGGCAC CTGCTATACA CCAGGCATTG TTCTGGGTGC TGGAGGAAGA  
 ACTGTGAGCA AGCCAGTCAG AATCCCTGCC CTCACAGAAC TTATATTCTA GCAGGAGATG  
 ACAGACAAGA AGCCATAAAC ATAATTTTAA AATAAAGCAG AGTCCCTATG AGTAACGAGG  
 TCAATAAACT TGGGCTGGGC GGCAGGCCCA ATGTGTGCCA GGGCCAGCTC ATACATGCTC  
 GCAAGAGTCT ACCAGCAAAT TTTCAGGAAT TTCGAGAACC AGTTGCTAAA TGCAGCCATC  
 ATTAATAAAT AAATTACATA AGCGTATAAT TACATAATTG ATTAATAAATA TTGTCAGTAA  
 ATACTCAAAA CTCAACTGTT GCTAATTATT TCAACTAATA CCTATGCTTG GGAGTGAGAT  
 ATGTCTCTTG TACTACGTCT GTAATGATGA GTTCTGTCAC ACCTCTTTCC AACTCCCCAA  
 CTCTGTCTGC ACCAGTAGCT TGACAATAGC CAAAGAAGAA GTATTTACTG CACTGAAATT  
 GAAAAACACT ATAGATAGGG CTTTGCCGGA CAGTCATTGC TAAACCTTTA CCAGGCACCC  
 TTGGATGGGT CTGCCTGGGA ATGACCTCAT GATCTTAGTG TCTGTCTTCT CAAAGTTCTG  
 TGCTTGAGTA CTGCAGAGTA TAGCTAAAAT AGAATGTTGT ACTCACCTTA TGTCTATGG  
 GGACAGCACA GTATTGGGGA ACCCTAAGGT GGCAGGTCTG GGACATGCAC GAAAGATTGC  
 TGGGAAGTAG AGGCTCCCTC CTTTCTCTCA TCCTCCACC CCATCCTCCA GTGTCTGGTA  
 ACCACCATTG TACTCTCTGC TTCTAAGAGT CTGAGTTTTT TAGATTTTCA ATGTAAGTGA  
 GATCATGCAG TAATTGTCAT TCTGTGTCTG ACCTATTTTCA CTAAACACAG TGTCTCTCCG  
 GTCCATCCAT GTTGTACAA ATGACAGGAT TTCTTTCTTT TATAAGGCAG AATAATATTA  
 AATTATACTG ATACTAATAT ATTACATTTT CTTTATCCAT TCATCCATCA ACAGACACAT


48/51

**FIGURE 42**

**FIGURE 43**

[illegible]

1	PSA	271	285	286	300	301	315	316	330	331	345	346	360
2	hK2	PERPSLVTKVVHYRK	WIKDTIVANP	-----	-----	-----	-----	261					
3	hK1	PEKPAVYTKVVHYRK	WIKDTIAANP	-----	-----	-----	-----	261					
4	HSCCE	PNKPSAVRVLVSVK	WIEDTIAENS	-----	-----	-----	-----	262					
5	zyme	PNDPGVYTQVCKFTK	WINDTMKKHR	-----	-----	-----	-----	253					
6	KLK-L6	KEKPGVYTNVCRYTN	WIQKTIQAK	-----	-----	-----	-----	244					
7	TLSP	PGYPGVYTNLCKYRS	WIEETMRDK	-----	-----	-----	-----	251					
8	KLK-L4	TRKPGVYTKVCKYVD	WIEETMKNN	-----	-----	-----	-----	250					
9	NES1	PDRPGVYTRVSRYVL	WIRETIRKYETOQQK	WLKGPQ				277					
10	KLK-L5	AQHPAVYTQICKYMS	WINKVIRSN	-----	-----	-----	-----	276					
11	neuropsin	DGIPGVYTYICKYVD	WIRMIMRNN	-----	-----	-----	-----	248					
12	protease	SDKPGVYTNICRYLD	WIKKIIGSKG	-----	-----	-----	-----	260					
		VGVPGVYTNLCKFTE	WIEKTVQAS	-----	-----	-----	-----	254					

**FIGURE 44**

## Sequence Listing

SEQ.ID. NO. 1

KLK-L1 na

1 tatctcatga gagagaataa gaacatgaaa agagaaagaa tgagagagag  
agagagaaaag  
61 aaaaaggaga gtggagtcta ggatctgggc aggggtctcc tccttgggtc  
cctagaccct  
121 gctgccagcc ccttctgggc cccaaccac tgcttgggtc gagttgaggc  
agcctgagag  
181 agttgagctg gaagtttgca gcacctgacc cctggaacac atcccctggg  
ggcaggccag  
241 cccaggctga ggatgcttat aagcccaag gaggccctg cggaggcagc  
aggctggagc  
301 tcagcccagc agtggaatcc aggagcccag aggtggccgg gtaagaggcc  
tggtgggtccc  
361 ccactaaaag cctgcagtgt tcatgatcca actctcccta cagctccatg  
tcgctggatt  
421 ctccagctct gtgccttctg tctccacatc tctctagaca gatctctcac  
tgtctctagt  
481 taggagtcac tgtctctagt taggggtctc tctgtctctc tgaatctata  
tctccatgct  
541 taactctcag actgtctctg aggatatctc tcaagcactc tgtctctccg  
gctctgattc  
601 tctgtgtgtc ttcctccat gcttgtttgt ggggtggctag acaccatctc  
tccccattca  
661 cagatggcta gatgtttct ctaaactttc ctttctacct agttctctct  
ctctctcttt  
721 tcccatctct ctctctcttt ttctctctca gtctctaaat ctgtctctct  
aggttctggg  
781 tccatggatg ggagaggggg tagatggtct aggctcttgc ctacctaata  
acgtcccaga  
841 gggagaaaag ggagggacaa agagagggat ggagagactt gggctgaaga  
tcccagaca  
901 cggctaagtc tcagtcctca tcccagggtg ctgacgtgat ggccacagca  
ggaaatccct  
961 ggggctgggt cctggggtag ctcctcttgc gtgtcgcagg tatctgagta  
tgcgtgtgtg  
1021 tgtctgtccg tgcttggggg cacagtgttt gttaatgttc aggtgtgact  
cagtgtctc  
1081 ttgcttgtga ctgcaaagct gcctgtgaga cggtagcgtg ttatccgtcc  
gccatggctg  
1141 tgcccctgca actccttgta tcgtggtaaa tttgtgtgtg gcagtgtgcc  
tgggtgtgtg  
1201 gttgtacctg tgagactctg acagtttgtg cctctgaata tctgggtggag  
tgacaacagt  
1261 gtaatgatga tatggggaca ggggaagccg aggggtgcagg agattgtgct  
tcctggggcg  
1321 tgatccattg ctgggaatct gtgcctgctt cctgggtctt cagtcctgag  
atccccctct  
1381 cccatcccca aggaactcac ctcacaggac tataaacgg tgttttgggtg  
tgcattggct  
1441 tgtggcttgg tgtgactgtg ggcaaggctg ggagaggata ggagtgactc  
ggcgcaggac  
1501 cgactctttg agcatcagtc tgcgcagaca agtgaccga tccttgcctc  
cagcaacaac  
1561 tccacccctt gagctttaat tcaccccgaa ggaccggatc ctaccgctat  
gagcctagac

1621 tctctgttg aacctctct gaccgtggct ttgcaccgag atggcaccag  
tctcacctcc  
1681 agagctcacc ccagagccct gactccgccc cagaagccct ggtccacact  
tctgagactg  
1741 cctctagcca taaccagct cttgaagcct tgatggcgcc cctgcgctgt  
aaccccaacc  
1801 ctaggagcac tgatcccgcc ttctcagccc acccccatgc cctgactctc  
ctcccaggag  
1861 ccctgactac cctgaatccc tgaccaggct cctgcaccgt gatcaccgcc  
cctgggagcc  
1921 ctaggcctat atcctggacc agccctgaa gctccgatca tgaccctgc  
accataaacc  
1981 cacccccagg agccctgggt ccgcccctg gggccgccc cagccctgac  
tcggccccc  
2041 aagagtcttg actgtctctg aagccctgac cagccctg ctcggttaacc  
cctccccaa  
2101 gagccctggg ccgcctctct gagcccttc ccagccctga ctccgccccg  
aggagccctg  
2161 actgtctctg aacctctgac cagccctg ctcggttaagc ccaccccag  
gaaccctggg  
2221 ccgcctctct ggtcccgatc ccatccctga ctccgcccctc aggatctctc  
gtctctggta  
2281 gctgcagcca aatcataaac ggcgaggact gcagcccgca ctgcagccc  
tggcaggcgg  
2341 cactggatcat ggaaaacgaa ttgttctgct cggcgctct ggtgcatccg  
cagtgggtgc  
2401 tgtcagccgc aactgtttc cagaagtga tgagaggta gggggagtgg  
gcagggcctg  
2461 ggtccggggg cggggcctaa tatcaggctc atcttggggg gtcaggggg  
aaacagcgt  
2521 gaaggctctg ggaggaggac ggaatgagcc tggatccggg gagccagag  
ggaagggtg  
2581 ggaggcgga atcttgcttc ggaaggactc agagagccct gacttgaat  
ctcagcccag  
2641 tgctgagtct ctagtgaact aaggcaagtt cttgtccctg aatttttgtg  
aatgaggatt  
2701 tgagaccatg gtttaagtagc tcttaggggtg tttagcgaag aggggtgggtg  
tggggttagg  
2761 agatggggat gggaatgggg ttgaagatga gaatggaggt aaggatgtag  
ttgccacaaa  
2821 actgacctgc cctccgtggc ccacagctcc tacaccatcg ggctgggcct  
gcacagtctt  
2881 gagcccgacc aagagccagg gagccagatg gtggaggcca gcctctccgt  
acggcaccca  
2941 gagtacaaca gaccttgct cgctaacgac ctcatgtca tcaagttgga  
cgaatccgtg  
3001 tccgagtctg acaccatccg gagcatcagc attgcttcgc agtgcctac  
cgcggggaac  
3061 tcttgctctg tttctggctg gggctctgctg gcgaacgggtg agctcacggg  
tgtgtgtctg  
3121 cctctttcaa ggaggtcctc tgcccagtcg cgggggctga cccagagctc  
tgcttcccag  
3181 gcagaatgcc taccgtgctg cagtgcgtga acgtgtcggg ggtgtctgag  
gaggtctgca  
3241 gtaagctcta tgaccgctg taccaccca gcatgttctg cgccggcgga  
gggcaagacc  
3301 agaaggactc ctgcaacgtg agagagggga aaggggaggg caggcgactc  
aggggaagggt  
3361 ggagaagggg gagacagaga cacacagggc cgcagggga gatgcagaga  
tggagagaca

3421 cacagggaga cagtgacaac tagagagaga aactgagaga aacagagaaa  
 taaacacagg  
 3481 aataaagaga agcaaaggaa gagagaaaca gaaacagaca tggggaggga  
 gaaacacaca  
 3541 cacatagaaa tgcagttgac cttccaacag catggggcct gagggcggtg  
 acctccaccc  
 3601 aatagaaaat cctcttataa cttttgactc cccaaaaacc tgactagaaa  
 tagcctactg  
 3661 ttgacgggga gccttaccaa taacataaat agtcgattta tgcatacggt  
 ttatgcattc  
 3721 atgatatacc tttgttgga ttttttgata tttctaagct acacagttcg  
 tctgtgaatt  
 3781 tttttaaatt gttgcaactc tcctaaaatt tttctgatgt gtttattgaa  
 aaaatccaag  
 3841 tataagtga cttgtgcagt tcaaaccagg gttgttcaag ggtcaactgt  
 gtaccagag  
 3901 ggaaacagt acacagattc atagaggtga aacacgaaga gaaacaggaa  
 aaatcaagac  
 3961 tctacaaaga ggctgggcag ggtggctcat gcctgtaatc ccagcattt  
 gggaggcgag  
 4021 gcaggcagat cacttgaggt aaggagttca agaccagcct ggccaaaatg  
 gtgaaatcct  
 4081 gtctgtacta aaaatacaaa agttagctgg atatggtggc aggcgcctgt  
 aatcccagct  
 4141 acttgggagg ctgaggcagg agaattgctt gaatatggga ggcagaggtt  
 gaagtgaatt  
 4201 gagatcacac cactatactc cagctggggc aacagagtaa gactctgtct  
 caaaaaaaaa  
 4261 aaaaaaaaaag actttacaaa gagatgcaga gacactgaga cagataaaca  
 agccacaaag  
 4321 gagacaaagg agagacagac aaacagaaac agacagacca caagcccaag  
 agaagcagcc  
 4381 agcattcagg acataggaca tcgggaagca ggattagatg aagtcaggga  
 tctggaatgg  
 4441 gacttccaac agatatgttg ctgggctatg ttgttattga tgatggttct  
 gtctttgttt  
 4501 ctcagtctca tttagttcct ttctgagccc atatccattt ccacctctct  
 gtgttttgaa  
 4561 ttctgactct ccctctcttc acaacagggt gactctgggg ggcccctgat  
 ctgcaacggg  
 4621 tacttgagg gccttggtc tttcggaata gcccgtgtg gccaaagttg  
 cgtgccaggt  
 4681 gtctaccca acctctgcaa attcactgag tggatagaga aaaccgtcca ggccagttaa

SEQ.ID. NO. 2

KLK-L1

SLVSGSCSQIINGEDCSPHSQPWQAALVMENELFCSGVLVHPQWVLSAAHCF  
 QNSYTIGLGLHSLEADQEPGSMVEASLSVRHPEYNRPLLANDLMLIKLDES  
 SESDTIRSISIASQCPTAGNSCLVSGWGLLANGELTGRMPTVLQCVNVSVVSE  
 EVCSKLYDPLYHPSMFCAGGGQDQKDSCNGDSGGPLICNGYLQGLVSFGKAP  
 CGQVGVPGVYTNLCKFTEWIEKTVQAS

SEQ.ID. NO. 3

KLK-L1

MATAGNPWGWFLGYLILGVAGSLVSGSCSQIINGEDCSPHSQPWQAALVME  
NELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVEASLSV  
RHPEYNRPLLANDLMLIKLDES VSESDTIRSISIASQCPTAGNSCLVSGWGLLA  
NGRMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGGGQDQKDSCNGDSG  
GPLICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTEWIEKTVQAS

SEQ.ID. NO. 4

Figure 4 Sequence

TGACCCGCTG TACCACCCCA GCATGTTCTG CGCCGGCGGA  
GGGCAAGACC AGAAGGACTC  
CTGCAACGGT GACTCTGGGG GGCCCCTGAT CTGCAACGGG  
TACTTGCAGG GCCTTGTGTC  
TTTCGGA AAA GCCCGTGTG GCCAAGTTGG CGTGCCAGGT  
GCCTACACCA ACCTCTGCAA  
ATTCAGT GAG TGGATAGAGA AAACCGTCCA GGCCAGTTAA  
CTCTGGGGAC TGGGAACCCA  
TGAAATTGAC CCCCAAATAC ATCCTGCGGA AGGAATTC

SEQ.ID. NO. 5

Table 8 Sequence

TGACCCGCTGTACCACCCCA

SEQ.ID. NO. 6

Table 8 Sequence

GAATTCCTTCCGCAGGATGT

SEQ.ID. NO. 7

Table 8 Sequence

GGTGATCTGCGCCCTGGTCCT

SEQ.ID. NO. 8

Table 8 Sequence

AGGTGTCCGGTGGAGGTGGCA

SEQ.ID. NO. 9

Table 8 Sequence

TGCGCAAGTTCACCCTCA

SEQ.ID. NO. 10

Table 8 Sequence

CCCTCTCCTTACTTCATCC

SEQ.ID. NO. 11

Table 8 Sequence

ACAATGAGCTGCGTGTGGCT

SEQ.ID. NO. 12

Table 8 Sequence

TCTCCTTAATGTCACGCACGA

SEQ.ID. NO. 13

KLK-L2 na

```
1 gggcccagag tgaaggcaag agaaggagtt gagagctccc tctgcaaagt
ggcttgagtc
61 tcccctgcct aaaatgcagg gagagggagg cagaaagaca gggaagagga
aggggtgggg
121 aagaaagaga gagagagaga gagacagaat aacacaacta cagaaacaca
gagagaacac
181 acagagagcc tgggacacag ggacacacag agtcagagag aaaagagaag
atagagaaaag
241 acacaaatgg agacacagag gtgtaaagaa agagagatta acagagtccc
agatacacgc
301 aaaggggcag aagcacagtt ttcaggggtg tgtctatgat catcttcttt
tttttttttt
361 tttttttttt tttttgagac ggagtctcgc tctgtcgccc aggctggagt
gcagtggcgg
421 gatctcggct cactgcaagc tccgcctccc gggttcacgc cattctctg
cctcagcctc
481 ccaagtagct gggactacag gcgcccgcga ctacgcccgg ctaatttttt
tgtattttta
541 gtagagacgg ggtttcaccg ttttagccgg gatggcctcg atctcctgac
ctcgtgatcc
601 gcccgctcgc gcctcccaaa gtgctgggat tacaggcgtg agccaccgcg
cccgccatg
661 atcatcttct tgactatgct gatgtgacaa gtacctaaag ccatcagact
ctacccttta
721 aatatgcagt ttgggccagg caccgtggct catgcctgta attccagcac
tttgggaggc
781 agaggtgggt gaatcacttg aggccaggag tttgagacca gcctggccaa
catggtgaaa
841 ctctgtcttt actaaaaaaaa aaaaaaaaaa aaaaaaatc agccgggtgt
cgtggggcac
901 acctgtaatc ccagctatgc tggaggctga ggcacgagag tcacttgaac
cctggaggcg
```

961 gaggttgtag tgggcccaga tcacatcacc gccctccagc ctgggagaca  
gagcaagact  
1021 ctgtctcaaa taaataaata aacaaacgaa caagcagttt gttgtacctt  
agttatatct  
1081 aaaaaaaaaa tgctgtcaac aaatagagca gaagtgaat aaaggaaaat  
aatgggcca  
1141 agaactctaa ggtatatattg acaaatcatt cagaaccttt aaaaaagaaa  
gaatcacaga  
1201 ggcataagaa gacagggagg aacagggaga cagaaacacc tgtggcccaa  
ggagaacaaa  
1261 acaaggctcc taagacagac aggaggagag agagagagag tgagttagag  
acagacagag  
1321 aaaaagacag agagagagag acagagacag agagacagag aggcgagagg  
gatagaaaga  
1381 gagagagggg ttgagagaga cagcagatat tgagagagac tcagaaagat  
agccgaggga  
1441 gaaccacaga gagatggaag aagactctga gaaaaaacca gagacaaaga  
tggaagagg  
1501 agtatcgagg gtgaacagac agtgggtggaa tgagcaaaat gcagagaaga  
aagcaagcaa  
1561 tccaggcgcc aagaatagtg acccagagtt ggtgagaagc cagatcctta  
aggctggggg  
1621 aggcagggaa ggggctggcc tggcttccgg agacctctcc ccattctccg  
ggccaggag  
1681 gtagggagtg acattccgga ctgggtgggg ggtgctctgg gggtagagat  
agggggagca  
1741 ggaggagcta ttgctaaggc ccgataggca cctcattgcc cgggaatgtg  
ccccaggag  
1801 cagtgggttg ttataactca ggcccgggtg ccagagccca ggaggaggca  
gtggccagga  
1861 aggcacaggc ctgagaagtc tgcggctgag ctgggagcaa atccccacc  
ccctacctgg  
1921 gggacagggc aagttagacc tggtaggggt ggctcagcag gcagggaagg  
agaggtgtct  
1981 gtgcgtctg caccacatc tttctctgtc ccctccttgc cctgtctgga  
ggctgctaga  
2041 ctctatctt ctgaattcta tagtgcttg gtctcagcgc agtgccgatg  
gtggcccgtc  
2101 cttgtggttc ctctctacct ggggaaataa ggtaggggag ggagggaag  
tgggttaagg  
2161 gtcccccgga tcgcctgggc ctcccaacc tctgacattc cccatccagg  
tgcagcgcc  
2221 atggctacag caagaccccc ctggatgtgg gtgctctgtg ctctgatcac  
agccttgctt  
2281 ctgggggtca caggtaacca gaactctggg gtgggagggg tgtgggattg  
ggaggactgt  
2341 ctctgaggca ctagagcgcc tgtcccctgg ggaactgtgt gagcctgggc  
atgactccgg  
2401 gaccgggtga atgtgagtct ctgtctgtac ttgtggttgt gcgacgtat  
gtggccctgt  
2461 gactgccacg gtgtgtgtcg gggaggggga tgccttttcc catatcaggt  
gactgtgcg  
2521 cagggtggac tgacctttg aggcgtgtgt tgtggttttg tgattgtgtg  
tgcatttaag  
2581 attgtgtgtg gctccacagc tgtgtgggtg aatgcatgta gactggggg  
tgttactgt  
2641 gtgtttggct gtgtgtgtg acttggcatt gtatatgact gcaggtatct  
gcagttcctg  
2701 tccctgaggt cccgggattg cgtgcaacaa aagtgggtcat caccatggaa  
agctgtgact

2761 gtgtgctgct tgcaggcgat tatgtgattg tggctgagtg tgacgttatg  
gatgcccgtg  
2821 tttgtgaccg tgtgactacc tgaagctctg tgtaggggtg actgtatgtg  
actgtgtgtg  
2881 tctgtgtgag gccgtgtaaa tgctactgta tgtgtgatgg tgcagctgtg  
tgtctggagt  
2941 ttctgtctct gcctggaggg atagaggggtg caggggtagc tatctctggg  
agatgggtgc  
3001 caggtgactg acttgcaagt tgtgcctgtg tgcagaagag tatgtggcag  
tctgaacatc  
3061 tgtgcacaca cggcatctgt gcgtggcact gagacactgt ggatgagggg  
gtgcgatccc  
3121 gctaggctgc ccgggagcgt gtgtacctgg agacagagct gtatgttagc  
tgcacctgtg  
3181 gaggcaacat gggcgtgtct gcagaactgc gtgcgtgctt ggctgttact  
gctgttgtgc  
3241 gcgtgggtct tggggtgagt tcgtgaatga tgggtggtgcc agggccatca  
gcaagggtaa  
3301 gaaccaggcc gggcgcggtg gctcacgcct gtaatcccag ccctttggga  
ggccgaggca  
3361 ggcggtacac ctgaggtcgg gagatcgagg ccagcctgac caacatggag  
aaccctgtct  
3421 ctactaaaaa taaaaaaat tagctggtgt ggtggcgctg gcctgtaatc  
ccagctactc  
3481 gggagactgg ggcagaaaaa tcgcttgaac ccgggaggtg gaggttgcgg  
tgagccgaga  
3541 tcgcgccatt gcactccagc ctgggcaaca agagcgaaac tccgtctcga  
aagaaaaaaa  
3601 gaaaaaaaaa agggtaagaa ccagtgaatg ggcacgggag gactgatgat  
ggagtggggc  
3661 atgcatgtag tctgtaggtc tgtgtgtgag aggaggagat tgacaggatt  
gagaaggcat  
3721 gttttcatct gagaattcag aaacctaggc ctgctcttcc cctccatgtg  
gccccctaag  
3781 ctgagccctt ctttcctggt cctgctttcg gaacctagc tccgcccatt  
agctctgacc  
3841 ccacctcctt tcctcaacca cgcccctagg ccagactcta gtggaccccg  
cctaaggcca  
3901 cacccttttg ggccaggctc cacccttat tctgtgggta ctttctagaa  
cccccttcaa  
3961 agtcagagct tttttttttt tttttttgga gacagtcttg ctctctctcc  
caggctggag  
4021 tgcagtggcg tgatctcggc tcaactgcaac ctctgcctcc caggttcaag  
tgattctcgt  
4081 gcctccacct cctgagtagc tgggattaca ggtgcgcgcc accacgcctg  
gctaattttt  
4141 gtgtcttttag tagagacagg gtttcacctt gttggccagg ctggtctcaa  
actcccaacc  
4201 tcagggtgatc cgcccacctc ggcctcccag agtgctgggg ttacaggcgt  
gagccaccgc  
4261 ccccagccca aagtcagagc tctttatagg agactctaac atgtaaccct  
gacctgggc  
4321 ctaactaagt caattccaaa ccccttctg cctccagccc tgacccact  
cactgaggcc  
4381 tgacccact tcttgagacc agttccatcc cttaaagccct ggtctccctc  
ccatccccag  
4441 gctccagccc ccacagcttt ggcactaccc ctgagcttgt ccaggaatcc  
tgtaccat  
4501 ttaccctca catgtagttc tagccaatc caggaatctg tgaggtccag  
ttagagtcca

4561 gtaaccctac ctgagcctgg gctctgtcct tgagcttgag cctgggcttg  
agaggtgcca  
4621 ctcttattct ccaggccctg cccctgcccc ctcagcatgt cagacacca  
ccctctagct  
4681 ggtctggcct cttgagtctg aaaccacccc ccagcccaag ccccgctct  
gagccccgc  
4741 caaccattt tccgttccca gagcatgttc tcgccaacaa tgatgtttcc  
tgtgaccacc  
4801 cctctaacac cgtgccctct gggagcaacc aggacctggg agctggggcg  
ggggaagacg  
4861 cccggtcgga tgacagcagc agccgcatca tcaatggatc cgactgcgat  
atgcacaccc  
4921 agccgtggca ggccgcgctg ttgctaaggc ccaaccagct ctactgcggg  
gcggtgttg  
4981 tgcattccaca gtggctgtc acggccgccc actgcaggaa gaagtgagt  
ggagttccaa  
5041 gaggaggggt ggtggggacg ggggaagtggg ggtgggggtg ggggaagtgg  
ggtgggggtg  
5101 tcatggaggt gagggtgtg ggggacggg aagtgggggt gggggtgtca  
tggaaggtga  
5161 ggggttgttg ggatgggttg gggatgtggg agcaggagga ggtcgagttg  
gggataggac  
5221 taaggatgga gttttgcggg ggagcaagg gggaggtga ggttgagag  
gggagagtgt  
5281 tgtggtagg aatgggaagg agccaaggat gggttggatt tggggttagg  
agcatatatt  
5341 tgttgaatgg tttgggatgg aggtggaatt gggattggct ttagaattgg  
gggtgggtga  
5401 aaatcgggct ggggtggaaa tgaagatagc atggagatag ggttgagatt  
gggagcagat  
5461 atagaatgaa ggatggggat tggagttttg ggtgggggtg gagatgggtg  
gatttgggct  
5521 tgagaatgca tatggtgatg gcttctgggt agggaaagaa ttaggggttg  
gaatgggatg  
5581 ggtttggaat tgtactggg atggggacag gcatgggatt ggagaccaag  
aggaggtga  
5641 ggatgggttg gggaccggg gtggggatgg ggggtgggct ggggctgggt  
gtggggttg  
5701 gattggcggt ggacgtggag atagagatca gggttggttg tgacctgccc  
catcttctc  
5761 agagttttca gagtccgtct cggccactac tccctgtcac cagtttatga  
atctgggcag  
5821 cagatgttcc aggggggtcaa atccatcccc caccctggct actccaccc  
tgccactct  
5881 aacgacctca tgctcatcaa actgaacaga agaattcgtc ccactaaaga  
tgtcagacc  
5941 atcaacgtct cctctcattg tccctctgct gggacaaaagt gcttggtgtc  
tggtggggg  
6001 acaaccaaga gcccccaagg tgagtgtcca ggttcttctt gataccgacc  
catctctgcc  
6061 gccttccatc tttctccact tctcattgtg ttctgtttg acagtgcact  
tccctaagg  
6121 cctccagtgc ttgaatatca gcgtgctaag tcagaaaagg tgcgaggatg  
cttaccgag  
6181 acagatagat gacaccatgt tctgcgccg tgacaaagca ggtagagact  
cctgccaggt  
6241 gaggacacct ctctttatc agcagataca cactgagtgc caactcggt  
acatggagc  
6301 ttgcaaatc ctgagaatcc agcaattgcc aagacagtca ggaccctgt  
tctcacagag

6361 ctcataccct agagtagtgg tgtttagtag aaataatgct gagctgctta  
tgtcattttcc  
6421 agtttttttag tagccacatt aaaacaggta aaaaaggctg ggcgagtg  
ctcacacctg  
6481 taatcccagc actttgggag gctgaggcag gcagatcacc tttggtcagg  
agtttgagac  
6541 tagcctggcc aacatggcga aactctgtct ctaaaaaaaaa atacaaaaat  
tagcctggca  
6601 tggtggcggg cgcctgtaat ctgagctgct caggaggccg agacacaaga  
atcacttaaa  
6661 cccaggaggt ggaggttgca gtgagctgag atcgtgccac tcaactcaac  
ctgggagaca  
6721 gagtgacact tttgtctcaa aaagaaaaa aaaaacaagt aaaaaagaaa  
caggtgaagt  
6781 taactttaat aaccaatgt atcccaaata caatcatttc aaagtgaat  
taatataaaa  
6841 caattatgaa tgagatactt tacattcttt tcttgttttc atattaagtc  
tttgaaagtg  
6901 agtatatatg ttatgctgac agcacatctc aatttggact agctacattt  
caggtgctca  
6961 gtagccacat gtggctagca gttactgtat tggatggcac ggatctagag  
ggaaagatca  
7021 gggctgtttt gtatggttg gtaggttgct cactgcataa agataccata  
tctaataagg  
7081 gcactccgtg ttacagatgt cagttttggc agttttcagg cgtgtggtag  
ttaagtgtct  
7141 tgtttcaaca aaatctgtaa tatgacagtt ttctagcaag tgctggtaaa  
atatcttgag  
7201 gaaggaaaag agaaatctgg taggtatttt tacaagagaa tatttaatac  
aggggattaa  
7261 ttgcaaagct gctggaagg gctggaggaa aaagttaaaa aataaaaaac  
tctgtggtca  
7321 agaactctgca taaatagggc aatttcagag agtggttaaag gttaaccca  
aaataaaaca  
7381 tggtttttagg atagtaaaca ataagggcc atattcaaaa aggtggtcag  
gggagcctcc  
7441 ttggagaggt ggcatttgag cagagaatgg atgacacaaa gaagctaaac  
tcgtgaagtt  
7501 taaggggaaa gaaaaggcac gtgcaaaggc cctgaggcag taaggaattt  
ggctgattca  
7561 aagaagaaga ggaaaccaat gcaactggag aacaaaagt ggggcaacag  
tagaaagtga  
7621 cgctggaggt gtaggcagg gcaaatgtct tgcaagtatt tcttggtcac  
caacacagag  
7681 cttccctatg ttctaattga agctgtatct gttgaggaag acagaattta  
aatcaaact  
7741 gttacatcaa ccagaccct tctctgtatt caggctcca agggatctag  
aaggacgtaa  
7801 gttaacaagc tctcattagc aggggtgtgtg tttcaacagt agttaggaag  
ctggggattc  
7861 aggagtactc cagtcccatg gctatgaaaa gctccccca aattgtacaa  
acctgacaaa  
7921 tgcaacacct cccagctct cccatttct tctctgtgcc ctgggtgtgg  
gggggtgggt  
7981 tgcgagggg aaaactttta acagaagaaa gcacatctcg gccgggcgtg  
gtggctcaca  
8041 cctgtaatcc caacactttg ggaggccgag gcgggtggat cactaggtca  
ggagatggag  
8101 accatcctgg ctgacacggt gaaaccctgt ctctactaaa aacacaaaaa  
attagccggg

8161 cgtggtggca ggcgcctgta gtcccagcta ctcgggaggc tgaggcagga  
gaatggcctg  
8221 aaccgggag gcggaacttg cagttagccg aggttgacc actgcactcc  
agcctgggca  
8281 acacagttag actccgtctc aaaaaaaaaa aaagaaaaga aaagaaatca  
catctcattc  
8341 aagtgggtggc atttaaaact atttagcctt tctgtaggca aggttagtat  
cttggttttc  
8401 cagacctcaa ggtgtttttt tgtttgtttt ttcataccgg tgtgtggtct  
gggtgtggcc  
8461 actaaaagct acaagcaaga aataataaca actacaacaa tactaatacc  
aatagtataa  
8521 aaataatagc atctggctaa ttgctggaca ctgttttaag tggtttgcac  
gcctcagctc  
8581 attaactcat ttacctgtta ttattggccc tattttacaa acaaggagcc  
aaggctcaga  
8641 gcagttaact aacagcctct caaaagaaac tctgcagaga tattaaattt  
aaaaaataat  
8701 gagagaaatt aaaccacaag aaagttgaaa ttagaggta caggcagcta  
agcttggttg  
8761 ctttgaaaca gtgtctgcta ctgggaaaaa ggcaagtctt ggctttccta  
ataattgata  
8821 ccaggactct gtaattcata ttttgcacgc atgtaagtaa gaaatgaagc  
cgggtgcaat  
8881 ggcacatgcc agtaatccca gcactctggg agactgaagt gggaagatca  
cttgagctca  
8941 ggagttaag accagcctgg gcaactaaaa attaaaaaaa taaaaatact  
aattgttttt  
9001 attttagtag attttattca taccacttac atcattattg tagtatgtac  
atattttatt  
9061 cttttctttt cttttctttt cttttttgag acggagtctc gctctgtcac  
ccaggctgga  
9121 gtgcaatggc accatatcag ctactgcag catgcgcctc ctgggttcaa  
gcattttctc  
9181 cacctcagcc tccaagtag ctgggataac aggcacccac caccatgcct  
ggctattttt  
9241 ttttttccgt agagatgggg ttccaccatg ttggccaggc tggctttgaa  
ctcctgacct  
9301 ccagtgatct gcctgcctcg gcctccaaa ttgctggtat tacagggtgtg  
agccaccgtg  
9361 cccagggtggg agatagacat ttctctctac ctcaaacaga ggtccactca  
agctactttt  
9421 cattttcttc ataaatatta gccgagtggc tattttgcac caggaatggt  
tccagggtgct  
9481 gtggatatgg catcaggcaa aacagaccaa aaacttcctg ccgcgtggac  
ctcatgttcc  
9541 ccaagtggaa gacaggcaat aaagagatag ataaatatgt agtaaattaa  
aaaaaaaaaa  
9601 aattagccgg gtgtggtggc ttgcacctgt agttccagct acttgggagg  
ctgaggtggg  
9661 agaattgctt gagcccaaac gtttgaggct gcggtaagcc atgactgcac  
tgctgcactc  
9721 cagacagcag cctgggtgac aaagcaagac gtttttgtca gaaagaaaaa  
aaaaagagac  
9781 gaagggagga aggagagaga aaggaaggaa ggaaggagaa agaaaggaag  
gaaggagaaa  
9841 gaaaggaagg aaggaaggag aaagaaagga agaaagagaa agaaagaaaa  
agaaagaaag  
9901 aaagaagaaa gaaaagagag aggaaggaag gaaagaagga aaagagggaa  
aaaaatgact

9961 gttgaagagc agtgagtatt attataggag ggtaattata gggaggtatg  
gggaattgaa  
10021 gacaggaaac acaaattagt ccaagcgaat ggatttctat tgggagtgat  
tctgccccta  
10081 gaagacactg gcaataccag gagacatttt tggttgtcac aactatatgg  
aggggcatta  
10141 ctggcaacta atggatagat gccaaagtgtg ctgttcaaca tgctatgatg  
cacacggcag  
10201 gcctccacaa caaaccatta tccagcttca gatgccca gtgcccagat  
cgaggaaacc  
10261 tcatccaggg gctgagaacc gtatttttgc agaagggagg tataaggatg  
ggttgggtgga  
10321 gaatggggaa ggaaggtgtg tgtccagtaa gagaaataag gcctgcacag  
gctggagggg  
10381 agagtgagag agaaagggag gcggagagat acacgatgag ggagacaggc  
tggaacagaa  
10441 agtagagacg aagattcgag atgtggagag gaagggtcac agaccccc  
gaaatgatgt  
10501 gtggacaaca ggaatctgga agaggaagat ggagtggaga gtgacaaatg  
gggtctaaag  
10561 gttgaacttg gaggccaggc atggtggctc acgcctgtaa tcccaacact  
ttggaggctg  
10621 aggtgggcca atcacttgag gccaggagtt cgagaccagc ctggccaaca  
tggtgaaacc  
10681 ccgtctctac aaaaaaata caaaaatta gccgggtgtg gtgatggaca  
cctgtagtca  
10741 cagctacttg ggaggctgag gcaggagaat tgcttgaacc cgggagatgg  
aggctgcagt  
10801 gagctgaggt caggccactg cgctccaacc tgggcaacag agtaagactc  
catctcaaaa  
10861 aaaaaaaagc tggatttga gtgaaatatt aataacattc tccctctctc  
tccttttgcc  
10921 tgtgtctcca tctctgtctt tttctgcatt tcttcatctc tgtactttcc  
atctctgtgt  
10981 gtctgttccc atctgcttct ccatctatgg gcattctctg gtctctcatg  
tctccttctg  
11041 ccacttttgc cacatctctg cctctctcat gcccccttt ctctcctgca  
gggtgattct  
11101 ggggggcctg tggctctgca tggctccctg cagggactcg tgtcctgggg  
agattaccct  
11161 tgtgcccggc ccaacagacc ggggtgtctac acgaacctct gcaagttcac  
caagtggatc  
11221 caggaaacca tccaggccaa ctctgagtc atcccaggac tcagcacacc  
ggcatcccca  
11281 cctgctgcag ggacagccct gacactcctt tcagaccctc attccttccc  
agagatgttg  
11341 agaattgtca tctctccagc ccctgacccc atgtctcctg gactcagggt  
ctgcttcccc  
11401 cacattgggc tgaccgtgtc tctctagttg aacctggga acaatttcca  
aaactgtcca  
11461 gggcggggggt tgcgtctcaa tctccctggg gcactttcat cctcaagctc  
agggcccatc  
11521 ccttctctgc agctctgacc caaattagt ccagaaata aactgagaag

SEQ.ID. NO. 14

KLK-L2 aa

MATARPPWMWVLCALITALLGVTEHVLANNVSCDHPSNTVPSGSNQDLG

AGAGEDARSDDSSSRIINGSDCDMHTQPWQAALLLRPNQLYCGAVLVHPQW  
LLTAAHCRKKVFRVRLGHYSLSPVYESGQOMFQGVKSIPHPGYSHPGHSNDL  
MLIKLNRIRPTKDV RPINVS SHCPSAGTKCLVSGWGTTKSPQVHFPKVLQCL  
NISVLSQKRCEDAYPRQIDDTMFCAGDKAGR DSCQGD SGGP VVCNGSLQGL  
VSWG DYPCARPNRPGVYTNLCKFTKWIQETIQANS

SEQ.ID. NO. 15

Table 11

GGATGCTTACCCGAGACAGA

SEQ.ID. NO. 16

Table 11

GCTGGAGAGAGATGAACATTCT

SEQ.ID. NO. 17

Table 11

GGTGATCTGCGCCCTGGTCCT

SEQ.ID. NO. 18

Table 11

AGGTGTCCGGTGGAGGTGGCA

SEQ.ID. NO. 19

Table 11

CCGAGACGGACTCTGAAAAC TTTCTTCC

SEQ.ID. NO. 20

Table 11

TGAAAAC TTTCTTCTCCTGCAGTGGGCGGC

SEQ.ID. NO. 21

KLK-L3

1 cttgaaccca ggaggcagag gttgcagtga gctgagatcg cgccactgta  
cttcagcctg

61 ggtgtcagag caatactccg ttttggaaaa caaacaacaa aacaaacaaa  
caaaaaacag

121 atggagcaac tgagagaggt cttgtgactt gcccagagtc acacacctca  
 tcactaatca  
 181 cacctaataca ttgagatttg gacacacatg gttcagttcc agagtccatg  
 ctccaaacca  
 241 tgacgacaca gtgagagaac attcaagggg agcccagacc cagcttcata  
 accaggcctg  
 301 tgagcaggag aaagtggaag ggatcgtaag tgcccagggg aggcaaagat  
 ggactctgcc  
 361 tgaggatctc agagatttcc tggaggaggg agaattgagg ttgggtgttg  
 aaggatgagt  
 421 gggagttcac caggaaaaga aggatatgga gaaagacatt cactcattca  
 atgaacatct  
 481 cctgaggact tctgcaagcc ctgttccgcc tggaacgggg tgatgctggg  
 acacagagat  
 541 gagtcagacc tgggcccagc cctccagaag ctgtccacct ggtgagaagg  
 aatgatgagg  
 601 agagaggcag ggaggatggg gtgatggaag ggacaatggg gtggggggca  
 gggagatgga  
 661 tgaaaaaat atatagcaaa tgttctcagg atttggcaaa gatcaggatg  
 tattaagaga  
 721 gagcacaggg cacttgctac ctggaagggt gggcacctgg gtccttgggt  
 ggtggagccg  
 781 tggggaaggg ggcagggtat gacaagagtg ggttaatcca gatggaacca  
 gatttctcaa  
 841 cattctagga gagggccttg tccttgtggg aagaggccca aatccccagg  
 gcagggaagg  
 901 ttctgcaagg tgtgtaaacc tgtgcagctg cctgtggtct ctgcctcact  
 ccactggat  
 961 ttccctcaat ctttcccggt ttctgtctcc tctccact ctcctctca  
 tcttgggtcc  
 1021 ttctgtgctt gtacctccct ctcttgtat cttttgctct tgtgtctgag  
 tctgactct  
 1081 gtcttcacc cctgcctcc tttctgggtg gtccccctgc acatccctcc  
 agcctgccgt  
 1141 gggaggttgg tctctgcaca ccaactgctt atccaaaata aacctgctgc  
 accccaggac  
 1201 cttaggcttc aaggatctcc ctcttttcc aggacacaaa agattctgta  
 tctttagacc  
 1261 taaggtgatg aggaatgagg tctccactc tgaagacccc agaggagggtg  
 ccacaaact  
 1321 ctccacacc ccagcactcc tctccattc agtcaagctc tggcccagca  
 agccgccagt  
 1381 tcatcccaaa aggggggtcc cctgcactt acctctctc ccaaggcccc  
 tgtcacagcc  
 1441 ccagggttc cccctcccc aggtacattt cccaaccccg attaatcaca  
 ggggcggccc  
 1501 catggaggag gaaggagatg gcatggctta ccataaagaa gcactggacg  
 ccgggtgcac  
 1561 gttccaggat ccagggtgcc aggggtcatg aagctgggac tcctctgtgc  
 tctgctctct  
 1621 ctgctggcag gtgaggctcc caggctggct gccccttcac ggctgtacta  
 aggtcacctt  
 1681 gctcttccct cccatcccag gcttctgcct cctgcctct aggttctca  
 gcactctctc  
 1741 cctgcctcc cagcctgctc ttcgctgacc ctttgtccc tcatccccac  
 ccagggcat  
 1801 ggctgggcag acaccgtgc catcggggcc gaggaatgtc gcccacactc  
 ccagccttg  
 1861 caggccggcc tcttccacct tactcggctc ttctgtgggg cgacctcat  
 cagtgaccgc

1921 tggctgctca cagctgcccc ctgccgcaag ccgtgagtga cccaggctgg  
ccatgctggg  
1981 gagggacaga ggctgggggt caggagaggg tgaggggtgc tttaggccag  
aagtgcggag  
2041 cctccacttc tgataccaca agttcaactc ttagaagtag gaagggtagc  
ctcccaaadc  
2101 ctaaaattct agagaccagc aatatctcat ttgagaagtc taagattcga  
aacttaggct  
2161 cttcgaatcc gagactgacc cagagaaatc cagaatcgta gaatcctaaa  
atcttgaatt  
2221 tatgaaatc tgcaatagcc tcagcaaatt ttagaatcat agattcgcag  
actattagaa  
2281 tcttagcagt ctgggtcagc actgcccaga ggaattatga tgccagccac  
atgtgttaagt  
2341 ttaaaattct ggtggacaca tttaaaaaat aaggaatgag taaaattaat  
tctaatagat  
2401 ttaacttgac atacccaaaa acttattttg acatgtaatc aatttttaaa  
tacgtatgaa  
2461 cgatacagtt tacttttgtt ttggtactaa gcctttgaaa tctgttctgt  
attttacaca  
2521 catagcctgt tacaaaatgg actagccaca tttcaagtgt tcaatagcca  
taatggctag  
2581 tgtgatccta gaatcttaaa ttcagagctt tctagattca ttgaatattg  
aaactcacag  
2641 tactagaatc tttgattcac agtatcctag aatattgaga ttcagataat  
tctgtagtct  
2701 taaactatct gaatcccaga ctcttaaatt tctaaggtta tagatttata  
gaatgatgac  
2761 attctagtct ttcttttttt tttttttttt tttttttgag acagagtctc  
cctctatctc  
2821 ccaggctgga gtgcagtggc acaatctcag ctactgcaa cctctgcctc  
tcgggttcaa  
2881 gcaattctcc tgcctcagcc tcctgagtag ctgggattac aggtatgcac  
caccatgcca  
2941 ggctattttt tttttttttt ttttttttagt agagacgggg gtttcacat  
attggccagg  
3001 ctggtcttga actcctgacc ttgtgatctg cccgcctcgg cctcccaaag  
tgctgggatt  
3061 acaggcgtga gccaccgccc ccagccaaaa ttctagtctt tttgtcctag  
aacattaaaa  
3121 ttctatgttc aaatcttaga ttaattcag ataatgtag aatcctggag  
tttttttgat  
3181 ccaggggaat ctggaatgtt agaactcttg attcataaaa ctctaaacct  
tgagcctcta  
3241 gattctagaa tcatggataa tagtgtgtcg gaatctgaga attctagaat  
cttaggttct  
3301 gggcattcta atagtatcct ggaatccacc tgatgcagga atcctctctc  
cattgcctct  
3361 gaaaagtgc catccatact gttccaattt tcttccctcc atgagtaaag  
cactgattgt  
3421 ggtaagagat gctgtgtggg aatttcccat catgcattgc tccatgatgg  
aacctcctt  
3481 aacttaagcc tatacatcag actgggagaa cgatgttcag atttcagccg  
aaagtgaagc  
3541 aggagaaatg cagagatatg aaggtggaag agagtgagag gcaggggaag  
ggtaggggga  
3601 tgaagggatg taggggtgag gactactttt ccagatccag agccaagaca  
gcaagaatga  
3661 cagagagaga cagacacaga tgtttctggt tccccaacct tgaattcgca  
gtcattagcc

3721 tgctgcctaa tgtcagaggt cagaggctgg ggaatggact tgtcatcccc  
gaaaggatcc  
3781 cagctgtcta gggcatggac cagaaatgaa acaagtgcgc tgagactgtg  
gtgagggctt  
3841 aaggtagac accaggaaga catgcattga aggggaagg atatgataga  
caggaaaagc  
3901 tgaggccaga gatgaccccc aatttgggga ttttccatat cccatcccct  
ttcatacaca  
3961 cgcacacgta tacacacaca ccacttagac atacagagcc gctcccacag  
aagccaccag  
4021 acctgtgggg gcaggggtgg ggcggttgtt atgtggtagg tgggggtccc  
cgtgcccaca  
4081 ccgttcctag ggaccaagt caccaccaag gctccagggt agtagggagg  
aagggtggctc  
4141 actcagcctg ggactaggag cgggggcttt gtggggagag ctacaaagat  
ggagacacac  
4201 aaaacatcag agtggggacc agggaccag aggaggtgtg tgcctcgctt  
aaaatcacag  
4261 taccctgggc cagacataga tgatgagggg gcagagaggg tgtgtggctt  
gcagaggggtc  
4321 acacagcacc ctgatggaca ggaaaagagg gctggggctg aaaggacttt  
tacctttccc  
4381 ccagcttgac ctctgaggcc tgtcccagca ggtatctgtg ggtccgcctt  
ggagagcacc  
4441 acctctggaa atgggagggg ccggagcagc tgttccgggt tacggacttc  
ttccccacc  
4501 ctggcttcaa caaggacctc agcgccaatg accacaatga tgacatcatg  
ctgatccgcc  
4561 tgcccaggca ggcacgtctg agtctgtctg tgcagcccct caacctcagc  
cagacctgtg  
4621 tctccccagg catgcagtgt ctcatctcag gctggggggc cgtgtccagc  
cccaagggtg  
4681 tgacctggcc cagaactctc tctgaaactt gctccctcac ccctctgtct  
ctgccttttc  
4741 atctctgtct tctccttttc tctctctct ctctctctgt cagtctatct  
atctgccaat  
4801 cgatatat tt aaccaaatat aagatgctag catttttaag atgtgccatt  
atttcatgaa  
4861 ctgcgaagaa gtggaagaag gaggaggagg agaagaaaaa aaggaggagg  
aggaaagatc  
4921 ccattagatc ccattgatta tataacacca ttttctggaa gacacattct  
aatttcagag  
4981 tgtttgtttg tttgtttgtt tgtttgtttt tgagacaggg tctcgctttg  
ttgctcaggc  
5041 tggagtgcag cgggtgtgatc acggctcatt gcagctttga actcctgggc  
tcaagtgatc  
5101 ctctcgctc aacctcccaa gtagctggga ttacagatat gcaccaccac  
atcccacacc  
5161 ggggtcattt ttttattatt tattattatt attattatta tctttttttt  
tgtattttta  
5221 gtagagacag aggtttcacc atattggcca ggctggctct aaattcctga  
cctgggtgatc  
5281 tgcccgctt ggactcccaa agtgctggga aaacaggcat gagccactgc  
accagccaa  
5341 aattctagtc ttttttaaat ctagtcatat cttagattta attcagataa  
tgttagaatc  
5401 ctggagtttt ttgatccagg ggaatctgga atgttagaat cttggattca  
taaaactcta  
5461 aacgttgagc ctctagattc tagaatcatg gatactagtg tgtcagaatc  
tgagaattct

5521 agaattcttag attctgggca ttctaatagt atcctggaat ccacctgatg  
caggaatcct  
5581 ctctccattg cctctgaaaa gtgaccatcc atactgttcc aattttcttc  
cctccatgaa  
5641 taaagcactg attctggtaa aagatgctgg gtgggaattt cccatcatgc  
attgctccat  
5701 gatgggacct cctttaactt aagccttatg ctaaaaattt ttattatatt  
tagcaaagat  
5761 gaggtcttgc tatgttgtcc aggctagtct caaactcctg gcctcccaaa  
gtgctgagat  
5821 tacaagtgtg agccactgta cctggcccag agatgtttaa atgtgaaatg  
cgttcatctt  
5881 agaatgggaa taagaccatg tctctcagag tcacggatca ctgaccatt  
agccaaattg  
5941 ggtcagtgga ttggaaaaac agtctgaatt tgttgctgcc aatatctaaa  
acttggaag  
6001 ttttatacaa aagccaggtt tctggattca cctgaaaaag ttgaagaac  
tcacattccc  
6061 aaaatagcaa gcattgggct gagtcaatgg aggctgcccc cttcagcaa  
gataagtctt  
6121 ctgattcact ccaatggacc caaatggctc ctgtctcctt gcacagcccc  
cgtccccgac  
6181 ttctgtttac caattctgtt tatcatatcc ctgatgcatt cggagcctgc  
acccatgtct  
6241 tatatagatg cacatgtgta ttatatatcc atatccacat ctatactgac  
tacactgtat  
6301 ctggtatctc tgtctatgtc tctgtctcca tcagtgaacca tcttcctgca  
aatctctttc  
6361 cttttatctc actgccttca ttccaccctt tgaggctctgg gtctttttct  
atctcttttt  
6421 tttttttttt taagagactg agtcttgctc ttgttgccca ggctggagtg  
cagtgggtgtg  
6481 atctcggtc actgcaacct ccacctcctg ggttttaagt gatcctcctg  
cctcagcctc  
6541 ccgagtagct gggactacag gtgtgcaaca gcatgcccag ctgatttttt  
gtattttcag  
6601 tagagacgga gtttcacat gtggccagg atggtctcaa tctcttgacc  
ttgtgatccg  
6661 cccgcctcag cctcccaaag tgctagggag ttatatatgc atctctctt  
atctcttggc  
6721 tctctgcatg catctttctg tttctcttcc ttctttctt tttttttttt  
tttttttttt  
6781 tttttttttt ttttttgaga cggagtcttg ctctgtctcc caggctggag  
tgcagtgacc  
6841 agtctcggt cactgcaacc tccacctccc aggttcaagt gattctcgtg  
cctcagcctc  
6901 ccgagtagct gggattacag gcgcctgcca ccatgcctgg ctaatttttg  
tatatttagc  
6961 agagatgggg tttcaccatg ttggctgggc tggctctaaa ctctgacct  
caagcgatcc  
7021 gccggcctcg gcctccaaaa cactgggatt acaggcatga gccacgggtg  
ccggccagcc  
7081 tctctctcta cttggccctc ttctccttg tctccatttg tttctctgt  
gtgctatgac  
7141 tgtctgtctg tcaactgtctc ttgtctctat ctttgagagt cctaaatgtg  
gtccatttg  
7201 tcttttgaa aagctgcagg gaggactcag ggcagtgggg tgctgagtgt  
gttgagaca  
7261 gttgcagatc cttgacagtt ctcttccttg acagcgtgt ttccagtcac  
actgcagtgt

7321 gccaacatca gcatcctgga gaacaaactc tgtcactggg cataccctgg  
acacatctcg  
7381 gacagcatgc tctgtgcggg cctgtgggag gggggccgag gttcctgcc  
ggtgagacct  
7441 tactctgggg aaaatgaggc tgtcctgcc agttttctag gatttagggg  
agcagagggg  
7501 tcggccccc gcccttctgg gtcaaatga gaaggagact gggatacctg  
gttctctggga  
7561 gaggacggga ccagggcctg gactccttag tgtaaaagag aaaaggctc  
gaggtccaga  
7621 cttctggatc tacaggagga gtgggctggg cgtccagagt ctgagtcctc  
ggggaggagg  
7681 aggttaggtc ctgcggggag gtgggccctc tgagctttta ctctgggtc  
tgaggaagaa  
7741 gaggctggag atggaggact ctcgatgtt ggaggaggaa ggggctggg  
cctttctggg  
7801 agggaggaag tggcccggtg aattgtcatg aacagagtgg cctaacagtt  
cctctgccct  
7861 tctctcgct acagggtgac tctgggggcc ccctggtttg caatggaacc  
ttggcaggcg  
7921 tgggtgtctg ggggtgtgag ccctgtcca gacccggcg ccccgagtc  
tacaccagcg  
7981 tatgccacta ccttgactgg atccaagaaa tcatggagaa ctgagcccg  
gcgccacggg  
8041 ggcaccttg aagaccaaga gaggccgaag ggcacggggt agggggttct  
cgtagggctc  
8101 cagcctcaat ggttcccgcc ctggacctc agctgccctg actcccctc  
ggactactaag  
8161 actccgcccc tgaggctccg cccctcagc aggtcaagca agacacagtc  
gcgccccctc  
8221 ggaacggagc agggacacgc ccttcagagc ccgtctctat gacgtcaccc  
acagccatca  
8281 cctccttctt ggaacagcac agcctgtggc tccgccccaa ggaaccactt  
acacaaaata  
8341 gctccgcccc tcggaacttt gccagtgagg acttcccctc gggactccac  
cccttgtggc  
8401 cccgctcct tcaccagaga tctcgccct cgtagtgtca gggcgagtc  
agctccgccc  
8461 acgtggagct cgggcgggtg agagctcagc cccttgtggc cccgtcctgg  
gcgtgtgctg  
8521 ggtttgaatc ctggcggaga cctgggggga aattgagga gggctctggat  
accttttagag  
8581 ccaatgcaac ggatgatttt tcagtaaagc cgggaaacct ca

SEQ. ID. NO. 22

KLK-L3

VHFTPINHRGGPMEEEGDGMAYHKEALDAGCTFQDPACSSLTPLSLIPTPGH  
GWADTRAIGAEECRPNSQPWQAGLFHLTRLFCGATLISDRWLLTAAHCRKPL  
TSEACPSRYLWVRLGEHHLWKWEGPEQLFRVTDFFPHPGFNKDLSANDHND  
DIMLIRLPRQARLSPAVQPLNLSQTCVSPGMQCLISGWGAVSSPKALFPVTLQ  
CANISILENKLCHWAYYPGHISDSMLCAGLWEGGRGSCQGDSGGPLVCNGTLA  
GVVSGGAEPSCRPRRPAVYTSVCHYLDWIQEIMEN

SEQ. ID. NO. 23

KLK-L3

MKLGLLCALLSLLAGHWADTRAIGAEECRPNSQPWQAGLFHLTRLFCGAT  
LISDRWLLTAAHCRKPYLWVRLGEHHLWKWEGPEQLFRVTDFPHPGFNKD  
LSANDHNDDIMLIRLPRQARLSPAVQPLNLSQTCVSPGMQCLISGWGAVSSPK  
ALFPVTLQCANISILENKLCHWAYPGHISDSMLCAGLWEGGRGSCQGDSGGP  
LVCNGTLAGVVSGGAEPSCRPRPAVYTSVCHYLDWIQEIMEN

SEQ. ID. NO. 24

Table 13

CATGCAGTGTCTCATCTCAG

SEQ. ID. NO. 25

Table 13

CATGGAGGAGGAAGGAGATG

SEQ. ID. NO. 26

Table 13

CTTCGGCCTCTCTTGGTCTT

SEQ. ID. NO. 27

Table 14

GACCCTGACATTGGACATCTA

SEQ. ID. NO. 28

TABLE 14

GCCACTGCCTGATGGAGACTG

SEQ. ID. NO. 29

TABLE 14

AACATCAGCATCCTGGAGAA

SEQ. ID. NO. 30

TABLE 14

CTTCGGCCTCTCTTGGTCTT

SEQ. ID. NO. 31

TABLE 14

GGGTCAGAGCTGCAGAGAAG

SEQ. ID. NO. 32

TABLE 14

GGGCCTGTCGTCTGCAATGG

SEQ. ID. NO. 33

TABLE 14

ATGGCCACAGCAGGAAATCC

SEQ. ID. NO. 34

TABLE 14

GGTCACTTGTCTGCGCAGAC

SEQ. ID. NO. 35

TABLE 14

CCCAACCCTGTGTTTTTCTC

SEQ. ID. NO. 36

GGCCCTCCTCCCTCAGA

SEQ. ID. NO. 37

TABLE 14

ATCCCTCCATTCCCATCTTT

SEQ. ID. NO. 38

TABLE 14

CACATACAATTCTCTGGTTC

SEQ. ID. NO. 39

TABLE 14

AGTGACACTGTCTCAGAATT

SEQ. ID. NO. 40

TABLE 14

CCCCAATCTCACCAGTGCAC

SEQ. ID. NO. 41

TABLE 14

GCTTCCCTACCGCTGTGCT

SEQ. ID. NO. 42

TABLE 14

CACTCTGGCAAGGGTCCTG

SEQ. ID. NO. 43

KLK-L4 NA

1 caggaggttg cacactgttc ctcccacctc gccactgcac cccaccaag  
gatggaattg  
61 gaggcggggg gcagattcca gggtcagggc tgtcaagagt gaatgaggcg  
aggagacatt  
121 caggagcaga gaggtttcag acgcggaggt tccgggcacg ccctcaacac  
ccccttcacc  
181 ttctcctcag gccccgcccg ccctgccttc ccctcccgat cccggagcca  
tgtggcccct  
241 ggccctagtg atcgccctccc tgaccttggc cttgtcagga ggtaagaatg  
cgcgggggtg  
301 gaggcgcggc ggccattcgg gacaatggta ggaggggtca ggccggaggg  
ggagggggcg  
361 tgggagccgc gagctccgc ccccgccac tcggggccgg gtccagtggg  
gacagctcag  
421 agctcttcct gcttgtccct gggtgacctg gtttcccggc tgaggttggc  
cctccgacct  
481 cagacccttc acctccaaa atacctcgc agcagcccct cccgcgttca  
aggttctctg  
541 tcctctctgg aaagctgaaa gacatgggtt cgcgtcctga cgctgccgt  
ttgagccagt  
601 agcctagcag ctgctttgtg cctaaattgt tttcatctgg aaaatgggct  
taatctataa

661 gtgcttacca gagaagggtca ctgtgaatat tgaaacgagg taatgcgtcg  
agccttcagt  
721 atgtgcgagg tagaaggagac ttgaaagtta gccacttagc cgttattact  
ttattagtag  
781 tattcctttt tttttttttt tttttttttg agatggaacc ttgctctgtc  
tcccaggctg  
841 gaaggcagtg gcacgatctt ggcttactga aacctccgcc tcccgggttg  
aagcgattct  
901 cctgcctcag cctcccagat agctgggatt acaggcgccc gccaccacgc  
ccaactaatt  
961 tttgtatttt cagcagagac ggggtttcgc catgttggtt aggctggtct  
cgaacttcta  
1021 acttcaagta gcccgctca gcctcccaa gtgccaggat tacaggcatg  
agccaccgag  
1081 cccggcctct agtattctgt cttcatactc agccccttc agaaccttct  
agattgttat  
1141 tttaatcctt ggggtgaccc caaacctatg tgacctcacc ccaaattggt  
agtccttaag  
1201 atccttatgg atctttccca tctttccctg ccgttgtagg caggttctct  
ggaaaccccg  
1261 ttcattgaatc atttattcat tcaacaaaca gctattaaac accggccact  
gtgctgggtg  
1321 ctgtacaagc agagacacag tccctgctct cagcacctgg agtctagcgg  
ggacagacgc  
1381 agatgttatt caaacaatta tccaaataat tagttaataa ttatcttgac  
atgaggtgaa  
1441 gacttcaagg agccaagcca ggggcctaga gatgtaatgg cggcttcccg  
accagaggcc  
1501 ttcccaaagg gcttgaccct tgagccaaga cctgaaaaag gagggatctg  
tgggtgcctg  
1561 gcacctggca ccctccttgg cctgaagggtg ggggtggcttt tctcctctgg  
cgacactccc  
1621 tggattcatg cccgtgccac tctgagtgcc cacaccctag gctaggagac  
ccacacgcta  
1681 cgccttggtg agtcctcaac aacctggcga ggtaggtgca ttgtaattac  
tccaatttca  
1741 tggcagagaa acctaggact caaagacaga aggtcctgc tccaatgaca  
ccggcgatgc  
1801 ctgagtcaga atcctaatac aggttggttt ccctgtccat atcctggact  
tgaggctctg  
1861 aaaaccattt ttataacttt tgacctaatc atttgcttaa agttagcttt  
ttttcttctt  
1921 ttttactca aacaaaagca tgttcaactt tatattactg tctgaatag  
agaatagaat  
1981 tctttgtcat aaatagaagg taaggaagga aataaatcct gcacaatgaa  
aagaaaataa  
2041 tatgtttatt ggggtggacc acctgaaatt gctgatactt gacccttttt  
gaccttccta  
2101 aaacaacttt tgcagatggg tcagtgtaat aaatgttagg tggcctgatg  
aggcttctgt  
2161 gtcctcctgg ctttgaaaag tgagctcagt gaggattagg gaggtgttaa  
aaccatatta  
2221 gcaccatcct gagactttat ccttgacaaa atcaggttta aaagagaact  
ggatgctggg  
2281 tcagcgtctg agtgtgcgat ttaacgttac ttaaattctca tctctctacc  
atctaaaatg  
2341 atcctgtgct caccgacaac ttctgtccct aactgcaaac cactgagcta  
atccaactgc  
2401 ttgccctgta gttggggaaa ctagctaggg aggcagaggg acctcctgtt  
gtagctaata

2461 attaataata acatttccca ctgactgagt gctctccatg ccacctgctg  
tgctgcacgg  
2521 tttgaaatgc aggatcatct tgaattcttc aactgcgcaa tgagagatga  
actattactt  
2581 tttctacttg acagctgggg aaactgaggc tgggtgatttg cataagggtca  
cacagtcaca  
2641 aaatggcatg catgttcagg attggattct cctgtccca cggaccctg  
ctgtgctttc  
2701 aatgccagac acagtgcctg gcacacacag catttattta ttgagcccc  
attgtgtgcc  
2761 aggcgctgtg ttaggtcctg ggaatatggt actgaataaa gcagttaagg  
tgctgtttgt  
2821 caatggagct tacagtcaaa gtggagagat ttttaaaaac gaatacatc  
aaatgtgaag  
2881 agaaatgaat agcaatcatt gttctgatga agaccaactg gaagaatgta  
atgggggagg  
2941 agtcgggacc aggagagtca acattagacc aggtggtcag ggaaggcctt  
tctgaagagg  
3001 agacatttga gctgacctct cagaattaag aaggaccag acatacaacc  
tctaaattct  
3061 gagggtcatc cagtagaata ttccatata gtatatatga aatatcctat  
atctgtgctg  
3121 tccaattatc cactagcccc ttcaggctat tgaacatttg aaatatggct  
ggtgtgactt  
3181 aagaactgaa tttttaattt agttttactt cattttaatt agtttaatt  
taaataagcca  
3241 catgtagcta gtggctacca tattaacaa cataggctcag gagaaaggac  
tgtgcagaga  
3301 gaggaatag caagtataaa atgtctagta tgggggcac caagatgatt  
taaattcttc  
3361 ttttcttta atgcctgggtg tgtttgaaga acaggcccat gaggctggac  
tagaggaagt  
3421 cagaagaaag aggttggaaga tgggggtcaaa gaggctggca agggccagac  
agcacagagt  
3481 cctgcacacc ttgggaaggc tttttggatt ttattttaaa gaaagttgag  
cctgggaaca  
3541 acatctgact ttctttgttt gaagagtcct cagcctactt tgagaagact  
ggatcggagg  
3601 gatgtaaaag tggaaggatt taggttaatg ttgtagtcac ttgggctaca  
gaagatgggg  
3661 catggacca gatgggtggca gaagtgtgga gataactgga tatttgggag  
ataaaaccaa  
3721 taggaactgg ttgtgagtga tgaaggaaag aagagaagca aagatgactc  
ccaggtttgg  
3781 ggctgagcac tgagggtggga aatactggag cgaacagttt tgattgagaa  
gaatcaagtt  
3841 gggaatacaa agcttaagat gcctgtaagg catccaaatc aacagtgttt  
gagttttgag  
3901 cttaaagaag agttcagggc tggagatgat tagcctatag ctggtattta  
aagccatgga  
3961 ggcaaccagt atatatgcag tgaaaggata gagagatggg tggaaagatg  
attggatgga  
4021 tgcatggatg gatatatgga tagatggatg gatggatggt tggattggat  
ggatggatgg  
4081 atggatggat ggatggatgg atggatggat ggatgaataa atggaccagt  
ggatggaggg  
4141 acagatgagt ggatggatgg ttggatggat ggatggatgg atggatggat  
agatgggttag  
4201 atgactacct aaatggatga atggatagat ggatgagtag acggatggac  
aatcaatag

4261 gatgaatggg ggatggatga ttggatagat tgatggatag atattgccta  
ggatggatgtg  
4321 taggtcagtc tcacttctac ctccctgaaat ccatcttctg gtagaatgat  
ataaaaaatg  
4381 catgtggaga gaaagtcagg ctccctgctta cctatcagca acatcctcat  
tttgtgaact  
4441 cttctgttaa cccccagtg aggatttggg acttccctgag aaaataatgt  
caccctttg  
4501 ccctaattca tctccacttg gtcaagaata gcaactgcc a taggtcggca  
aattcatctt  
4561 cagttcctgg tcaccaggg caataatccg acccttacc caaaccaga  
aaccacaacc  
4621 ccagggctcc tctgccccct ggatcccagt tttctaaca tctctcttct  
ttaccaggtg  
4681 tctcccagga gtcttccaag gttctcaaca ccaatgggac cagtgggtt  
ctcccaggtg  
4741 gctacacctg cttccccac tctcagccct ggcaggctgc cctactagt  
caagggcggc  
4801 tactctgtgg gggagtcctg gtccacccca aatgggtcct cactgccgca  
cactgtctaa  
4861 aggagtatgt gggggccggg ggagcatggg gtagggatga gaatgggact  
gggattgtgg  
4921 atggggtaga gttggatttg aggatggagt tggagttagg gttggggatg  
gacatgggag  
4981 tgagaatgag gtttgggggt gagatatggg gattgggtat gggaatagaa  
tcaaagtagg  
5041 ggatttggat gggattgaag ttgaggatgg gggagatgta tttggagatg  
aggaaggtag  
5101 gatggagaag aagttagggt ggggatggga agagggtggg gctgggatgg  
ggatggaaat  
5161 gggtcatct tctttcctaa ccaccttct tctgcacca cagggggctc  
aaagtttacc  
5221 taggcaagca cgccctaggg cgtgtggaag ctggtgagca ggtgagggaa  
gttgtccact  
5281 ctatccccca ccctgaatac cggagaagcc ccaccacct gaaccacgac  
catgacatca  
5341 tgcttctgga gctgcagtc cgggtccagc tcacaggcta catccaaacc  
ctgccccctt  
5401 cccacaacaa ccgcctaacc cctggcacca cctgtcgggt gtctggctgg  
ggcaccacca  
5461 ccagccccca gggatgcac ccacacaggt ggcctgaggg cccataggag  
tggtcgggga  
5521 aacaggggca gagatgggag ggaaggctct aggtagggtc ctttatatat  
aaaaatataa  
5581 ataagtaaat aaatatatat atttaaagtt agctgtatcc tttatataaa  
tataaattca  
5641 tgaatatata aaaatatgag tatataaatt catgaatata tagaaatata  
aatagatcta  
5701 atatatgaat atattatatg atgtatatta tgtattatat agtaatataa  
ttatatatta  
5761 tacaaaaagt atacaaatta aatgtatttt ataaattata aaatttatca  
attatgtatt  
5821 ttaaatatgt atttctgcat aatgtatata ttatatataa tctatattta  
aattatatat  
5881 tataaatgta ttttataaat gtatacattt atatatttat atactgtaaa  
tgaattttat  
5941 catttataat atataaatca tacatataaa atgtttatat ttctataatt  
tataaaatgt  
6001 ttaatatatt aaatatgggt attaataaaa tgtctaataa ttcaatgtaa  
taattaattc

6061 tatatcatta cttagtaagt ataatacatt atatatgtga atataaagtt  
gatgtatata  
6121 ccgacaagag ccctttgcat ctccctagca atccctgact ctctcccagc  
ctcatgtttg  
6181 tatctttctc ctcaacatgc cctgtctctc ttcttaccat tctatccaac  
tctcccgtaa  
6241 ctcttcccat cctgttctct gcttttccca tctttaattc tctatttctg  
accatctccc  
6301 tattccaact cctctctctc aactttctct cccaccgct ggctccacca  
ctctccttat  
6361 caaccttcca ttctcttgtc ccttccctcc ttgtccttcc ctccactttt  
ctctcatct  
6421 ctcccttgcg ctctctccca tgtccctcca tatttctgtc acttccgttg  
ctttaccag  
6481 ataggtgtc atctcttctc ccatctttct ctcccatct caattttcta  
tctactcttt  
6541 acccattcaa ctgcctatt tcaccttcat cccatctct atccaggctg  
gataccttag  
6601 accttctctt tcttctcccc agtgaattac cccaaaactc tacaatgtgc  
caacatccaa  
6661 cttcgctcag atgaggagtg tcgtcaagtc taccaggaa agatcactga  
caacatgttg  
6721 tgtgccggca caaaagaggg tggcaaagac tcctgtgagg tgaggccggg  
aggctggtgg  
6781 gtgccttga caggatagaa agccagaatg gaagtgcag atgctgggga  
aaaagctttg  
6841 ttccagcct taggggaacc aatctttata agataaatg tcccctcaca  
taggaggta  
6901 agacaaaaag gggtagccag ggatggcagg aataattcat cataagcccc  
agctttgact  
6961 gagtggctgc caagatccct gtgttgagat gcataaagg tggattctt  
tcacttgtga  
7021 gtgatagaca accaactcaa actggcttaa acaaatgca ggcttttga  
actgaaaatc  
7081 caggttgtct ggctttaggc acagatggat ccaggatgc aaattgtgtg  
tttggaattc  
7141 tgtctttctt ttaactctca gctcttcttt attctgtttt ggcttcatte  
tcggttagat  
7201 tcttcccatg acaagatggc ccagcagct ttgagcttac atcctaccct  
ctaggcaacc  
7261 ctattagaaa gagaacctct cttttccaat agttcacaca aaagtcttaa  
gcatgattct  
7321 cactaggctg acctaatgca tgtgtcttga gccatcactc caccagagct  
gtgggattct  
7381 ctgatgggcc aagcctgagt cacatagtta actgtgggtg ctggagaggg  
gcagggacaa  
7441 actgcatgga ttggaagtgg agaagggcag ttccccaat gaaaaaatca  
ggagaggctg  
7501 ttaccaaaat aaggggaaat ggccaagtac agtagttcat gcctgtaac  
ccagcacttt  
7561 gggaggctga ggtgagagga ttacttgagc ccaggagttt gagaccagcc  
tgggcaacat  
7621 agtgagactc tgtctctaca aaaagaaaaa aaagttttta aattagccag  
gtgtggtgga  
7681 gtacaactgc agtcctagtt actcgggagg ctgaggcaga aggactattt  
gaaccagga  
7741 gttcaaggct gcagtgaggt atgatcatgc cactgcactc cagcctgggt  
gatagagcaa  
7801 ggcctgtct ctaaaacaaa aagaaataaa tagagcaaga cactgtctct  
aataaataaa

7861 taaataaaaaa tttaaaaatg aatgtttaat tttttaaaaaa taagaggaaa  
tggatactac  
7921 atgagcaaaaa aatagccttc atcaataaag aagttgagat tggattcagt  
gagaaagagt  
7981 atgatactat attaatgata tgtgccttga tcgattagtg atgtctgcct  
tgggccagg  
8041 aagagaaata gacttacacg tgtgttgcac accctgcca gatatgaatg  
ggttcaactca  
8101 atagttagag acacaaatga gccttaaata ggagcagggt cagctggtgt  
ggggcagggg  
8161 gtgatttagt accagggaaa caaaaatggg tatgaagtaa gttgttacca  
ttttaatgaa  
8221 actgaggaac agagaaaaac acagaaattt ctctgtgtct ctctttctct  
ggcctatct  
8281 ctgtctttct gtccctatct ctgtctcttg ctgtctgtcc ctctgtgtt  
gtcttcttgt  
8341 ctgtttctca ctgtcttcat tgctttctct cacactgtgt gtgtctgact  
ctgcctctct  
8401 gactctctct ctctgtgtgt gtctctctcc atctttcact ctctccccac  
acctccctgt  
8461 ccctgccttg tttagcccca gcaaggacc acctctctct ctctttcttt  
ccccaactca  
8521 ggggtgactct gggggccccc tggctctgtaa cagaacactg tatggcatcg  
tctcctgggg  
8581 agacttccca tgtgggcaac ctgaccggcc tgggtgtctac acccgtgtct  
caagatacgt  
8641 cctgtggatc cgtgaaacaa tccgaaaata tgaaaccag cagcaaaaat  
ggttgaaggg  
8701 cccacaataa aagttgagaa atgtaccggc ttccatcctg tcaccatgac  
ttctcactat  
8761 ggtctgctta gcccttctct gctccttatt ccagtggtc catttgaacc  
agtgatccat  
8821 gtcttgaaaa atgtcactc tcagctaaca ttccatgttt cagaagcatt  
caggcactgc  
8881 caggcttgca gtctcccaga tgttgcatcc ctgaaacatc tcaacaacct  
gaatgtccca  
8941 acccagacaa tggcccagggt ctctcaactt catcagtgtg gcttctatga  
gccagatca  
9001 ccacctgaac gttctgtctg tggcacattc ttaaataatt ccatcagccc  
atctcaacaa  
9061 tatatgtcct ataatggac catccttgac aacatcctct aactcttcaa  
gtatttatct  
9121 aatgccagta tcctagacct tctatTTTTT gactcaaga aggctctaga  
ctccatgat  
9181 agttcactct gaaaatatct tcttatgcc acaatcttct gccctgacaa  
cattctgtgt  
9241 acctctgtga ctaccacag ctaacattgg atcctcagaa tatttcattc  
tcacactgtt  
9301 atgggtgtct cagaagtccc aacccaacct acatcccaca ttcttccaat  
acccacctc  
9361 tgccaacatt ccctctctga atcaatggca ccctagtctc tagagttata  
gggttcagta  
9421 taccaaaggg tcttcttgcc tgaactttat tgtctacaa atattccgtc  
ttgtatcccc  
9481 tccatgaaca tccttgggtca gtgtcccttg ctgttacatc tttgtgcatg  
acctaaaaat  
9541 gtagtgcaaa tccttgcttt ggacaagtta taaaactcac agtctctgtg  
ctttctcatc  
9601 tgtaaaatgg gttcataatt ttttttaatt gtaacattat tacaagaata  
aatgtcaagc

9661 atttatcact attattatatt gcatggttcc cataaaatat taccttagaa  
tggttaataac  
9721 agcccttcga atttgagag tgtccaaaaa aagtgttgca ctgatttatt  
ttcctcagga  
9781 gacatttctt cagtgttgac tatgtgcaag cactctcctg ggtgttgta  
aatatagttt  
9841 atttactcaa caaatatttg tacctatcaa gagccaggca ctgttgaga  
gacaagtgat  
9901 aaccaatgag ttaaacagat aaaaacttct gccctttag aacttacatt  
ctttcaaga  
9961 agtctccata acaatgaata aagaaatagg ctgtcagggtg gtgctgcaag  
ccatagcaag  
10021 aaatgaaaca agggccatat gtggtagctc atgcctgtaa taccaacact  
gggaggccaa

SEQ.ID.NO. 44

KLK-L4 AA

ESSKVLNTNGTSGFLPGGYTCFPHSQPWQAALLVQGRLLCGGVLVHPKWVL  
TAAHCLKEGLKVYLKGKHALGRVEAGEQVREVVHSIPHEYYRRSPHLNHDH  
DIMLLELQSPVQLTGYIQLPLSHNNRLTPGTTCTRVSGWGTTTSPQVNYPKTL  
QCANIQLRSDEECRQVYPGKITDNMLCAGTKEGGKDSCEGDSGGPLVCNRTL  
YGIVSWGDFPCGQPDPRPGVYTRVSRVVLWIRETIRKYETQQQKWLKGPQ

SEQ.ID.NO. 45

KLK-L4 AA

MWPLALVIASLTALSGGVSSKVLNTNGTSGFLPGGYTCFPHSQPWQA  
LLVQGRLLCGGVLVHPKWVLTAHCLKEGLKVYLKGKHALGRVEAGEQVRE  
VVHSIPHEYYRRSPHLNHDHIMLLELQSPVQLTGYIQLPLSHNNRLTPGTT  
CTRVSGWGTTTSPQVNYPKTLQCANIQLRSDEECRQVYPGKITDNMLCAGTKE  
GGKDSCEGDSGGPLVCNRTLYGIVSWGDFPCGQPDPRPGVYTRVSRVVLWIRE  
TIRKYETQQQKWLKGPQ

SEQ.ID.NO. 46

Table 15

AACTCTACAATGTGCCACA

SEQ.ID.NO. 47

Table 15

TTATTGTGGGCCCTTCAACC

SEQ.ID.NO. 48

Table 15

GGATGGTCCATTTATAGGAC

SEQ.ID.NO. 49

Table 15

AGGCTGCCCTACTAGTGCAA

SEQ.ID.NO. 50

Table 15

ATATTGCCTAGGTGGATGTG

SEQ.ID.NO. 51

Table 15

AAGACTTCAAGGAGCCAAGC

SEQ.ID.NO. 52

Table 15

GACCCTTCACCTCCCAAAT

SEQ.ID.NO. 53

Table 15

CTAGTGATCGCCTCCCTGAC

SEQ.ID.NO. 54

Table 15

GGTGATCTGCGCCCTGGTCCT

SEQ.ID.NO. 55

Table 15

AGGTGTCCGGTGGAGGTGGCA

SEQ.ID.NO. 56

KLK-L5 na

1 attgagggg gatccagca ggtccattt gttcagattc ttctgggcct  
ttctgtgttg  
61 cttttcttcc tctgcaggac acctgtcaca tgagggtctt caggggaaaa  
ggaattcttag  
121 tgtctgtgac tgcttcaaag gagaagtaag gggagaggag aggaaagcag  
gaggaggttg  
181 gagagaacct cttgttcttg aggtcttcca atctccttca gctcaaagca  
ctcagcatgc  
241 tgaagttcca gactctgggc tatcacgttc tgacatgcaa cacaggcaac  
acccagctc  
301 catccacgtt tctccaaaag cacaggcatg acgtcatatg gtgacaaaac  
cccttgtcca  
361 aaggaagccc ataggatacg ctaattctag attcacaat actctagagg  
aactcacaca  
421 atgggatggg ccagtgcctc acacagagta tgaggcctcc caccttggtt  
gaatatcttt  
481 ttcttttttt tctttttttt tttttttgag acggagtctc actctgtcac  
ccaggctgga  
541 gtgcagtggc tcgatctcgg ctactgcaa cctctgcctc ccaggttcaa  
gtgattctcc  
601 tgccctagcc tctggagtag ctggaattac aggtgcccac caccacaact  
ggctaatttt  
661 tgtattttta gtagagacga gggccacca tggtgaccag gcttgtctcg  
aactcctggt  
721 ctcaagtgat ctgccacct cggcctagtg ctgggattac agacgtgagt  
caccacgccc  
781 ggccccacct tggttaaatt tctgaaaatc atttggtaaa gtgaggaccc  
ctccagctga  
841 gacactgcca ggaaacagct attgagtctc ttagcaccca cagcattaac  
acaaacccaa  
901 aacatttttag gcctcgttga gttctggagg caaatattc ctcatctaca  
aatttatctt  
961 attttatctt ttaataaaag ttattattat tttttttat agagagacag  
ggtcttgctc  
1021 tgtgaccag actagagtgc agtggtgtga ccatgggtca ctgcagcctc  
cacctcctgg  
1081 gctcaagcga tctcccacc tcagcctgcc aagtagctgg gaccacaggt  
atgcccctcc  
1141 ccaggctaa ctttttttta ctttttggg agatgaggtc tcaactatgtt  
gcccagctg  
1201 gtcttttttt tttttttttt ttttttgag acgggagtgc aggtggagt  
gcccaggtg  
1261 gagtgcagtg gcacaatcat aactcactgc agcctcaaac tctggggctc  
aagtgatect  
1321 ccacctcag cctcccacat aactaggact acaggcctgt gtcaccataa  
tgctgggtta  
1381 attttttttag tttttgtttg tagaaacagg gtctcactat gttaccaggt  
ctggcctcta  
1441 actcctggcc tcaatccatc ctcccctc aacctccaa agtgctagga  
ttatagccac  
1501 gagccacat gccgggccc atacttatat ttacttgtag tgagaacact  
taaaccctac  
1561 tcggtcagta attttcaagt acacaatata ttgttactaa ctatatacat  
ggtattttta  
1621 atttgcata atgtcctta aaatcagcga gccgcctta tttttgtat  
tcaattttat  
1681 tggatataaa ttccacatca gtaaacctga ttctcttaa aatctacaca  
gaaaaaaaaa  
1741 aagagagaga ggttgagttg ggtagttgtt ggtatttttg tttttgttc  
tttgcctatt

1801 cagttgaggt ttattgtaaa acttgacacc agaaaaagga agaagtggcg  
tttttggtgct  
1861 gtgagcgtgt gacgtgtgtc ctttccagag aaaggacagt catgggtgctt  
ttttatcctc  
1921 tctgccccaa gaaggaaagc tcttaacagc cagcaggagg ctttgtaggg  
accagcgtaa  
1981 tcaacgccag tcgcgctgac caatgatgag ttaaagcagt taggtcgttt  
ctaagagcaa  
2041 atcaaaagct aaggttctgt gattctgaaa atgagacacg gacagagact  
ggagacccag  
2101 agagaaagtg aaggactaaa agacagtcac aggggtgggag ttgtctctcc  
tctgttttgt  
2161 tctgggtttt ttgttttgtt tgtgcgctct gtctgaccgc ttttctttt  
tttcttttct  
2221 tttctttttt tttttttttt ttttgagatg gagtcttgct ctgccgcca  
ggctggagtg  
2281 cgggtggcaca atctcggtc actgcaagct ccgcctccca cgttcaaggg  
attctcctgc  
2341 ctcagcctcc cgagcagctg ggattacagg catgcaccac cacactcagc  
taatttttgt  
2401 attttttagta gagatggggt ttcaccatat tgaccaggct ggtcttgaac  
tctggcgctc  
2461 aagtgacttg ccgcctcgg cctcccaaag tgttgggatt acaggcgtga  
gccaccgcgc  
2521 ctggccctga ctgcttttct ccttggtttg ttgtcaatc ccccttctc  
tgagccgaat  
2581 tccctttttg ttctattttt ctctctctgt cccctctctc tctcttctt  
tctctcttc  
2641 cattctctct agatgaagca aaaactcaga taaaccagca cagaggccag  
gtatggtggc  
2701 tcacacctgt aatccccgga ctttgggaag ccaaggcagg caggttgctt  
gaggccagga  
2761 gttcaagacc agcctggccc acatggtgaa acccctctc tactaaaaat  
acaaaaatta  
2821 gccggacatg gtggcacgtg cctgtaatcc caggtactca agaggtggag  
gttgacagtga  
2881 gcggagatca cagccctgca ctccagcctg ggtgacagag cgagactcca  
tctcaaaaacg  
2941 aaaaacaaaa aacagcacia agttcccttg tctgtgact cattctctct  
ctctcttct  
3001 accatttctc ctccctgtg tctttttttt ttctctctgt gggttttatt  
taagcaatag  
3061 aagttcttag caaagaaaaa ctttatggaa ttagattgat ccacttcata  
tgtacatata  
3121 tgaactcagt tcagaaactc tcttctaccc ctgcctgac acctatttgg  
aagtctgttc  
3181 ctccaactct tcttctcttt ctgggactct ttctagcttg ggcttctgc  
ccctccgctc  
3241 cactctctg ctttcacagc ctctccttcc ccctgccct cccctgcact  
gcatgggat  
3301 gggccccagg tgtccaaggt ctccccaccc tcctttgtca ctggagtcag  
gattagaacc  
3361 cagctcccta gtcacctga gtcacagtc ctggggctgc tgacgggctt  
gcagaggaga  
3421 gagggagtgg ggctgggtct tcccacctg ggtccttctc tccttcccca  
ctccgtttag  
3481 ctgtaaagct caattaagtg tgattagctg agaagagttt ctgcagaatt  
agagcacgcc  
3541 ccaccctgt cttcgtgggc cccttccctt aaccgggaaa ctggatgggc  
caggacaaaag

3601 agagttaaga gctttgtcag tggctctgtct ggagcgacag atggaaggaa  
agggaccggt  
3661 tgagcaacat gacaggtggc tgaggagcca ggtgcagagt ggtagagttg  
gctggcggag  
3721 tggccagcac atgagaagac aggcaggtag gtggacggag agatagcagc  
gacgaggaca  
3781 ggccaaacag tgacagccac gtagaggatc tggcagacaa agagacaagg  
tgagaaggag  
3841 gtaggcgact gccaatgagg gagtgcacac caggggagca ggtagagaga  
ggacaagcag  
3901 gtcacccctt tggtgacctt caaagagaag cagagagggc agaggtgggg  
ggcacaggga  
3961 aagggtgacc tctgagattc cctttttccc ccagactttg gaagtgacc  
accatggggc  
4021 tcagcatctt tttgtcctg tgtgttctg gtgagttctc ccggagcagg  
gagagggcag  
4081 gactgcgact ggatccctc acccccatga ggaggcccca ccaccctccc  
catctcagct  
4141 ctggcccccga gcctgggtgt gaggaggaga ggggctttct ctgtgcctcc  
atttacctgc  
4201 agctctcagg gtactgtc cctcggctc cctattttt tgatccctct  
tcccttctgt  
4261 cctctctga atctctgtct ctccatttcc ctctatgtg taagcatctt  
tctccttggg  
4321 tgtctttgat gtttcatggt ctttttctat cactgggtct ctctctctt  
ctctctctt  
4381 ctgctctctc tttctcctct ctctctcctg cctgtttctc tctctactc  
tgtgtgtctc  
4441 tccatctctg tatcttttct tctctctct gaccatgcc cctgtctgtc  
tccagggctc  
4501 agccaggcag ccacaccgaa gattttcaat ggcactgagt gtgggcgtaa  
ctcacagccg  
4561 tggcagggtg ggctgtttga gggcaccagc ctgcgctgcg ggggtgtcct  
tattgaccac  
4621 aggtgggtcc tcacagcggc tcaactgcagc ggcaggtaa tcccttccgt  
gggtgggcga  
4681 agggaggact atgggaaggc aagcgctggg ggtaggatca caaggaggag  
tggtgcccac  
4741 tggaagaag ctgacctctc aacaagagag tctgaggtta gaccaggagt  
ggaacttctc  
4801 tagcagtggg cctgggggtg tgctgggcag ggtgaggtat gttgggtgga  
gggccgggga  
4861 gggctcctgga acctgccctc ctgcctctcc cattcctgca tgtacctt  
cttctctata  
4921 tgacatctgc cactcacccc agccattcct tgaccagtc tgggcccg  
gccaggtct  
4981 caccgaagct ctttttctt tctttttt tatttttttg agacagggtc  
tcgctctgtc  
5041 gccaggtctg ctgtgcaatg gcgtgatcac agctcactgc tgtctctgcc  
tcccaggtc  
5101 aagtgattct cctgccccag cctcctgagt agctgggatt acaggcaccc  
gccaccatgc  
5161 ccagctaatt tttgtatttt ttgtagagac agggttttgc catgttgcc  
aggctgggtc  
5221 cgaactcctg gcctcaaagt acctgcccgt cttggcctcc caaagtgtg  
ggattacagg  
5281 tgtgagccac tgcacccggc caacatgacc caaactctt gtgcaactc  
agaatctatg  
5341 cctggcacct ctctgggct cagtagactg atgttctgga attttttct  
ttttcttct

5401 tttttttttt ttttgagac agagtcttgc tctttctgtc atccaagctg  
gagtgcagtg  
5461 atgctatctt ggctcactac agcctcaacc acctgggctc aagtgatcct  
cacacctcag  
5521 cctcccaagg agctaagact acaggcctgc gccaccacac ctggctaatt  
tttaaatttt  
5581 ttttgtagag acagggtttt gctatgttac ccaggctggg ctcaaactcc  
tcagctcaag  
5641 caatcttcct gccttgacct cccaaagtgc tgggattaca ggcatgagcc  
actgtgcctg  
5701 gcctggaact ttttttgtga aaggggagat cagatgcaaa gaaacagaga  
ctcagggaga  
5761 gagagggcca gcagcaggat gcagagaggc cattcatcaa cccactcggt  
caatcatgaa  
5821 cccactcgtc cagcatgag catggagggc acatgctccg tgccaggcgg  
tgggaataag  
5881 gcagtgaaca aggtccactg atgtccctgc cttcatgggc ttcaccagcc  
gagagaatca  
5941 gaaagagagg cctggcgcggtg tggtcacac ctgtaatccc agcactttgg  
gaggccgagg  
6001 cgggcggtatc acttgagggtc aggagtttga gaccagcctg acacacatgg  
tgaaacctta  
6061 tctctactaa aaatacaaaa attagctggg catggtggca tgcttctgta  
atcccagcta  
6121 cttgggaggc tgaggcagggt gaattgcttg aacctgggag gtggagggtg  
tagtgagcca  
6181 agatggtgcc actgcactcc agcctgggag acagagcgag actcgggtctt  
gaaaaaaaaa  
6241 aaaaaaaaaa aaaggagaga gagagacaca gatgcaggga catggtagga  
gaaacaggga  
6301 acacccaaga tggaaagagg gtgatggagg ttgggaataa gagcctgtaa  
gagagactcg  
6361 gagaatgaga gttgcgggtg agaggacaga cagtgagggg cagaacagtg  
gggagcggca  
6421 ggagcgcctg agtgtccgtg gaggggtgca aggtggggga ctgcgtgcct  
gccaccgct  
6481 cagccgtcgc caccggcagc aggtactggg tgcgcctggg ggaacacagc  
ctcagccagc  
6541 tcgactggac cgagcagatc cggcacagcg gcttctctgt gacctatccc  
ggctacctgg  
6601 gagcctcgac gagccacgag cagcactcc ggctgctgcg gctgcgcctg  
cccgtccgag  
6661 taaccagcag cgttcaacct ctgcccctgc ccaatgactg tgcaaccgct  
ggcaccgagt  
6721 gccacgtctc aggttggggc atcaccaacc acccagggag taagggggcc  
agggccaggg  
6781 gtcaggggtc aggatgggta caagtctggg atgcagggcg agaggtcgaa  
tcatgacacc  
6841 tcagaggaag gatgggtaaa gggtcagggt gtgggatggg acatcaggat  
catggtttgg  
6901 ggtcagagat tatggtggat tggggctctt ggagccaaag ggggttaaagg  
actgggtatg  
6961 aagtcaggga tcagaggtca gaggtcagag tgtgtcagag gtcacacac  
tgagcaaaa  
7021 ggcatatata tatatatatg tatgtatagg atatgggcat tgtgggtcat  
gggtctgggg  
7081 ttagaggtca ccgtagaatt aaggtcatgg gatccagagg ttgtacaatc  
tggtcaaaat  
7141 ctgaggatgg aaattgggat tctatccaaa atcacatatc tgagattgga  
ggtcatagcg

7201 tttggggtgt ggggcccga gtttggggtc atggaggctg gggcccaata  
aactaggatc  
7261 aggggacact ggcgttgga gcaagtgggt ttggaagatg cagagctgag  
gttggagggt  
7321 aaggtaaaga caggacatg gggtcaggag acagaagata tgagatcaag  
ctgggatcat  
7381 aaggtaataa gacagaaggt caaagatcac agtagctggc attgaagagg  
gtcagggtctg  
7441 gattcgttgt ctctgacgct ggagagacaa gaaagttctt gagttatgcc  
actcaaagtc  
7501 aaatgtcaaa gatcaaagag accgtcaatc atctggggtc atgattcata  
tgaaattaag  
7561 tcataaatat gtaacttggg ggtttcggga ttgtagtaca ggtcggtagg  
gggcaggggt  
7621 attgacatgg atggggccaca tccagggaag agggacgtgg cctcaaagtg  
gggagattta  
7681 ggggaccctg cagcacgcat gttctctctc cagaccatt cccggatctg  
ctccagtgcc  
7741 tcaacctctc catcgctctc catgccacct gccatggtgt gtatcccggtg  
agaatcacga  
7801 gcaacatggt gtgtgcaggc ggcgtcccgg ggcaggatgc ctgccagggtg  
agccagtga  
7861 ggcagcgtgc gtggtcacca ggacaggaag tgaaggggag gggctggaag  
caggagggga  
7921 actgatggag gatgaatcag ggaaagggga tgctgcagag agacggggtc  
aaaaaggaag  
7981 ggagaggctg gttacggagg ctcacacctg taatcccagc actttgggag  
gccgaggcgg  
8041 gcggatcact tgaggtcagg agttcaagac aagcctggcc aacacgggtga  
gactctgaat  
8101 ctactaaaaa taccagaatt agccgggggt ggtggtgcaa gcctgtggcc  
ccagctactt  
8161 ggaaggctga ggcaggagaa tcgcttgatc ccgggaggcg gaggttgag  
tgagctgaga  
8221 tcacgccact gcactccagc ctgggcgaca gagccagact ctgtctcaaa  
acaaaataat  
8281 taataataat aataataata ataataataa taatggagga gaggccag  
ataagggagg  
8341 gagagagaca gggagtaaaa gggaggaccg gggaatggag gagggggagg  
ggcagggaga  
8401 gagaggagg aagggaacag agaaggaaag atggggcagg ggttacagag  
agagacagca  
8461 aaacagacgg agaggactgg gagcccagac agggaaccag ctgtttctg  
ggctctaagt  
8521 ctttcccata ccactctca gttggtgctg tcccagactg agagagattt  
gaggatggcg  
8581 gtctctcccc tcattggtca gggccccagc cattgtcctt gagagaacte  
tgtgcttttg  
8641 atggagtcct gccaccttc cctgggattg gtcatttttg atggcactct  
ctcccccat  
8701 tggtcagaac cccaggcatt gtccttgaga gaacctctat cctttatgga  
gtcccacct  
8761 cctcccctgg gattggtcat tgataatagt gttctctctc ctcatgggtc  
agggccccag  
8821 ccattgtcct tgagagaatg ctgactctt tatgttgtct tgacagcctc  
ccctgagatt  
8881 ggtcattaat gactgtgctc tctctctca ttggtcaggg cccagccat  
tgtccttgag  
8941 agaacctctg tcctttatgg agttccacc ttcttccctg ggattggccc  
ctagagacag

9001 tgggttcttct ctttttggtta gccattgcca ttgtcctccg ggaaagtgat  
tatactcttt  
9061 tgtctaataga ccagacttgg agccctcccc aaggcccagg actggggtga  
agggttgggg  
9121 aggaaaacag aaataagatg tctcccttgt tcagacagta cttctcttcc  
cttccagggt  
9181 gattctgggg gccccctggt gtgtggggga gtccttcaag gtctggtgtc  
ctgggggtct  
9241 gtggggccct gtggacaaga tggcatccct ggagtctaca cctatatattg  
caagtatgtg  
9301 gactggatcc ggatgatcat gaggaacaac tgacctgttt cctccacctc  
cacccccacc  
9361 ccttaacttg ggtaccctc tggccctcag agcaccaata tctcctccat  
cacttcccct  
9421 agtccactc ttgttggcct gggaacttct tggaacttta actcctgcc  
gcccttctaa  
9481 gaccacagag cggggtgaga gaagtgtgca atagtctgga ataatataa  
atgaaggagg  
9541 ggccatgtct gtccatttga agtcctcatg ctgggtgaga ctggaagaag  
gactcagcag  
9601 tttccctatc tcataggagt agaaacagag ctcaaataag gccaggcaca  
gtggctcaca  
9661 cctgtaatcc catcactttg ggaagctgag gcagggtgat cacctgaggt  
caggaactcg  
9721 ggaccagcct ggtcaacata gtgaaacccc aactctacta aaaatgcaaa  
aattagccag  
9781 gcatggtggc gcatgcctgt aatcccagct actcaggagg ctgagacagg  
agaatagcat  
9841 gaaccctgta ggcagaggct gcagcgagcc gagattgaac cattacactc  
cagcctgggc  
9901 gacagagcga gactccatct caaaaacaaa caaacaaaaa acccagtgtc  
caaataggat  
9961 gagggctctc cctgagtagt tactcagaaa tggagtagaa aaagttactt  
ttaataatat  
10021 aggccgggtg cagtggccca cgcctgtaat cccagcactt tgggaggccg  
agggtggagg  
10081 atggcttgag ctcagatttc gagatcagcc tggcaacaca gtgaaatctt  
gtcactacaa  
10141 aaacacaaaa aattagctgg gtgtggtggt gcgtgcctgt agtccagct  
acttggaag  
10201 ctgagggtgg aggatcacc gagccgggga ggtggaggct gcaaagagcc  
gagatcatgc  
10261 cactgcactc cagcctgggc aataaagtga gaccttgtct caaaaacaaa  
aaccagcaa  
10321 tataaataag acacatgttt cttcatctgg cataatagaa atagtgccca  
gagcttataa  
10381 gcttttcaag agtcacaaaa agaccgaaa aagaaaaaga aaattgttag  
ctccaaaata  
10441 ccagatgaaa gctgcaaagt caacatttat gaccatttaa tccaatgtcc  
ataaaacgta  
10501 gcattctttc cactagccaa ctgcagtta ctttcttgta atgaagcata  
cattgtatct  
10561 ttaatgtggg acgtggcttt gttctaataa gacgaagggt ggagtgcagg  
cttggaagc  
10621 aggagagctc agcctacgtc tttaatcctc ctgccaccc cttggattct  
gtctccactg  
10681 ggactcaaga ggtgaggaga gaccatctcc ccaaagcac tgaagggaaa  
ctggaggagg  
10741 gagggagtga ggggtgatca taccagcgga ggcacatttg ctgagcccc  
ccgcagtctg

10801 ctcttttccaa gtggaccctc ctggaagcct gatcccaacc tcccctgcaa  
 gcaggtctgt  
 10861 cacccccac tctcagatga agaaactgag ccttgaggg gtggagtccc  
 ttgtcccccac  
 10921 gtcataagg tagtcatagt agtaggaaga ggaagcacct aggtttgagg  
 ccagggtgg  
 10981 ctgctgtcag aacctaggcc ctcccctgcc ttgctccaca cctggtcagg  
 ggagagaggg  
 11041 gaggaagcc aagggaagg acctaactga aaacaaacaa gctgggagaa  
 gcaggaatct  
 11101 gcgctcgggt tccgcagatg cagaggttga ggtggctgcg ggactggaag  
 tcatcgggca  
 11161 gaggtctcac agcagccagt aagtgaacag ctggactcgg gctgcctggg  
 cggcagggag  
 11221 aagcgggcag ggggaagggtc agcagaggag cgaggcccca gaggagccct  
 ggggtggagc  
 11281 acagccaagg gctctgttcc ctttcttga ctcggcttcc acaggccctg  
 acctgcctcc  
 11341 cccaccctcc ggtcctgccc ctgtgcctgg cagcagcccc acctgtgtga  
 catcccagca  
 11401 cccccccct ctccttgcaa aggagaaggg agcggcctag gggaggccag  
 gggccaccc  
 11461 gggctggggc tgtggagagg gagggtgg gacgggagga aaaagagaga  
 cggagattag  
 11521 atggaagaag agggatttca agacaaattg ccagagatgc agtcagagag  
 actgactgag  
 11581 agacacaaag atagaaggaa ttagagaaag ggccacacag agccagacag  
 agagagaaga  
 11641 gtggagatgg agacaggac gaggacagag aaaggcagac agacacatag  
 ggacagaaag  
 11701 agaaaaatca cacaagtca gaattactga atgacaggga atgacacata  
 gaacgagaca  
 11761 cagattcaga gactcagggc agggaaagga aggctgcaga cagacagaca  
 gacagaggga

SEQ.ID.NO. 57

KLK-L5 AA

LSQAATPKIFNGTECGRNSQPWQVGLFEGTSLRCGGVLIDHRWVLTAAHCSG  
 SRYWVRLGEHSLSQLDWTEQIRHSGFSVTHPGYLGASTSHEHDLRLRLRLP  
 VRVTSSVQPLPLPNDCATAGTECHVSGWGITNHPNPFDPDLLQCLNLSIVSHA  
 TCHGVYPGRITSNMVCAGGVPGQDACQ

SEQ.ID.NO. 58

KLK-L5 AA – alternatively spliced

MGLSIFLLLCVLGLSQAATPKIFNGTECGRNSQPWQVGLFEGTSLRCGGVLID  
 HRWVLTAAHCSGRPIPGSAPVPQPLHRLPCHLPWCVSRENHEQHGVCRRRPG  
 AGCLPG

SEQ.ID.NO. 59

KLK-L5 AA – alternatively spliced

MGLSIFLLLCVLGLSQAATPKIFNGTECGRNSQPWQVGLFEGTSLRCGGVLID  
HRWVLTAAHCSGSRYWVRLGEHSLSQLDWTEQIRHSGFSVTHPGYLGASTS  
HEHDLRLLRLRLPVRVTSSVQPLPLPNDCATAGTECHVSGWGITNHPNPPFD  
LLQCLNLSIVSHATCHGVYPGRITSNMVCAGGVPGQDACQGDSGGPLVCGG  
VLQGLVSWGSVGPCGQDGIPGVYTYICNSTLVGLGTSWNFNSCQPF

SEQ.ID.NO. 60

KLK-L5-AA

MGLSIFLLLCVLGLSQAATPKIFNGTECGRNSQPWQVGLFEGTSLRCGGVLID  
HRWVLTAAHCSGSRYWVRLGEHSLSQLDWTEQIRHSGFSVTHPGYLGASTS  
HEHDLRLLRLRLPVRVTSSVQPLPLPNDCATAGTECHVSGWGITNHPNPPFD  
LLQCLNLSIVSHATCHGVYPGRITSNMVCAGGVPGQDACQGDSGGPLVCGG  
VLQGLVSWGSVGPCGQDGIPGVYTYICKYVDWIRMIMRNN

SEQ.ID.NO. 61

Table 17

TCAGCCAGGCAGCCACACCG

SEQ.ID.NO. 62

Table 17

TTGGTGATGCCCCAGCCTGA

SEQ.ID.NO. 63

Table 17

CCACACCGAAGATTTTCAAT

SEQ.ID.NO. 64

Table 17

GCCCCTCCTTCATTTATA

SEQ.ID.NO. 65

## KLK-L6 NA

1 atcgtgtaat caccgccaca tccagtgcaa agctgattcg tcaccacaga  
gcagctccct  
61 cctgccaccc catccctggg tcccaagaga accctttctt aaaagaggga  
gttcttgacg  
121 ggtgtggtgg ctcatgcctg taatccttgc actttgggag gccaaaggag  
gtggatcatt  
181 tgagggtcagg agtttgagac cagactggcc aacatggtga aaccctgtct  
ttactaaaaa  
241 tacaaaaaaa tgagcggggc atggtggtgg gtgcctatag cccagctac  
tcaggaggct  
301 gaggcaggag aatcgcttga acccaggagg cagaggttgc agtgagccga  
gattgagcca  
361 ctgcactcca gccggggcta aagagtgaga ctctgtctca aaaaaaaaaa  
aaagaaaaag  
421 aaaaaaagaa aaaaaataa aataaataa taaataaaat aaatttaaaa  
atttaaaaat  
481 aaagaggggg ttcttgtgtt gatgccgagc ctgaaccaag gcagaggagg  
ccgggaaggc  
541 ttcccaaggc cttcagctca aagcaggagg gcccatagtt aaacagaaac  
agttcaggaa  
601 tcacagaaag gcacctgggg agagatgggt gtgtggctcc agatgcagg  
gccagacag  
661 tgcgtcccca ggtgtacaga cagaccagg ccaagctcca gctcaaagag  
ccagcctagg  
721 ggggtgccga ggtggaggga ggctgagtca ggctgaggcc ggggaacagt  
tggggtagcc  
781 aaggagggca agcagcctcc tgagtcacca cgtgggtccag gtacggggct  
gccaggcccc  
841 agagacggac acaagcactg gggaatttaa ggggctaggg gaggggctga  
ggagggtagg  
901 cctccccca aatgaggatg gaaccccccc aactccagaa cccccctgca  
ggctggccag  
961 aatccttccc catctcatte actctgtctc tctgtctctc tgccgtctcc  
tattttgaat  
1021 ttccaacccc gtctgttaag actgtccttc tgtctctgaa tctctgtccc  
cttctcttcc  
1081 tgggtctctc tccctctccc tctgggtctc tgtcccccctc tctgggtctc  
tgtcactctc  
1141 tctttgcac tccagctctc actttgtctc tgcacctagc agatcccaag  
ctggggaatg  
1201 ccagttcttg caccaacctt cctgtctcct gctggggcct ctgctcccc  
atctctcagg  
1261 agtcgaaagt gagaaagcaa ggtgggcagc tctgtctcag gtccaggat  
ctcccgccca  
1321 cctcctgccc gtctctatc ccacccctcc tctccatctc tccctgggcg  
tgccatctct  
1381 catctaggcc tccgtctcct ctgtcattgt ccccatcccc tgtagggtgc  
catecttccc  
1441 gtctccccctc tgccatcggc ctgcctgtcc catcctcttt ctcccaccat  
gtcccgttct  
1501 ctccacgtc tcatgcccgc actgccttca tcatcatcgc tgtgtttctg  
tgtgtgtttg  
1561 tgggtgagtgc cgcagtgtgg gggcgtctcg gcctctctcc tctctctcca  
ctgttttctc  
1621 tttctgtgtg tctgtttcca ttctatctcc acctcttccc ctccgtcttt  
tgcttttcta  
1681 tctccacttc tccacacccc tctctccctg cgtctctgtg tctccctctt  
cctctgtctt

1741 gtttttttcc caccgtctgc ctcttctgtt ccctgtcaca tccaacttcc  
accggtttct  
1801 ccagctctct cctcagttcc ttctctcatg agcacacctg cctctgtgct  
cgtattcctg  
1861 gactcctctc tctccactgt catatcttct cattcatttt cccagtctct  
ctctgtctct  
1921 tgctctcccc ctctctgtca ctctgtctct gtctctctct ttctctctct  
ctctctgtgt  
1981 ctctctgtct ggtctctctct ctgtctctct ctccatctct ctctctctct  
ccccccgtc  
2041 accctgtctc tgtctctctc tgtctgtgtg tctctctgtc tttctctctc  
tccatctctc  
2101 tctgtctctc tctctctctc tctctctctc cctctctccc tctctccgtg  
actcctctc  
2161 tcagtcctac tcttctctcc tctctcagcc cttctgtgcc ctttctctg  
aactcccca  
2221 ccttggttct ctgactccac cactagatcc accacctcca gcaactggga  
accctcccct  
2281 gccacacctg ccttggggtc cctctccagg attccttcta gattatagca  
tcttccctgg  
2341 gcgggttctc atgaacaatt gtggctgctt ttttggccag acaggggagg  
gaggggatgg  
2401 gatcagggag tcttggaatg ggaactaggc aataaaaaaa aaaaaatgtc  
agaagcaggg  
2461 cggcgggagg tgggggcagg gccagctgtc cttaccaggg ataaaaggct  
ttgccagtgt  
2521 gactaggaag agagacacct cccctccttc cttcatcaag acatcaagga  
gggacctgtg  
2581 ccctgtctca catctctcca cctgccgccc gcagagcctg caggccccgc  
ccccctcgtc  
2641 tctggtccct acctctctgc tgtgtcttca tgtccctgag ggtcttgggc  
tctgggtaag  
2701 tgccccctgc tgtctctgcc tctcagcccc cggttctgtt gaaggttcct  
tctctctcac  
2761 tttttctctg catttgacag gacctggccc tcagccccta aaatgttctc  
cctgctgaca  
2821 gcacttcaag tcttggtat aggttaagaga acggttgggt atgacacaag  
ggggtcccct  
2881 ggagactctg agaagagatg gggatgggtc cttggggccc ctggatgctc  
atggtgacct  
2941 cataagaaag agcagggagt gggttggggg tcatgggtggg ggaacgtgct  
ggaggcctaa  
3001 attcctagtt gtggaggtgc tagggaattg tggggccggg gagagaggtg  
tttataaggt  
3061 ctggtgcaaa atacataagg aatcttaggg aactattagg tcttgagtgg  
gtcatagcag  
3121 aaagatcacg gggctctacc tgactgtgtt aggaagaaa caatgtcaga  
aagatgtttt  
3181 gttgtcagag ggaaggtgga gaaggatgat gggatggcgg gatcgtggca  
tggggtggcg  
3241 ggatcgtggc atgggtgtgt gaggtggatg ggggcaagtg tggggcaaga  
gatggcggat  
3301 ccttggggtc cactgagtg ggaacgttgg ggaggagaca gggaggtcct  
tgaatgtgtt  
3361 ggggaaggac tcattggggg gaaatgtggc atatttcgag aagtgtcac  
agaaattatg  
3421 ggagcataga gctaagggtc gtagatgtag caaggccctg gataaggtgg  
ccacggcaca  
3481 aaataagaga tgctacggag gtgacttggg aggtgagtca gaaagctctc  
cgtgctgggg

3541 caataacggg gtcaatattg ggcattgtctc accctgggtg ggacagatag  
aggcgggcag  
3601 tttaggggtt agacaaaaag gaaggggatt tgtcagtttt ggaatcctac  
aaacttgttg  
3661 agtggagagt gtttgtcat ctactttccc cacccaatcc tgtccactcc  
tagccatgac  
3721 acagagccaa gaggatgaga acaagataat tgggtggccat acgtgcaccc  
ggagctccca  
3781 gccgtggcag gcgccctgc tggcgggtcc caggcgccgc ttcctctgcg  
gaggcgccct  
3841 gctttcaggc cagtgggtca tcaactgctgc tcaactgggc cgcccgtaa  
tgacccctc  
3901 cctgtccct gtacctagtg aattccagag tctaaagccc tagagctgag  
ctgagaacct  
3961 ggatctctgt atagaacca atgtagtggc tggctcctgg tttgaggtct  
agagaagagc  
4021 ctggaacaaa aacacagctc gggatgtggg ctctccata aatctcgac  
tcagcatagg  
4081 ttctgaaagc agatgggcag cttggaacct atggacctgc tgagaaccga  
acatctgatc  
4141 cagtgttct tccagaggcc acacattaca tcgagaccaa gcttagccca  
ttccagattg  
4201 gtggctgaat tcaggacccc gtctacattc agaaactcag gacactacgt  
agaactcaga  
4261 gccagttca ggacctgcag tctagccata aatccagaac tagaacgtg  
ctcacagctg  
4321 gaacatacaa ctctaagaat agaggcaaaa cctggaggct gtttcacacc  
caaggtttag  
4381 ttcagagtct agtctatagc tccgctatga gcagacttca acccagtgtt  
tgaatcccag  
4441 aatgtggcgg gtgcgggtgc tcatgcctat aatcctagca ctttgggatg  
ctgaggcagg  
4501 cagatcacct gaggtcagga gtctgagacc agcctgagca acatagagaa  
accctgtctc  
4561 tactaaaaat gcaaaattag ccaggcatgg tggcacatgc ctgtaatccc  
agccactcg  
4621 gaggtgagg caggagaatc acttgaacct gggaggcgga ggttgcagtg  
agtcaagatc  
4681 gcaccattgc actccaggct aggcaacaag agcgaaactc catatcaatc  
aatcaatcaa  
4741 taaatcccag aatgcagatc ctaatcagaa gccccatata aaacctagac  
ccctcctaaa  
4801 ttctagatct gaacttcaa ccagacccc agccaagagg tcaaatgcc  
tataagccat  
4861 atctatgcca taaacaggtc agtctagaac ctagagatca aagctcaggc  
cagagtctag  
4921 aatataaagg ccagaatgca aaccagactc tagaatcttg gatccgggcc  
ataacctaga  
4981 gctccaacta gaacccagag cccaacctga ggtcaagggc tagggccaga  
gtccagaacc  
5041 aagagcccta taatccaata tgaaacagac ctgtagaggc tgggtgcggt  
ggctcacgcc  
5101 tgtaatccca gcactttggg aggtgaggc gggagaatca cttgaactgg  
gagttggagg  
5161 tcgagagtga gctgagatcg tgccactgca ctccagccta ggtgacagag  
cgagactcca  
5221 tcacaaaaaa aaaataaata aataaatcaa gtcataatcc aggttcgatc  
tagaatcctg  
5281 atcttagcat agagtcaaaa gttaagatg tctagaactc agaaccagg  
ctagaacag

5341 aatggtgcct actccggaat atcagttccg atttagagcc tagactcata  
acgcagtttc  
5401 gcttaggact caatgcaccg agcccagcac agaccctggc acggagccaa  
gctctcccaa  
5461 tcatcacctt cttcccaagc caggagctgg agcccagccc aagagcggaa  
ggagaggcag  
5521 ctggggctgg gccgagagaa tgccctggcc atggggaagg gcacaggagg  
ccaagaatgc  
5581 tcggcctgca gttagtgaga agcaggctag acctcgggga agactcgtca  
cccggccagg  
5641 gaaccgggct ggagggtggg gaggagtctc tggctcagac cctgagcagc  
gcttctcttg  
5701 ggggtcgtgg ccaggatcct tcaggttgcc ctgggcaagc acaacctgag  
gaggtgggag  
5761 gccaccacgc aggtgctgcg cgtggttcgt cagggtgacgc accccaacta  
caactcccgg  
5821 acccacgaca acgacctcat gctgctgcag ctacagcagc ccgcacggat  
cgggagggca  
5881 gtcaggccca ttgaggtcac ccaggcctgt gccagccccg ggacctcctg  
ccgagtgtca  
5941 ggctggggaa ctatatccag ccccatcggg gaggactcct gcgtcttgga  
aagcagggga  
6001 ctgggcctgg gctcctgggt ctccaggagg tggagctggg gggactgggg  
ctcctgggtc  
6061 tgaggaggga ggggctgggc ctggactcct gggctcaggg gaggaggggg  
ctgaggcctg  
6121 gactcctggg tctcaaggag gaggagctgg gcctggactc atacgtctga  
gggaggaggg  
6181 gctggagcct ggactcctgg gtctcaagga ggaggggctg ggcctggact  
tctgggtctg  
6241 agggaggagg ggctggggac ctggactccc gggctcaggg gaggagggac  
tgggggtctg  
6301 gactcctggg tctgaggag gaggggctgg gggcctggac tcctgggtct  
gagggaggag  
6361 gtgctggggc tggactcctg ggtcggaagg aggaggggct gggggcctgg  
acccttgggt  
6421 cttatgggag ggtagacca gttataacce tgcagtgtcc cccagccagg  
taccgccct  
6481 ctctgcaatg cgtgaacatc aacatctccc cgatgaggt gtgccagaag  
gcctatccta  
6541 gaaccatcac gcctggcatg gtctgtgcag gagttcccca gggcgggaag  
gactcttgct  
6601 aggtgaaggc caggatggga gctgtggtag ggattatttg ggactgggat  
ttaagcaaat  
6661 gatgtcagga gcatggaagt ctgcagaggt cttcagaaga gagtgaaccg  
caggcacaga  
6721 gagattccga tagccaggcc accctgcttc ctagccctgt gcccctggg  
taatggactc  
6781 agagcattca tgccctcagtt tcctcatctg tcagggtggga gtaaccctct  
taggtagtt  
6841 ggtggaatgg gatgaggcag gttggggaaa gatcgagag tggcctctgc  
tcatatgggt  
6901 ctgggaaagg ctgtgctgag gcttctagaa atcttaatgc atccttgagg  
gaggcagaga  
6961 tggggaaata gaaaaagaga gacacacaaa tgttctacag ttggagcgaa  
cagagagggg  
7021 cctggtgaga ttcaagggac aggcagggtgc acacagagac agagccagac  
ccagcggaga  
7081 gggaaggaag tgccccgacc tccggggctg agacctcaga gctggggcag  
gactgtgtcc

7141 ctaactgtcc accagtgtct ctgcctgtct cctgtgtct gttctcggg  
ttctctgtgc  
7201 catggtggct ctggctacct gtccatcagt gtctccattt ctgttcctcc  
ccctcaggg  
7261 gactctgggg gacccctggg gtgcagagga cagctccagg gcctcgtgtc  
ttggggaatg  
7321 gagcgtgcg ccctgcctgg ctaccccggt gtctacacca acctgtgcaa  
gtacagaagc  
7381 tggattgagg aaacgatgcg ggacaaatga tggctctcac ggtgggatgg  
acctcgtcag  
7441 ctgcccaggc cctcctctct ctactcagga cccaggagtc caggccccag  
cccctctcc  
7501 ctcagacca ggagtccagg cccccagccc ctctccctc agaccggga  
gtccaggccc  
7561 ccagcccctc ctccctcaga cccaggagtc caggccccag cccctctcc  
ctcagaccg  
7621 ggagtccagg cccccagccc ctctccctc agaccagga gtccaggccc  
cagtcctcc  
7681 tccctcagac ccaggagtcc agggccccag cccctctcc ctcagacca  
ggaatccagg  
7741 cccagcccct cctccctcag acccaggagc cccagtcccc cagccccctc  
tccttgagac  
7801 ccaggagtcc agggccagcc cctcctcct cagaccagga agccccagtc  
cccagcatcc  
7861 tgatctttac tccggtctg atctctcct tcccagagca gttgcttcag  
gcgttttctc  
7921 cccaccaagc cccaccctt gctgtgtcac catcactact caagaccgga  
ggcacagagg  
7981 gcaggagcac agaccctta aaccggcatt gtattccaaa gacgacaatt  
ttaaacacgc  
8041 ttagtgtctc taaaaaccga ataaataatg acaataaaaa tggaatcatc  
ctaaattgta  
8101 ttcattcatc catgtgttta ctttttattt tttagacaa ggtcttgctc  
agtctcctgg  
8161 tgaaatgctg taacgcaatc atagctcact gcaaccgtga cctcctgggc  
tccagtgatc  
8221 ctctacctc agcctccga gtagctggga ccacaggtgc ccgtaccat gccccgtac

SEQ.ID.NO. 66

KLK-L6 AA

MTQSQEDENKIIGGHTCTRSSQPWQAALLAGPRRRFLCGGALLSGQWVITAA  
HCGRPILQVALGKHNLRRWEATQQVLRVVRQVTHPNYNSRTHDNDLMLLQL  
QQPARIGRAVRPIEVTQACASPGTSCRVSQWGTISSPIARYPASLQCVNINISPD  
EVCQKAYPRITPGMVCAGVPQGGKDSCQGDSSGGLVCRGQLQGLVSWGM  
ERCALPGYPGVYTNLCKYRSWIEETMRDK

SEQ.ID.NO. 67

KLK-L6 AA

MFLLLTALQVLAIAMTQSQEDENKIIGGHTCTRSSQPWQAALLA  
GPRRRFLCGGALLSGQWVITAAHCGRPILQVALGKHNLRRWEATQQVLRVV  
RQVTHPNYNSRTHDNDLMLLQLQQPARIGRAVRPIEVTQACASPGTSCRVSQ  
WGTISSPIARYPASLQCVNINISPDEVQKAYPRITPGMVCAGVPQGGKDSC

QGDSSGGLVCRGQLQGLVSWGMERCALPGYPGVYTNLCKYRSWIEETMRD  
K

SEQ.ID.NO. 68

Figure 9

prostase

MATAGNPWGWFLGYLILGVAGSLVSGSCSQIINGEDCSPHSQPWQAALVME  
NELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVEASLSV  
RHPEYNRPLLANDLMLIKLDES VSESDTIRSIASQCPTAGNSCLVSGWGLLA  
NGRMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGGGHDQKDSCNGDSG  
GPLICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTEWIEK

SEQ.ID.NO. 69

Figure 9

EMSP

MATAGNPWGWFLGYLILGVAGSLVSGEMSPSCSQIINGEDCSPHSQPWQAAL  
VMENELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVEAS  
LSVRHPEYNRPLLANDLMLIKLDES VSESDTIRSIASQCPTAGNSCLVSGWG  
LLANGRMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGGGHDQKDSCNG  
DSGGPLICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTEWIEK 249

SEQ.ID.NO. 70

Figure 9

KLK-L2

MATARPPWMWVLCALITALLGVTEHVLANNDVSCDHPSNTVPSGSNQDLG  
AGAGEDARSDSSRIINGSDCDMHTQPWQAALLLRPNQLYCGAVLVHPQW  
LLTAAHCRKKVFRVRLGHYSLSVYESGQMFQGVKSIPHPGYSHPGHSNDL  
MLIKLNRRIRPTKDVRPINVSSHCP SAGTKCLVSGWGTTKSPQVHF PKVLQCL  
NISVLSQKRCEDAYPRQIDDTMFCAGDKAGRDSCQGDSSGPPVCNGLSLQGL  
VSWG DYPCARPNRPGVYTNLCKFTKWIQE

SEQ.ID.NO. 71

Figure 9

zyme

MKKLMVVLSLIAAAWAEQNKL VHGGPCDKTSHPYQAALYTS GHLLCGGV

LIHPLWVLTAAHCKKPNLQVFLGKHNLQRESSQEQQSSVVRAVIHPDYDAAS  
HDQDIMLLRLARPAKLSELIQPLPLERDCSANTTSCHILGWGKTADGDFPDTI  
QCAYIHLVSREECEHAYPGQITQNMLCAGDEKYGKDSCQGDSGGPLVCGDH  
LRGLVSWGNI PCGSKEKPGVYTNVCRYTNWIQK

SEQ.ID.NO. 72

Figure 9

neuropsin

MGRPRPRAAKTWMFLLLLGGAWAGHSRAQEDKVLGGHECQPHSQPWQAA  
LFQGGQQLLCGGVLVGGNWVLTAAHCKKPKYTVRLGDHSLQNKDGPEQEIPV  
VQSIPHPCYNSSDVEDHNHDLMLLQLRDQASLGSKVKPISLADHCTQPGQKC  
TVSGWGTVTSPRENFDTLNCAEVKIFPQKKCEDAYPGQITDGMVCAGSSKG  
ADTCQGDSGGPLVCDGALQGITSWGSDPCGRSDKPGVYTNICRYLDWIKKTL  
SPMRILQLILLALATGLVG

SEQ.ID.NO. 73

Figure 9

TLSP

GETRIIKGFECKPHSQPWQAALFEKTRLLCGATLIAPRWLLTAAHCLKPRYIV  
HLGQHNLQKEEGCEQTRTATESFPHPGFNNSLPNKDHRNDIMLVKMASPVSI  
TWA VRPLTLSSRCVTAGTSC LISGWGSTSSPQLRLPHTLR CANITIEHQK CEN  
AYPGNITDTMVCASVQEGGKDSCQGDSGGPLVCNQSLQGIISWGQDPCAIR  
KPGVYTKVCKYVDWIQE

SEQ.ID.NO. 74

Figure 9

PSA

MWVPVVFLTSLVTWIGAAPLILSRIVGGWECEKHSQPWQVLVASRGRAVCG  
GVLVHPQWVLTAAH CIRNKS VILLGRHSLFHPEDTGQVFQVSHSFPHPLYDM  
SLLKNRFLRPGDDSSHDLMLLRLSEPAELTDAVKVMDLPTQEPALGTTCYAS  
GWGSIEPEEFLTPKKLQCVDLHVISNDVCAQVHPQKVTKFMLCAGRWTGGK  
STCSGDSGGPLVCNGVLQGITSWGSEPCALPERPSLYTKVVHYRKWIKD

SEQ.ID.NO. 75

Figure 9

KLK2

MWDLVLSIALSVGCTGAVPLIQSRIVGGWECEKHSQPWQVAVYSHGWAHCG  
GVLVHPQWVLTAAHCLKKNSQVWLGRHNLFEPEDTGQRPVSHSFPHPLYN  
MSLLKHQSLRPDEDSSHDLMLLRLSEPAKITDVVKVLGLPTQEPALGTTTCYAS  
GWGSIEPEEFLRPRSLQCVSLHLLSNDMCARAYSEKVTEFMLCAGLWTGGKD  
TCGGDSGGPLVCNGVLQGITSWGPEPCALPEKPAVYTKVVHYRKWIKD

SEQ.ID.NO. 76

Figure 9

KLK1

MWFLVLCLALSLGGTGAAPPIQSRIVGGWECEQHSQPWQAALYHFSTFQCGG  
ILVHRQWVLTAAHCISDNYQLWLGRHNLFDDENTAQFVHVSESFPHPGFNMS  
LLENHTRQADEDYSHDLMLLRLTEPADTITDAVKVVELPTEEPEVGSTCLASG  
WGSIEPENFSFPDDLQCVDLKILPNDECKKAHVQKVTDFMLCVGHLEGGKDT  
CVGDSGGPLMCDGVLQGVTSWGYVPCGTPNKPSVAVRVLSYVKWIED

SEQ.ID.NO. 77

Figure 9

trypsinogen

MNPLLILTFVAAALAAPFDDDDKIVGGYNCEENSVPYQVSLNSGYHFCGGSLI  
NEQWVVSAGHCYKSRIQVRLGEHNIEVLEGNEQFINAAKIIRHPQYDRKTLNN  
DIMLIKLSRAVINARVSTISLPTAPPATGTKCLISGWGNTASSGADYPDELQC  
LDAPVLSQAKCEASYPGKITSNMFCVGFLEGGKDSQCQGDGGPVVCNGQLQ  
GVVSWG-DGCAQKNKPGVYTKVYNYVKWIKN

SEQ.ID.NO. 78

Figure 17, 27, 36, 43

PSA

MWVPVVFLTLSVTWIGAAPLILSRIVGGWECEKHSQPWQVLVASRGRAVCG  
GVLVHPQWVLTAAHCIRNKSIVLLGRHSLFHPEDTGQVFQVSHSFPHPLYDM  
SLLKNRFLRPGDDSSHDLMLLRLSEPAELTDAVKVMDLPTQEPALGTTTCYAS  
GWGSIEPEEFLTPKKLQCVDLHVISNDVCAQVHPQKVTKFMLCAGRWTGGK  
STCSGDSGGPLVCNGVLQGITSWGSEPCALPERPSLYTKVVHYRKWIKDTIVA  
NP

SEQ.ID.NO. 79

Figure 17, 27, 36, 43

KLK2

MWDLVLSIALSVGCTGAVPLIQSRIVGGWECEKHSQPWQVAVYSHGWAHCG  
GVLVHPQWVLTAAHCLKKNSQVWLGRHNLFEPEDTGQRVPVSHSFPHPLYN  
MSLLKHQSLRPDEDSSHDLMLLRLSEPAKITDVVKVLGLPTQEPALGTTCTYAS  
GWGSIEPEEFLRPRSLQCVSLHLLSNDMCARAYSEKVTEFMLCAGLWTGGKD  
TCGGDSGGPLVCNGVLQGITSWGPEPCALPEKPAVYTKVVHYRKWIKDTIAA  
NP

SEQ.ID.NO. 80

Figure 17, 27, 36, 43

KLK1

MWFLVLCALSLGGTGAAPPIQSRIVGGWECEQHSQPWQAALYHFSTFQCGG  
ILVHRQWVLTAAHCSIDNYQLWLGRHNLFDDENTAQFVHVSESFHPGFNMS  
LLENHTRQADEDYSHDLMLLRLTEPADTITDAVKVVELPTEEPEVGSTCLASG  
WGSIEPENFSFPDDLQCVDLKILPNDECKKAHVQKVTDFMLCVGHLEGGKDT  
CVGDSGGPLMCDGVLQGVTSWGYVPCGTPNKPSVAVRVLSYVKWIEDTIAE  
NS

SEQ.ID.NO. 81

Figure 17, 27, 36, 43

protease

MATAGNPWGWFLGYLILGVAGSLVSGSCSQIINGEDCSPHSQPWQAALVME  
NELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVEASLSV  
RHPEYNRPLLANDLMLIKLDESVS-  
ESDTIRISIASQCPTAGNSCLVSGWGLLANG—  
RMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFACAGGGHDQKDSCNGDSGGP  
LICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTEWIEKTVQAS

SEQ.ID.NO. 82

Figure 17, 27

trypsinogen

MNPLLILTFVAAALAAPFDDDDKIVGGYNCEENSVPYQVSLNSGYHFCGGS  
LI  
NEQWVVSAGHCYKSRIQVRLGEHNIEVLEGNEQFINAAKIIRHPQYDRKTLNN  
DIMLIKLSRAVINARVSTISLPTAPPATGTKCLISGWGNTASSGADYPDELQC  
LDAPVLSQAKCEASYPGKITSNMFCVGFLEGGKDSQCGDSGGPVVCNGQLQ  
GVVSWG-DGCAQKNKPGVYTKVYNVYVKWIKNTIAANS

SEQ.ID.NO. 83

Figure 17, 27, 36, 43

neuropsin

MGRPRPRAAKTWMFLLLLGGAWAGHSRAQEDKVLGGHECQPHSQPWQAA  
LFQGGQQLLCGGVLVGGNWVLTAAHCKKPKYTVRLGDHSLQNKDGPEQEIPV  
VQSIPHPCYNSSDVEDHNHDLMLLQLRDQASLGSKVKPISLADHCTQPGQKC  
TVSGWGTVTSPRENFPDTLNCAEVKIFPQKKCEDAYPGQITDGMVCAGSSKG  
ADTCQGDSGGPLVCDGALQGITSWGS DPCGRSDKPGVYTNICRYLDWIKKIIG  
SKG

SEQ.ID.NO. 84

Figure 17, 27, 36, 43

zyme

MKKLMVVLSLIAAAWAEQNKLVHGGPCDKTSHPYQAALYTSGHLLCGGV  
LIHPLWVLTAAHCKKPNLQVFLGKHNLQRRESSQEQQSSVVRAVIHPDYDAAS  
HDQDIMLLRLARPAKLSELIQPLPLERDCSANTTSCHILGWGKTADGDFPDTI  
QCAIYIHLVSREECEHAYPGQITQNMLCAGDEKYGKDSCQGDSGGPLVCGDH  
LRGLVSWGNIPCGSKEKPGVYTNVCRYTNWIQKTIQAK

SEQ.ID.NO. 85

Figure 27

EMSP

MATAGNPWGWFLGYLILGVAGSLVSGEMSPSCSQIINGEDCSPHSQPWQAAL  
VMENELFCSGVLVHPQWVLSAAHCFQNSYTIGLGLHSLEADQEPGSQMVEAS  
LSVRHPEYNRPLLANDLMLIKLDES VSESDTIRSISIASQCPTAGNSCLVSGWG  
LLANGRMPTVLQCVNVSVVSEEVCSKLYDPLYHPSMFCAGGGHDQKDSCNG  
DSGGPLICNGYLQGLVSFGKAPCGQVGVPGVYTNLCKFTEWIEKTVQAS

SEQ.ID.NO. 86

Figure 27

TLSP

MRI-LQLILLALATGLVGGETRIIKGFECKPHSQPWQAALFEKTRLLC  
GATLIAPRWLLTAAHCLKPRYIVHLGQHNLQKEEGCEQTRTATESFPHPGFNN  
SLPNKDHNRNDIMLVKMASPV SITWAVRPLTLSSRCVTAGTSC LISGWGSTSSP  
QLRLPHTLRCANITIEHQKCENAYPGNITDTMVCASVQEGGKDSCQGDSGGP  
LVCNQSLQGIISWGQDPCAITRKPGVYTKVCKYVDWIQETMKNN

SEQ.ID.NO. 87

Figure 27

HSCEE

MARSLLLPLQILLLSLALETAGEEAQGDKIIDGAPCARGSHPWQVALLSGNQL  
HCHSCCEGGVLVNERWVLTAAHCKMNEYTVHLGSDTLGDRRAQRIKASKSF  
RHPGYSTQTHVNDLMLVKLNSQARLSSMVKKVRLPSRCEPPGTTCTVSGWG  
TTTSPDVTFPDLMCVDVKLISPQDCTKVYKDLENSMLCAGIPDSKKNACNG  
DSGGPLVCRGTLQGLVS WGTFFCGQPNDPGVYTQVCKFTKWINDTMKKHR

SEQ.ID.NO. 88

Figure 27

NES1

MRAPHLHLSAASGARALAKLLPLLMAQLWAAEAALLPQNDTRLDPEAYGAP  
CARG SQPWQVSLFNGLSFH CAGVLVDQSWVLTAAHCGNKPLWARVGDDH  
LL-LLQG-EQLRRTT RSVVHPKYHQSGPI LPRRTDEHDLML LKLARPVV-  
PGPRVR ALQLPYR-CAQPGDQ CQVAGWGTTAARRVK YNKGLTCSSITLSP  
KECEVFYPGVVTNNM ICAGLDR-GQDPCQS DSGGPLVCDETLQGI LSWG-  
VYPCGSAQHPAVYTQICKYMSWINK VIRSN

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**